

Macromolecules made of subunits



(Lipids are not polymers)


Nucleic Acids

A D N A R N



@ 2001 Sinauer Associates, Inc.

Nucleotide



General structure of nucleotides






Figure 8-1a
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company

the numbering convention for the pentose ring




Numbering without prime


Nitrogenous bases





A nucleic acid strand:




(c) Polynucleotide











(a) Nucleotide components

Copyright @ Pearson Education, Inc., publishing as Benjamin Cummings.



Purines NH2 O C N C


Adenine Guanine A G



(c) Polynucleotide

(a) Nucleotide components

Copyright @ Pearson Education, Inc., publishing as Benjamin Cummings.



Purines

CH_{2 O} Ó-Phosphate group Pentose sugar (b) Nucleotide Guanine G Adenine носн_а он он Ribose (in RNA) (c) Polynucleotide (a) Nucleotide components


From Campbell

Nitrogenous base

Sugar-phosphate backbone **Bases** 5' phosphate end Thymine (T) Polarity of nucleic acid strands: Adenine (A) Cytosine (C) Phosphate **DNA** nucleotide 3' OH end Sugar (deoxyribose) Guanine (G)

Copyright @ Pearson Education, Inc., publishing as Benjamin Cummings.

DNA




Deoxyribo

Nucleic

Acid


DNA:

The 2 strands are complementary antiparallel




0.34 nm


Note the sizes




NUCLEIC ACIDS



NUCLEIC ACIDS (IN DNA)



NUCLEIC ACIDS (IN DNA)



Purine


Pyrimidine

Purine


Pyrimidine

ribose **Bases** phosphate Sugarphosphate backbone


NUCLEIC ACID STRUCTURE



A nucleotide contains a phosphate group, a sugar molecule, and a nitrogen-containing base.



Ribose carbons are numbered from 1' to 5'.




Pentose = 5-carbon sugar

Heptanucleotide


5' terminus

5' C A T G C A G 3'





3' terminus

NUCLEIC ACID STRUCTURE



A nucleotide contains a phosphate group, a sugar molecule, and a nitrogen-containing base.








Why DNA strands must be antiparallel?

Guanine and Cytosine can pair only when antiparallel

Parallel strands: G and C cannot form hydrogen bonds.









Copyright @ 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.


Campbell 8th



Copyright @ 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.


Campbell 8th






Copyright @ 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.


Campbell 8th





Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Bases pairing rule: purine with pyrimidine



Purine + purine: too wide

Pyrimidine + pyrimidine: too narrow

Purine + pyrimidine: width consistent with X-ray data

RIBONUCLEIC ACID (RNA)

RNA STRUCTURE


The sugar molecule

backbone contains

an extra oxygen.

in the RNA

There are three important structural differences between RNA and DNA.



RNA has only one sugar-phosphate backbone, while DNA has two.

Instead of thymine, RNA has a similar base called uracil.

Uracil

5′


RIBONUCLEIC ACID (RNA)

RNA STRUCTURE

 The sugar molecule in the RNA backbone contains an extra oxygen.


on carbon 2'

Instead of thymine,
 RNA has a similar base called uracil.



= oxygen

Differences between DNA and RNA



Sugar: desoxyribose ribose


Base: Thymine Uracil


Structure: double stranded Single stranded (mostly) (mostly)

RIBONUCLEIC ACID (RNA)

RNA FUNCTION

RNA acts as a middleman molecule. It takes instructions for production of a protein from DNA, moves them to another part of the cell, and directs the building of a protein.





(a) Replicating DNA Overview Replicating DNA strands Copy (replication) **REPLICATION** of each complementary DNA strand (b) DNA transcription Gene **TRANSCRIPTION** of a DNA template strand (c) Messenger RNA **Expression** Messenger RNA

(d) Polypeptide

Amino acid

Codon

traduction

TRANSLATION of

messenger RNA

Valine Leucine Serine Histidine Cysteine Glutamine Leucine Alanine

