Chapter 10:

The Structure and Function of DNA


covers


DNA replication

DNA transcription

mRNA translation

Chapter 11 covers the regulation of gene expression







Gene expression in an eukaryotic cell:

Eukaryotic cells are cells with a nucleus.

Translation takes place in the cytosol.

cytosol + organites = cytoplasme






Translation by ribosomes ocures in 3 steps :

• initiation

• elongation

termination

Genetic code




Note interesting facts:


- 61 sense codons + 3 STOP codons
- Tryptophan is encoded by only 1 codon
- arginine, leucine and serine are encoded by 6 codons
- synonymous codons : codons for the same amino acids

TRANSLATION of a messenger RNA growing polypeptide ribosome mARN (messenger) Scanning electron microscope Synthèse de protéines (MET, image colorisée). On distingue les ribosomes en bleu, l'ARN messager en rose et les protéines synthétisées en vert.

Translation




Ribosomes: made of rRNA and proteins




Eukaryotic ribosome


(prokaryotic ribosomes are similar but different enough to be specifically inhibited by antibiotics)

Eukaryotic ribosome











actual structure

tRNA (transfer RNA): carrier of amino acids






Attaching a specific amino acid to a tARN

Only 1 enzyme for a 2-step reaction

The accuracy of translation depends mainly on these enzymes. There are 20 enzymes, one for each amino acid.

Initiation of translation (in eukaryotes)

- → determines the **reading-frame**
- small subunit + initiator tRNA (with Met)
 - -> bind to cap
- 2. "scan" for first AUG = Met
- 3. large subunit joins






Figure 10.18 The initiation of translation





Figure 10.19


How do ribosomes and tRNA add amino acids during translation?

Translation: elongation





One elongation cycle:



Elongation: tRNAs arrive bound to EF-Tu





Figure 6-67 (part 1 of 7) Molecular Biology of the Cell (© Garland Science 2008)








Figure 6-67 (part 3 of 7) Molecular Biology of the Cell (© Garland Science 2008)



Note the position of the in-coming tRNA




Marshall et al. Annu. Rev. Biochem. 2008.77:177-203.

One elongation cycle: 3 steps


2. Formation of a peptide bound

What about the energy?

Is the reaction exergonic or endergonic?




One elongation cycle:



Translocation:


the model where the large subunit moves forward followed by movement of the small subunit is outdated.



Termination of translation

when ribosome reaches a stop-codon:

- 1. no complementary tRNA
- 2. release factors bind at A-site
- 3. detach and release polypeptide



Eukaryotic messengers are monocistronic.

If you engineer a bicistronic messenger only the upstream cistron is translated.



Now you should understand why only the first cistron is translated

Many ribosomes on a single mRNA

