


The usual case: balanced, biallelic expression




Sickled cells disease




Electrophoresis:





Linus Pauling, 1949: HbS has different charge.








Fig. 7.27 a-d. Zone electrophoresis diagram of hemoglobins at pH = 6.9. a Normal homozygote (AA); b patient with sickle cell anemia (SS); c sickle cell trait (AS); d mixture of equal parts of HbA and HbS. Arrow, starting point of electrophoresis. (From Pauling et al. 1949 [195])

Genetic dosage is important

Trisomy 21 47, XX +21





Each chromosome is used equally:


2 chromosomes : protein concentration = 100% 3 chromosomes : protein concentration = 150 %

Chromosome 21


47 millions base pairs


352 protein coding genes

In trisomy 21, 352 proteins are expressed about 150%.




Normal (disomic) cell:






Trisomic cell.


For genes located on the X chromosome women (X X) should produce twice as much of a protein compare to men (only 1 X)




Observations do not confirm the expectation.




Women and men have the same amount of G6PD because women and men have only 1 active X chromosome.



X-inactivation





Difference between calico and tortoiseshell


(b) (a) There is no melanocyte



Ewart BERTRAM

A Morphological Distinction between Neurones of the Male and Female, and the Behaviour of the Nucleolar Satellite during Accelerated Nucleoprotein Synthesis


Murray BARR (1908-1995)



XX Fig. 1 XY Fig. 2 Fig. 3 XX

In 1949, Murray Barr and Ewart Bertram identified a highly condensed structure in the interphase nuclei of somatic cells in female cats that was not found in male cats.

This structure became known as the **Barr body**.



(a) Nucleus with a Barr body

In 1960, Susumu Ohno correctly proposed that the Barr body is a highly condensed X chromosome.

Barr body

(inactivated X chromosome)






Figure 4-16a

Genetics: A Conceptual Approach, Third Edition
© 2009 W. H. Freeman and Company



Triple X (47, XXX)





2 Barr bodies (2 inactivated X)

One lady with this anomaly was Ewa Klobukowska, a 1964 Olympic sprint bronze medallist who was the first woman to fail the sex chromosome test during the 1967 European Cup. "One chromosome too many to be declared a woman for the purposes of athletic competition" was the statement, and she was obliged to return all her medals. She gave birth to a baby the following year.



→ 47,XXX women have normal children!!

Feminity test:

(competition)

Barr body

= inactivated X chromosome

Normal woman:




46, **X** X











Official document attesting gender.

Certificat de féminité pour les sportives :



L'athlète mentionnée ci-dessus a subi un examen médical approuvé et la **chromatine sexuelle** s'est révélée positive.

Complete androgen insensitivity




Maria PATINO

Tested female in 1983 in Helsinki; failed the gender test in 1985

46,XY female

Santhi Soundarajan



She had to give back her silver medal (800 m).


Is it fair ???



Gender test


(competitions)

Normal woman:










No Barr body

The opsines familly: 8 % of men are red / green color blind chromosome 7 Chromosome X gène S gène M gène L opsine sensible au bleu opsine sensible au vert opsine sensible au rouge


X-linked recessive




Recessivity is defined only in women.


Note: a few carriers may be mildly affected due to skewed X-inactivation.

Ishihara Test For Color Blindness






L'appariement n'est pas affecté par la méthylation :



5-methylcytosine base-pairs with guanine in exactly the same way as unmodified cytosine.


CpG Methylation



DNA replication (mitosis or meiosis)

Maintenance methylation

CpG Methylation



DNMT = **DN**a **M**ethyl **T**ransferase

DNA can be cut by restriction nucleases




Figure 10-4 Essential Cell Biology, 2/e. (© 2004 Garland Science)

Isoschizomeres

	cut	no cut
Hpa II recognizes	5'- C C G G – 3' 3'- G G C C – 5'	5'- C ^m C G G – 3' 3'- G G ^m C C – 5'
Msp I recognizes	5'- C C G G – 3' 3'- G G C C – 5'	5'- C ^m C G G – 3' 3'- G G ^m C C – 5'
	cut	cut

Hpa II and Msp I recognize the same restriction site: they are isoschizomeres.

However there is one important difference :

Hpa II is **sensitive** to DNA methylation Msp I is **insensitive** to DNA methylation

