

Figure 34-3 The amino acid sequence of calf thymus histone H4.


This 102-residue protein's 25 Arg and Lys residues are indicated in red. Pea seedling H4 differs from that of calf thymus by conservative changes at the two shaded residues: Val $60 \rightarrow$ Ile and Lys $77 \rightarrow$ Arg.

The underlined residues are subject to **post-translational modification**:

Ser 1 is invariably *N*-acetylated and may also be *O*-phosphorylated; Lys residues 5, 8, 12, and 16 may be *N*-acetylated; and Lys 20 may be mono- or di-*N*-methylated.

[After DeLange, R.J., Fambrough, D.M., Smith, E.L., and Bonner, J., J. Biol. Chem. 244, 5678 (1969).]

DNA replication in eukaryotic cells



POLY- MERASE	FUNCTION	PROOF- READING*
Pol α (alpha)	Contains <i>primase</i>Initiates DNA synthesis	-
Pol β (beta)	Repair	_
Pol δ (delta)	 Elongates Okazaki fragments of the lagging strand 	+
Pol ε (epsilon)	Elongates the leading strand	+
Pol γ (gamma)	Replicates mitochondrial DNA	+

Nucleosome structure






Figure 18-2bc Biological Science, 2/e



Condensed chromatin



Decondensed chromatin





Figure 18-4 Biological Science, 2/e


Heterochromatin forms localized clusters Heterochromatin Nucleolus Nucleus Heterochromatin Cytoplasm www.ergito.com

Nuclear content appears heterogenous :



b. Darkly-stained heterochromatin and lightly-stained euchromatin

Nuclear content appears hétérogenous :




Nucleolus: ribosomes assembly

b. Darkly-stained heterochromatin and lightly-stained euchromatin


Compaction de l'ADN au cours du cycle cellulaire

Chromatin and Condensed Chromosome Structure




interphase

metaphase (mitosis)



Cell cycle:



Mitotic phase Cell cycle: Mitosis G₂ Final growth and G₁ Centrioles G_1 activity before replicate mitosis DNA replication **Tutorial 9**

FIGURE 1–13 Sequence of events during the cell cycle. Immediately following mitosis (M) the cell enters a gap phase (G1) before a DNA https://wps.prenhall.com/wps/media/objects/1109/1135980/9_1_1_1_plain.html

Interphase

parts 3 and 4




Figure 2_16 Human Molecular Genetics, 3/e. (© Garland Science 2004)

on metaphase chromosomes or interphase nuclei.



Human metaphase chromosomes stained with propidium iodide

Alkaline DNA denaturation







Figure 1.16 Dénaturation du DNA par l'addition d'une base. L'addition d'une base à une solution de DNA en double hélice initialement à pH 7 provoque la séparation de la double hélice en simples brins. Le processus est à moitié complet un peu au-dessus de pH 9.







Karyotype



Expose to ultraviolet (UV) light.
 Take picture of fluorescent chromosomes.




Add hybridization probes labeled with fluorescent dye and wash away unhybridized probe.




FISH: Fluorescence in situ hybridization

Nucleus in **interphase**




2 probes

Chromosomes in **metaphase**




1 probe





Sonde centromérique







FIGURE 10-18 Fluorescent in situ hybridization (FISH) of human metaphase chromosomes. The probe, specific to centromeric DNA, produces a yellow fluorescence signal indicating hybridization. The red fluorescence is produced by propidium iodide counterstaining of chromosomal DNA.



Telomers:

