




One genome - many epigenomes







One genome - many epigenomes




Packaging of Chromatin inside the Nucleus



Epigenetic Mechanisms control Gene Activity




Epigenetic Mechanisms control Gene Activity




Repressed gene


H3K27me3





How is cellular identity established?




How is cellular identity established?














Histone modifications are gradually installed before zygotic genome activation















The repressive histone modification H3K27me3 is not remodelled in the early embryo










The repressive Histone Modification H3K27me3 is inherited











Higher Order Chromatin Structure is organized before ZGA





RNAi screen for chromatin factors involved establishing 3D genome organisation







HP1 binds H3K9me3 and compacts Heterochromatin






HP1 binds H3K9me3 and compacts Heterochromatin








HP1 recruitment triggers Establishment of Chromatin 3D Organization
















Understanding Human Brain Development


21 Days




Formation of Neural Tube





35-40 Days




Regionalization









Understanding Human Brain Development using Neural Organoids




Human Neural Organoids recapitulate embryonic Development









Reconstructing developmental transitions in human brain organoids




Organoids form different brain regions




Organoids form different brain regions




Mapping histone modifications in single cells with CUT&Tag



Prepare single cell Cut&Tag libraries using 10x Genomics





Binding and Tagmentation in situ inside the nucleus



Integrating RNA expression and histone modifications

Multimodal integration of scRNA and scCUT&Tag data in high-resolution clusters







Integrating RNA expression and histone modifications

Multimodal integration of scRNA


Taloncochales (Tal)

Multimodal integration of scRNA and scCUT&Tag data in high-resolution clusters






Building a model of gene regulation during brain development




Building a model of gene regulation during brain development




Check the regulatory pattern of any gene during early brain development!

EpiScape







Investigating Chromatin Regulation during Differentiation





Pseudotime alignment of RNA and histone modifications







Pseudotime alignment of RNA and histone modifications








Pseudotime alignment of RNA and histone modifications







Cell adhesion, signaling receptor activity, negative regulation of multicellular processes JARID2, FOXO1, CDH1


Epithelium development, central nervous system development, WNT and BMP signaling SKI, SFRP2, GPC3, LHX5, ZIC5, ZIC2, OTX2, IGF1R, IGF2BP1

Regionalization, telencephalon development FEZF2, PAX6, EMX2, NR2F2, ARX Cerebral cortex development POU3F3, POU3F2, LHX2, DCLK2, LHX2, ZEB1 Neuron development, axonogenesis, CRMP1, MEF2C, STMN2, NEUROD2, GRIA2, MAPT, PLXNA2, DPYSL5











Min-max scaled gene expression/activity 0 1


Zenk & Fleck et al. 2024, Nature Neuroscience


Chromatin accessibility precedes transcription at neuronal genes







Chromatin accessibility precedes transcription at neuronal genes










Functional role of Epigenetic Modifications in Organoid Development









Functional role of Epigenetic Modifications in Organoid Development










































EpiGN Lab



We use human neural organoid systems to study...

- 1. Brain development and cell differentiation
- 2. Molecular mechanisms of neurodevelopmental disorders
- 3. Maintenance of chromatin structure in postmitotic nuclei

