

ARTICLE

Received 5 Nov 2015 | Accepted 3 Jun 2016 | Published 3 Aug 2016

DOI: 10.1038/ncomms12149

OPEN

1

Whole-animal genome-wide RNAi screen identifies networks regulating male germline stem cells in *Drosophila*

Ying Liu^{1,*}, Qinglan Ge^{1,*}, Brian Chan¹, Hanhan Liu¹, Shree Ram Singh¹, Jacob Manley¹, Jae Lee¹, Ann Marie Weideman¹, Gerald Hou¹ & Steven X. Hou¹

Stem cells are regulated both intrinsically and externally, including by signals from the local environment and distant organs. To identify genes and pathways that regulate stem-cell fates in the whole organism, we perform a genome-wide transgenic RNAi screen through ubiquitous gene knockdowns, focusing on regulators of adult *Drosophila* testis germline stem cells (GSCs). Here we identify 530 genes that regulate GSC maintenance and differentiation. Of these, we further knock down 113 selected genes using cell-type-specific Gal4s and find that more than half were external regulators, that is, from the local microenvironment or more distal sources. Some genes, for example, *versatile* (*vers*), encoding a heterochromatin protein, regulates GSC fates differentially in different cell types and through multiple pathways. We also find that mitosis/cytokinesis proteins are especially important for male GSC maintenance. Our findings provide valuable insights and resources for studying stem cell regulation at the organismal level.

¹ Basic Research Laboratory, National Cancer Institute at Frederick, National Institutes of Health, 1050 Boyles Street, Building 560, Room 12-70, Frederick, Maryland 21702, USA. * These authors contributed equally to this work. Correspondence and requests for materials should be addressed to S.X.H. (email: hous@mail.nih.gov).

tem cells are crucial components of many animal tissues and organs. Germline stem cells (GSCs) are unique because of transmitting genomic information to future generations. GSCs in *Drosophila* testis provide an excellent opportunity to study the defining mechanisms of stem cells at the cellular and molecular levels (reviewed in refs 1,2). GSCs in the Drosophila testis reside at the apical tip of the testis that maintain spermatogenesis^{1,2}. Each GSC is enclosed by two somatic cvst stem cells (CvSCs). A recent study suggests that CvSCs encystment promotes GSCs abscission during GSC division due to inhibition of cytokinesis³. Both the GSCs and CySCs are attached to a cluster of postmitotic somatic cells called the hub⁴, via cadherin-mediated cell adhesion^{5,6}. The somatic hub serves as a niche, which expresses the signalling ligand for the Drosophila Janus kinase-signal transducer and activator of transcription (JAK-STAT) pathway, Unpaired (Upd). Upd instructs the JAK-STAT pathway in neighbouring GSCs and CySCs to regulate their self-renewal. Besides the JAK-STAT pathway, there are other regulatory pathways that also control the fate of GSCs and CySCs at the testis niche^{1,3,7-20}.

In CySCs, JAK-STAT signalling and its putative targets (such as such as zinc-finger homeodomain-1 (zfh-1) and chinmo) both act intrinsically to regulate CySC self-renewal and maintenance, and non-autonomously regulate the self-renewal of adjacent GSCs^{21,22}. Epidermal growth factor receptor (EGFR)-Ras/Raf signalling also acts both intrinsically to regulate CySCs and non-autonomously to regulate adjacent GSCs, but has the opposite effects of JAK-STAT signalling^{23–25}. In addition, a Polycomb group component, Enhancer of Zeste, acts intrinsically in CySCs to control CySC activity and identity, and non-autonomously to prevent adjacent GSC ectopic self-renewal and conversion to a somatic cell identity²⁶. However, it is unclear how these pathways are interconnected in CySCs to both control CySCs intrinsically and regulate GSCs non-autonomously.

In GSCs, the mother (old) centrosome always anchored close to the niche during GSC division, whereas the daughter centrosome moves to the reverse side of the cell, congregating a mitotic spindle perpendicular towards the hub^{27,28}. GSCs actively monitor centrosome orientation through a centrosome orientation checkpoint $(COC)^{8,27,29}$. The centrosome component Centrosomin and the cell polarity kinase Par-1 are essential components of the COC. The polarity protein Bazooka (Baz) provides a platform for the correct centrosome orientation and the COC monitors centrosome orientation by controlling the Baz-centrosome association. GSCs with COC defects enter mitosis without the correct centrosome orientation, leading to misoriented spindles and symmetric stem cell divisions 30,31. Further, the mother centriole nucleates excess microtubules compared with the daughter centriole that may promote asymmetric Apc2 (Adenomatosis polyposis coli 2) delivery to the cortex where GCSs contact the hub²⁸. In addition, at the cortex, Apc2 and E-cadherin together anchor the spindles of mitotic GSCs perpendicular to the hub³¹. As a result, only one daughter cell is directly connected to the hub and receives JAK-STAT signal to sustain its stem cell identity, whereas the other daughter cell, at the opposite end of the mitotic spindle, experiences a much weaker signal and begins differentiation. Recently, we found that nucleoporin and spindle matrix protein Tpr/Mtor (megator) regulate GSC asymmetric division and maintenance¹⁶. Loss of Mtor function affects expression and localization of Apc2 and E-cadherin. Further, we found that Mtor is essential for proper centrosome orientation, mitotic spindle formation and chromosome segregation. Our results suggest that nuclear matrix-SAC (spindle assembly checkpoint) axis controls maintenance and asymmetric division of GSC through

the Mtor-Mps1 (monopolar spindle 1)/Mad2 (mitotic arrest deficient 2) pathway¹⁶.

Recent developments in genome-wide RNA interference (RNAi) techniques in *Drosophila* have enabled the knockdown of nearly complete sets of genes involved in cellular processes in living animals 32–34. In addition, genome-wide RNAi screens have been performed to identify regulatory networks in several types of stem cells 35–38. However, stem cells are regulated not only intrinsically but also by extracellular cues from the local environment 39; these previous screens focused only on identifying intrinsic regulators. Furthermore, signals from distant organs also regulate stem cell/progenitor maintenance 40,41 and organ–organ communications are very important in regulating organismal growth and ageing 42–44.

The fruit fly *Drosophila* uniquely enables the systematic study of stem cell biology at the organismal level. To comprehensively identify genes and pathways that regulate GSC fates from different cell types in the whole organism, we perform a genome-wide transgenic RNAi screen through ubiquitous knockdowns of genes in adult *Drosophila* and then examine male GSC defects. Here we identify 530 genes whose RNAi-mediated knockdown affects stem cell maintenance and differentiation. Of these, we further knock down selected genes using cell-typespecific Gal4s and find that more than half are external regulators of GSC fate that originate from either the local microenvironment or distant organs. Moreover, we identify genes that can differentially regulate GSC fates from different cell types and through multiple pathways. Our data provide valuable insight and a useful resource for studying stem cell regulation at the organismal level.

Results

Developing the high-throughput screen. To systematically analyse the function of individual genes in the male GSC, we screened the existing Vienna *Drosophila* RNAi Center (VDRC) and the Bloomington *Drosophila* Stock Center (BDSC) collection of long double-stranded RNA (dsRNA) and short small hairpin RNA (shRNA) lines. The RNAi methodology has definite restrictions^{32–34}. First, the P-element-based UAS-hairpin constructs incorporate haphazardly into the genome and the level of hairpin expression is influenced by its chromosomal location. Second, the RNA level is reduced only to a variable degree by the RNAi-mediated knockdown that, in some cases, results in negligible effect. Third, null mutations of a large number of non-essential genes do not cause a phenotype (FlyBase).

To reduce the overall false-negative rate and conduct an efficient screen, we first performed a pilot experiment in which we selected 1,000 RNAi lines at random. Each of these lines was crossed in duplicate to actin GAL4 driver fly line (*Act5-Gal4*) and to *Act5-Gal4*, *UAS-GFP/+*; *tub-Gal80*^{ts}/+ drivers. The progeny from the cross with *Act5-Gal4* were screened for lethality and any visible adult phenotype. Those from the cross with *Act5-Gal4*, *UAS-GFP/+*; *tub-Gal80*^{ts}/+ were scored for GSC phenotypes. We found that 90.4% of the RNAi lines with GSC phenotypes were lethal in the cross with *Act5-Gal4*. Thus, in the following screen, we first crossed all the RNAi lines with *Act5-Gal4* to test for lethality and then crossed only the lethal lines with *Act5-Gal4*, *UAS-GFP/+*; *tub-Gal80*^{ts}/+ to screen for GSC phenotypes (Fig. 1a).

Genome-wide RNAi screen for male GSC using *Act-Gal4*. To identify candidate genes involved in male GSC regulation, we performed a systematic RNAi screen *in vivo*. We ubiquitously expressed *UAS-RNAi* in adult *Drosophila* using the *Act5-Gal4*, *UAS-GFP/*+;tub-Gal80^{ts}/+ driver⁴⁵. Crossing of virgin females




Figure 1 | Transgenic RNAi screen. (a) Workflow of the GSC RNAi screen. **(b)** Pie chart summarizing the screen results. Fly lines shown in the table are those that gave a lethal phenotype when crossed with *Act-Gal4*. **(c)** Pie chart summarizing phenotype observed in all the 720 lines. *Act*^{1s} > *RNAi* male flies were dissected after 7 days at 29 °C. Their testes were stained with antibodies (vasa, 1B1 and Arm) and analysed by confocal microscopy. The phenotypes were divided into two major categories (GSC decrease and increase). Percentage of fly lines showing no phenotypes are also shown in the chart. **(d)** Confidence of 530 genes identified from the screen. High-confidence genes were identified by two or more independent RNAi lines. Medium-high-confidence genes were identified by one RNAi line, but they shared a complex with high-confidence hits. Medium-low-confidence genes were identified by one RNAi line, but they shared a complex with other low-confidence hits. Low-confidence hits were identified by one RNAi only.

(Act5-GAL4, UAS-GFP/+;tub-Gal80^{ts}/+) with two to three males from UAS-RNAi fly lines were set at 18 °C. Male adult flies from F1 generation were shifted to 29 °C, to induce RNAi expression. After 7 days, testes of F1 males were dissected and examined for phenotypes, and only abnormal testes were stained with antibodies and examined under confocal microscopy (Fig. 1a).

GSCs in the adult Drosophila testis are regulated both intrinsically and externally by signals from the hub and CySCs (reviewed in ref. 1). In addition, trachea-derived decapentaplegic (Dpp) regulates stem cells in the adult Drosophila midgut⁴⁰ and olfactory receptor neurons regulate blood progenitor maintenance by controlling γ -aminobutyric acid secretion⁴¹. Furthermore, organ-organ communication plays important roles in regulating organism growth and ageing 42-44,46. Thus, signals from both the local environment and distant organs could be important for GSC regulation and much of the complexity of stem cell regulation would be lost if we limited our screen to one cell type. Therefore, we first performed the genome-wide transgenic RNAi screen by the ubiquitous knockdown of genes with Act-Gal4 and then characterized the functions of selected genes in individual cells with cell-type-specific Gal4 (such as germ cells specific driver, Nanos (Nos)-Gal4 or CySC and early cyst-cell driver (c587-Gal4). This is similar to first performing a classical ethyl methanesulfonate- or P-element-mediated screen and then characterizing the functions of selected genes in individual cells by mosaic clonal analysis.

In total, we screened 19,979 transgenic lines of either dsRNA or shRNA from both the VDRC and the BDSC (Fig. 1b and Supplementary Data 1), representing 13,019 of the 14,139 protein-coding genes (92%) in release 5.7 of the *Drosophila* genome⁴⁷. Among the 19,979 transgenic lines tested, 9,106

(45.6%) lines, corresponding to 6,873 genes, were lethal, once expressed by the *Act5-Gal4* driver.

We then expressed the 9,106 transgenic RNAi lines with *Act5-Gal4*, *UAS-GFP/+*; *tub-Gal80*^{ts}/+ (Fig. 1a) and analysed 5–10 flies for each line for GSC phenotypes (Fig. 1c). A total of 720 promising lines scored from the first-round screen, corresponding to 530 genes (Fig. 1b,d and Supplementary Data 2), were repeatedly screened and stained with molecular markers to confirm the phenotypes (Figs 2–5).

Through a BLAST search for all of the other available lines in the VDRC and BDSC, which were generated from independent or mini-overlapping sequences for all 530 genes, we found only 319 dsRNA lines (319 genes) have other available lines in the VDRC and BDSC. Among these 319 genes, the phenotypes of 150 genes were confirmed by two or more independent lines; the other 169 genes had either no other independent line or other independent line but lacked a phenotype, possibly due to low knockdown efficiency (Supplementary Data 1). The phenotypes of a number of genes, including the following four, were confirmed using only two independent lines, even if more than two independent lines were obtained: CG1098/Madm (MLF1-adaptor molecule)— BL316644 (GSC loss)/V27346 (GSC loss)/BL42529 (no phenotype), CG7467/osa—BL31266 (GSC loss)/V7810 (GSC loss)/ BL38258 (no phenotype), CG8553/SelD (Selenide)—V35959 (GSC loss)/BL29553 (GSC loss)/V100820 (no phenotype) and CG32183/Ccn-V38208 (GSC loss)/BL31715 (GSC loss)/ V101518 (no phenotype). This result showed that some independent lines had low knockdown efficiency. Among 211 shRNA lines, the phenotypes of 10 were confirmed in two or more independent lines. We did not further confirm the other 201 shRNA lines, because lines generated from shRNAs usually have better efficiency and fewer off-targets than dsRNA lines³⁴.




Figure 2 | Representative RNAi screen phenotypes using Act^{ts}-Gal4. (a) Diagram of GSC lineage and marker expression. GSCs and GBs have round spectrosomes (green dots) and spermatogonia contain branched fusomes (green branched lines). GSCs are enclosed by two CySCs (pink). (b) Wild-type (WT) control. (c) Knockdown of *tum* resulted in GSC loss and premature differentiation into spermatogonial cells (white arrow). (d) Knockdown of *Nedd8* resulted in GSC loss and death of differentiated cells. (e) Knockdown of *bab2* (*bric a brac 2*) resulted in GSC loss and GSCs that directly differentiated into spermatogonial cells (yellow dotted line) and spermatocytes (light blue arrows). (f) Knockdown of *Taf8* resulted in GSC differentiation defects. (g) Knockdown of *Cka* (*Connector of kinase to AP-1*) resulted in GSC overproliferation at the expense of terminal differentiation. (h) Knockdown of *Vein* (*Vn*) resulted in GSC differentiation defects. Testes of the indicated flies were dissected, stained with antibodies to Vasa (red, marks all germ cells including GSCs), Arm (green, hub cells at the apex), 1B1 (green, marks round spectrosomes and branched fusomes) and DAPI (blue, marks the nuclei), and analysed by confocal microscopy. Asterisks indicate hub cells. White arrows near hub cells indicate GSC and green arrows indicate GBs (in b,d,e), and light blue arrows (b,c,e,h) indicate spermatocytes. White dotted lines indicate GSCs (near hub in a) and GSC-like cells (in f-h). Scale bars, 10 μm.

Quality evaluation. Six lines of evidence suggest that our screen identified GSC regulators with high confidence. First, it is known that the JAK-STAT and EGFR-Ras/Raf signal transduction pathways are major pathways that regulate male GSCs, often with opposite effects²¹⁻²⁵. We identified four positive regulators in the JAK-STAT pathway whose knockdown resulted in GSC loss, nine positive regulators in the EGFR-Ras/Raf pathway whose knockdown caused GSC differentiation defects and four other components of the EGFR pathway, whose knockdown resulted in GSC loss (Supplementary Data 2). Second, many of the other genes found in our screen were previously shown to regulate GSC functions (Supplementary Data 2). Third, many of the identified hits (385 genes), whose products are components of protein complexes, showed high phenotypic similarity (Fig. 1c and Supplementary Data 2 and 3). Fourth, of 530 genes identified, 319 were positive targets from dsRNA lines, among which we verified 150 genes by at least two or more independent lines (Supplementary Data 1 and 2); the other 211 genes were positive targets from shRNA lines, which are known to have good efficiency and few off-targets³⁴. We also verified some of the genes identified in the RNAi screen by mutant clone analysis (Figs 3p,q and 4d,g). Furthermore, many of the GSC-specific genes were lethal in the primary screen, but did not show GSC phenotypes. Fifth, among the remaining 141 low-confidence genes, 78 (78/141 = 55%) were knocked down with shRNA lines and 41 (41/141 = 29%) were identified in at least 1 of 3 previous stem cell RNAi screens (neuroblast³⁵, female GSC³⁶ and intestinal stem cell (ISC)³⁷; Supplementary Data 4). Finally, we showed efficient knockdown of a select set of genes by quantitative PCR analysis (Supplementary Data 5).

Genes whose knockdown resulted in GSC loss. In the wild-type testes, five to nine GSCs reside at the apical niche (Fig. 2a,b). RNAi-mediated knockdowns of genes required for GSC self-renewal and maintenance resulted in GSC loss. We identified 300 such genes (Figs 1c and 2c–e, and Supplementary Data 2).

JAK-STAT pathway. The JAK-STAT signalling pathway was the first one reported to regulate stem cell maintenance in the *Drosophila* testis^{24,48}. In the testis, the ligand Upd is expressed in hub cells and activates JAK-STAT signalling in adjacent stem cells for the maintenance of both GSCs and CySCs. When JAK-STAT signalling is depleted from testis cells, both GSCs and CySCs are completely lost. In contrast, when the signalling is amplified by overexpressing upd throughout the testis apex or by overexpressing a constitutively activated form of JAK kinase, hopscotch tumorous lethal (hop^{Tum-l}) in somatic cells, the testes are filled with self-renewing GSCs interspersed among the CySCs^{1,15,21}. In this screen, we found that knockdown of the receptor gene dome (domeless), the Drosophila stat gene stat92E and two new positive regulators in the pathway, CG6946/glo (glorund)⁴⁹ and CG31716/Cnot4 (Cnot 4 homologue)⁵⁰, resulted in GSC loss. Surprisingly, the knockdown of Drosophila PIAS (protein inhibitor of activated STAT-1) gene (dPIAS) also resulted in GSC loss. dPIAS was initially identified as a negative regulator of the JAK-STAT pathway and later found in the heterochromatin protein complex involved in epigenetic modification^{51–53}. Thus, dPIAS may regulate GSCs through a JAK-STAT-independent pathway.




Figure 3 | Cell-type-specific knockdowns of mitotic/cytokinesis genes. (a) A schematic diagram showing expression pattern of Nos-Gal4 in adult testis. Knockdown of Smc2 (b), APC10 (c), Cenp-C (Centromeric protein-C) (d) and tum (e) in germ cells (Nos-Gal4) resulted in dramatic GSC loss. (f) A schematic diagram showing expression pattern of c587-Gal4 in adult testis. Knockdown of Smc2 (g), Cenp-C (i) and tum (j) in CySC (c587-Gal4) resulted in GSC differentiation defects (white dotted lines, g,i,j); however, knockdown of APC10 (h) in CySCs resulted in GSC loss. (k) A schematic diagram showing expression pattern of Bam-Gal4 in adult testis. Knockdown of Smc2 (l) and APC10 (m) Cenp-C (n) and tum (o) in spermatogonial cells (Bam-Gal4) resulted in normal (l,m) or polyploid (n,o, yellow arrows) cells. (p,q) FRT^{42D}-SMC2^{f06842} mosaic clones of GSCs 2 days (p, white dotted lines with white arrows, β-galactosidase (green) negative) after clonal induction (ACI). Armnegative GSC clones were detected in 4 out of 8 testes 2 days ACI but in 0 out of 20 testes 7 days ACI. Therefore, SMC2 is intrinsically required for GSC maintenance. Testes of the indicated flies were dissected, stained with antibodies to Vasa (red, marks all germ cells including GSCs), Arm (green, hub cells at the apex), 1B1 (green, marks round spectrosomes and branched fusomes) and DAPI (blue, marks the nuclei) in b-o, and analysed by confocal microscopy. White arrows near hub cells indicate GSC and green arrows indicate GBs (in l-o). Asterisks indicate hub cells. The flies were cultured at 29 °C for 7 days in b-o. Scale bars, 10 μm.

Mitosis/cytokinesis genes. In mammalian cells after anaphase onset, the mitotic spindle is completely reorganized into an array of interdigitating and antiparallel microtubules, known as the central spindle. Many signalling and cytoskeletal proteins cooperate to assemble and organize the central spindle. These include the microtubule-associated protein regulatory of cytokinesis 1 (PRC1, Drosophila Fascetto/Feo), the chromosomal passenger complex and several kinesin-like motors: KIF4A (Drosophila kinesin-like protein at 3A/Klp3A), KIF14 (Drosophila Nebbish/ Neb), KIF20A (Drosophila Subito) and KIF23 (Drosophila Pavarotti/Pav). KIF4A transports PRC1 to the spindle midzone, KIF14 interacts with PRC1 and KIF23 is important for bundling central spindle microtubules to form a protein complex known as centralspindlin. Centralspindlin is a conserved heterotetramer composed of KIF23/Pav and RacGAP1/MgcRacGAP (Drosophila Tumbleweed/Tum) dimers. The central spindle plays a key role in cleavage furrow formation. In addition, the formation of an active

zone of RhoA at the equatorial cortex both stimulates profilinmediated actin polymerization by binding the diaphanous (Dia) formin-homology proteins and activates Rho-associated kinase, which phosphorylates the myosin regulatory light chain (*Drosophila* Spaghetti squash/Sqh), to promote myosin contractility. These events together trigger actomyosin filament assembly at the cleavage furrow and contractile ring formation (reviewed in refs 54–56).

In this screen, we identified 69 genes that had been either described as mitotic genes in FlyBase or identified in two previous screens for mitotic genes^{57,58}. Knockdown of these 69 genes resulted in GSC loss (Figs 2 and 3, and Supplementary Data 6 and 2). This group of genes included *sqh*, *pav*, *tum*, *dia*, *feo*, *pebble* (*pbl*) and *neb*. As knockdowns of these cytokinesis genes resulted in GSC loss, these factors are probably important for GSC maintenance. We further tested 16 of these genes in spermatogonial cells (using *Bag of marbles* (*Bam*)-*Gal4*) and




Figure 4 | TCTP differentially regulate GSCs from different cells. (a) Knockdown of *Tctp* using *Act-Gal4* resulted in GSC differentiation defects. (b) Knockdown of *Tctp* using *Nos-Gal4* resulted in GSC loss. (c) Knockdown of *Tctp* using *c587-Gal4* resulted in GSC differentiation defects. *FRT*⁸²⁸-*piM* mosaic clones of GSCs 2 days (d) and 7 days (e) ACI. *FRT*⁸²⁸-*Tctp*^{h5p} mosaic clones of GSCs 2 days (f) and 7 days (g) ACI. In control flies, Arm-lacZ-negative GSC clones were detected in 33 out of 37 testes 2 days ACI and in 51 out of 64 testes 7 days ACI; in *Tctp*^{h5p} flies, Arm-lacZ-negative GSC clones were detected in 37 out of 53 testes 2 days ACI and in 8 out of 63 testes 7 days ACI. Thus, *Tctp* is intrinsically required for GSC maintenance. Testes of the indicated flies were dissected, stained with antibodies to Vasa (red, germ cells including GSCS), β-galactosidase (green) and DAPI (blue, marks the nuclei) in d-g, and Vasa (red), Arm (green, hub cells at the apex), 1B1 (green, marks round spectrosomes and branched fusomes) and DAPI (blue, marks the nuclei) in a-c, and analysed by confocal microscopy. The flies were cultured at 29 °C for 7 days in a-c. White arrows near hub cells indicate a GSC clone (d-f) and yellow dotted lines in d-g indicate GSC clones. White dotted lines in a,c indicate GSC-like cells. A yellow arrow in b indicates a remaining germ cell. Asterisks indicate hub cells. Scale bars, 10 μm.

found that the knockdown of 14 genes (14/16=88%) with *Bam-Gal4* caused polyploidy (Supplementary Data 6 and Fig. 3n,o,q), a phenotype usually associated with cytokinesis defects⁵⁶, indicating that most of the 69 mitotic genes regulate cytokinesis. Knockdown of these genes in GSCs may block GSC cytokinesis, causing polyploidy, followed by elimination of the defective GSCs through cell death.

We further crossed 32 of the 69 genes to cell-type-specific Gal4s and found that 25 (25/32 = 78%) resulted in strong GSC loss with Nos-Gal4 (GSCs and early germline cysts) and various phenotypes with c587-Gal4 (CySCs and early cyst cells) (Supplementary Data 6 and Fig. 3). We also generated SMC2 (Structural maintenance of chromosomes 2) mutant mosaic clones of GSCs and found that SMC2 is required for GSC maintenance (Fig. 3p,q). These data together suggest that most of this group of genes intrinsically regulate GSC maintenance. We also identified 27 mitotic genes whose knockdowns resulted in GSC differentiation defects (Supplementary Data 2). These 27 genes are mostly ribosomal proteins or translational factors identified in the two previous screens for mitotic genes^{57,58}; they may indirectly regulate mitosis by controlling the expression of key mitotic factors, as previously suggested or they may regulate GSC differentiation through mitosis-independent functions.

In a genome-wide RNAi screen focusing on *Drosophila* midgut ISCs³⁷, we previously identified eight genes that regulate the mitotic cell cycle or mitotic cytokinesis. However, knockdown of these genes blocked mitotic cell division and disrupted the diploid status of ISCs, resulting in ISCs with excessive cell growth and large polyploid nuclei. It will be interesting to examine why disruption of the mitotic cell cycle resulted in GSC loss but excessive growth and large nuclei in ISCs. Notably, there were many more mitotic genes identified in the male GSC screen than in screens of ISCs, female GSCs and neuroblast^{35–37}, suggesting that the mitotic genes play a more important role in male GSC regulation.

Dlg-Scrib-EGFR and Toll pathways. Our screen identified two cell polarity complex components discs large (dlg), scribbled (scrib), four EGFR signalling pathway components (krn/Keren, Lk6/Lk6 kinase, Mkp3/Mitogen-activated protein kinase phosphatase 3 and vav) and two Toll signalling pathway components (Nedd8/Neural precursor cell expressed, developmentally downregulated 8 and dl/dorsal). Knockdowns of these eight genes all resulted in GSC loss (Supplementary Data 2). The tumour suppressor genes dlg and scrib encode septate junction proteins with functions in epithelial cell polarity. Dlg is required for GSC




Figure 5 | Vers interacts with EGFR and Dpp to regulate GSC and CySC. (a,b) Knockdown of vers in CySC lineage c587-Gal4) caused GSC differentiation defects. (c) Wild-type (control) testis. (d) Knockdown of vers in CySC lineage (c587-Gal4) caused CySC expansion. (e) Wild-type (control) testis. (f) Knockdown of vers in CySC lineage (c587-Gal4) activates both EGFR and Dpp signalling. Testes of the indicated flies were dissected, stained with antibodies to Vasa (red), Arm (green, hub cells at the apex) and 1B1 (green, marks round spectrosomes and branched fusomes) in a,b, Vasa (red) and Zfh1 (green, marks CySCs) in c,d, Smad3 (red) and Cic (green) in e,f, and DAPI (blue, marks the nuclei), and analysed by confocal microscopy. White dotted lines in a,b indicate GSC-like cells. White arrow near hub cells indicates a GSC and green arrow indicates a CySC (c). Yellow dotted lines in f indicate Smad3-positive cells. Asterisks indicate hub cells. The flies were cultured at 29 °C for 7 days. Scale bars, 10 μm.

and CySC maintenance by establishing and maintaining the male stem cell niche in somatic cells^{59,60}. Scrib and the cell polarity complex may participate in the process involving Dlg. As described in a later section, EGFR signalling mainly functions in somatic cells to regulate GSC abscission³. The four EGFR-signalling-related components may affect GSC maintenance by negatively regulating EGFR signalling. The Toll pathway's function in GSC regulation is novel, therefore; it will be interesting to examine how this pathway regulates GSC self-renewal or maintenance.

In addition to the genes described above, we also identified 218 other genes (Supplementary Data 2) whose RNAi-mediated knockdowns resulted in GSC loss. These 218 genes include several genes that function in nucleosome, nucleoporin, chromatin binding and epigenetic modification (Supplementary Data 2), but most of the remaining genes' functions in GSC regulation are new. These genes provide a rich resource for future studies on GSC self-renewal and maintenance.

Genes regulate GSC expansion or differentiation defects. RNAi-mediated knockdowns of genes required for GSC differentiation or of inhibitors of GSC self-renewal resulted in GSC differentiation defects or expansion. For simplicity, we will call them differentiation defects from now on. We identified 230 genes whose knockdown resulted in GSC differentiation defects (Figs 1c and 2f-h, and Supplementary Data 2).

The EGFR-Raf-MAPK and Tor pathways. The EGFR-Raf-MAPK pathway was the first pathway reported to regulate GSC

differentiation^{24,25}. The germ cells associated with somatic cells mutant for the Raf and Egfr genes fail to differentiate and accumulate as early-stage germ cells, suggesting that the EGFR-Raf-MAPK pathway in somatic cells is required for GSC differentiation 24,25. In addition, germ cells are known to release the EGFR ligand Spitz, to induce the somatic cell EGFR-Rac pathway, which regulates the germ cell enclosure, by somatic cells⁶¹. In somatic cells, Rac1 is positively regulated by the guanine nucleotide exchange factor Vav and negatively regulated by the small GTPase Rho1 (ref. 61). The encystment of GSCs by CySCs promotes abscission during GSC division after a regulated block in cytokinesis³. In this screen, we identified nine positive regulators (CG9375/Ras85D (Ras oncogene at 85D), CG10491/vn (vein), CG10334/spi, CG15793/Dsor1 (Downstream of raf1), CG1044/dos (daughter of sevenless), CG4063/ebi, CG5053/ RASSF8, CG3403/Mob4 and CG10754) in the EGFR-Raf-MAPK pathway whose knockdown resulted in GSC differentiation defects.

Our screen also identified two components (*raptor* and *Tor*) in the Tor (target of rapamycin) signalling pathway. Knockdowns of these two genes resulted in GSC expansion or differentiation defects (Supplementary Data 2). The Tor pathway's function in GSC regulation is novel. It will be interesting to examine how this pathway regulates differentiation.

Other genes involved in GSC differentiation. Besides the genes in the EGFR and Tor signalling pathways, and the 27 mitotic genes described above, we also identified 5 microtubule motor complex components, 9 mediator complex components,

11 components involved in vesicle transport/secretion, 5 nucleoporin components and 162 other genes. These genes will be a rich resource for future studies on GSC differentiation.

Signal from local niche and distant organs regulate GSC fate. To investigate the cell-type specificity of the above-identified genes, we knocked down 113 genes by crossing their corresponding RNAis with both Nos-Gal4 and c587-Gal4 (Supplementary Data 6). Among the 113 lines that resulted in strong abnormal GSC phenotypes with Act-Gal4, only 43 lines (38%) had abnormal and similar GSC phenotypes with Nos-Gal4 (Supplementary Data 6), suggesting that more than half of the genes (62%) identified in the screen regulate GSC fates nonautonomously. Of the 113 lines, 74 (65%) had abnormal (either gain or loss) GSC phenotypes with c587-Gal4, suggesting that CySCs or early cyst cells are important in regulating GSC fates. In addition, 73 (65%) of the 113 lines had abnormal GSC phenotypes with Nos-Gal4 and c587-Gal4, similar to their phenotypes with Act-Gal4, suggesting that 65% of the genes regulate GSC fates intrinsically or externally from CySCs or local early cyst cells, and the other 35% regulate GSCs from distant organs. We also found that 28 of the 113 genes had no phenotypes with both Nos-Gal4 and c587-Gal4, suggesting that these genes regulate GSCs outside of GSCs and CySCs. We selectively crossed the RNAi lines of these 28 genes with Upd-Gal4 (hub cells) and Elav-Gal4 (pan neurons). None of the 10 lines with Upd-Gal4 had GSC defects, whereas 9 of the 23 lines with Elav-Gal4 did have defects, indicating that these 9 genes regulate GSCs from neurons and 14 genes regulate GSCs from other organs.

Tctp differentially regulates GSCs from different cells. We further characterized CG4800, a Drosophila translationally controlled tumour protein gene (Tctp). Its product was initially identified as a guanyl-nucleotide exchange factor for Drosophila Rheb GTPase, essential for growth and proliferation⁶², and was recently characterized as a regulator of ataxia telangiectasiamutated kinase activity to control genome stability and organ development in Drosophila⁶³. Tctp knockdown with Act-Gal4, Nos-Gal4 and c587-Gal4 resulted in GSC differentiation defects, GSC loss and GSC differentiation defects, respectively (Fig. 4a–c). We further generated mosaic clones with mutant Tctp and found that Tctp is required for GSC maintenance (Fig. 4d–g). Therefore, Tctp is required in GSCs for GSC maintenance and in CySCs for GSC differentiation.

Vers has multiple roles in regulating testis stem cells. Heterochromatin protein 1 has several conserved roles, including gene silencing transcription, DNA replication and DNA repair⁵¹. In this screen, we identified l(3)j2D3 (CG6801, lethal (3) j2D3), which encodes a component of the heterochromatin protein 1 protein complex. We renamed the gene 'versatile' (vers) based on its versatile functions described below. vers knockdown by transgenic RNAi with an Act-Gal4 driver resulted in GSC differentiation defects (Supplementary Data 2). To examine the function of vers in the germline versus the soma, we knocked down *vers* using cell-type-specific Gal4s and two independently generated *UAS-vers*^{RNAi} transgenic fly lines. Depleting *vers* in the CySC lineage (*c587*>*vers*^{RNAi}; Fig. 5 and Supplementary Figs 1 and 2) caused GSC differentiation defects (Fig. 5a,b). We further examined CySC proliferation and changes in c587>vers^{RNAi} flies by staining with antibodies to Zfh1 (marks CySCs and their immediate cyst cell daughters; Fig. 5c,d), phospho-Histone 3 (Fig. 6a,b) and Tj (traffic jam, a transcription factor expressed in CySCs, early cyst cells and hub cells; Supplementary Fig. 2b,c).

Compared with wild-type testes, the numbers of Zfh1-positive, p-H3-positive and Tj-positive cells were either dramatically increased in the testes of c587>vers^{RNAi} flies, suggesting that vers knockdown caused a dramatic expansion of CySCs and GSC differentiation defects. We also stained the c587 > vers^{RNAi} flies with antibodies to Cic (capicua, a transcriptional repressor downregulated by EGFR signalling)⁶⁴ and Mad (Mothers against dpp, a transcriptional factor in the Dpp signalling pathway). Compared with wild-type testes, the numbers Cic-positive and Mad-positive cells were dramatically increased in the testes of c587 > vers^{RNAi} flies (Fig. 5e,f), suggesting that vers knockdown caused the downregulation of EGFR signalling and the activation of Dpp signalling. We made an anti-Vers antibody, stained the testes and midguts of adult Drosophila, and found that Vers is ubiquitously expressed and localized to the nucleolus (Fig. 6c,d).

We found that depleting vers in the germ cell lineage using Nos-Gal4 resulted in a significant decrease (Student's t-test, P < 0.05) in the number of GSCs associated with the hub compared with the wild-type control (Supplementary Fig. 1a-d). To examine the function of vers in GSCs, we generated negatively marked GSC clones of wild-type or vers^{j2D3} flies⁶⁵ using the FLP (flippase)/FRT (flippase recognition target) mosaic analysis technique⁶⁶. Testes with LacZ (arm-lacZ)-negative clones were counted 2 and 7 days after clone induction (ACI). As expected, the FRT^{80B} control testes contained many LacZ-negative GSCs and their differentiated progenies (Supplementary Fig. 3a,b,e) at 2 and 7 days ACI. The FRT^{80B}-vers^{j2D3} testes at 2 days ACI also contained LacZ-negative GSCs and their differentiated progenies (Supplementary Fig. 3c,e). However, at 7 days ACI, LacZ-negative vers homozygous mutant GSCs had recovered at negligible levels (Supplementary Fig. 3d,e). Staining testes containing vers mosaic clones with anti-Stat92E antibodies revealed similar levels of Stat protein in the wild-type and *vers*-mutant GSCs (Supplementary Fig. 2a), indicating that Stat signalling is not required for Vers's function in GSCs.

These data together suggest that the heterochromatin protein Vers has multiple roles in regulating testis stem cells. It regulates GSC maintenance through a Stat-independent pathway in GSCs and regulates GSC differentiation and CySC expansion, possibly by upregulating the EGFR signalling pathway and/or downregulating the Dpp signalling pathway in somatic cells.

Comparing our screen to previous stem cell screens. We compared our results with those of previous screens focused on neuroblast³⁵, female GSCs³⁶ and ISCs³⁷, to identify both common and unique factors regulating the self-renewal and differentiation of different stem cell systems (Fig. 6e). Among the 530 genes identified in the male GSC screen, 7 were identified in all four screens, 12 were identified in the neuroblast, ISC and male GSC screens, and 45 were identified in the neuroblast, female GSC and male GSC screens (Fig. 6e and Supplementary Data 4). Most of the shared genes are regulators of basic, intrinsic cellular processes, such as messenger RNA splicing, chromosome assembly and disassembly, the cytoskeleton, the mitotic chromosome and cytokinesis (Supplementary Data 4). Of the 530 genes, 286 (54%) were male GSC-specific genes and were likely to include external regulators from both the local environment and distant organs. Future characterization of the common stem cell intrinsic regulators will increase our understanding of basic stem cell biology and further study of the external regulators will help reveal how stem cell behaviours are modulated by both the local microenvironment and signals from distant organs.




Figure 6 | Cell proliferation in vers knockdown and Vers expression. (a) Wild-type (control) testis. (b) Knockdown of vers in the CySC lineage (c587-Gal4) resulted in an expansion of mitotic cells. Vers expression in the testis (c) and in the midgut (d). (e) Comparison of male GSC, ISC, neuroblast and female GSC RNAi screens. Number of genes identified in the male GSC, ISC, neuroblast and female GSC screens. Seven genes were found in all four screens; 9 genes were shared by male GSC, female GSC and neuroblast screens. Venn diagram was prepared from: http://bioinfogp.cnb.csic.es/tools/venny/index.html. Testes of the indicated flies were dissected, stained with the antibodies to phospho-Histone 3 (pH3) (red, marks mitotic cells) (a,b), Vers (red, all type of cells in the testis and midgut), Arm (green, hub cells in testis (c) and epithelial membrane in the midgut (d)) and DAPI (blue, marks the nuclei), and analysed by confocal microscopy. Asterisks indicate hub cells. Yellow arrows in d indicate ISC/EB (enteroblast) cells and white arrows indicate enterocyte (EC) cells. The flies were cultured at 29 °C for 7 days in a-d. Scale bars, 10 μm.

Discussion

Drosophila male GSCs provide an excellent genetic system for studying niche-stem cell interactions. To comprehensively identify genes and pathways that regulate GSC fates in the whole organism, we performed a genome-wide transgenic RNAi screen through the ubiquitous knockdown of genes in adult *Drosophila* and identified 530 genes that regulate various fates of adult *Drosophila* testis GSCs.

As we screened the RNAi lines eliciting an adult lethal phenotype, we might have missed some adult testis-specific genes in this screen. A number of published genes that function in testis stem-cell regulation were missing from the genes identified in our screen, including *zfh1* (ref. 22), *chinmo*²¹, genes in the Hh (hedgehog) signal transduction pathway^{23,67}, *baz* and *par-1* (ref. 30). In this screen, we focused only on genes whose RNAimediated knockdown affected GSC numbers. Loss-of-function of the above genes does not affect the GSC numbers; therefore, our screen did not identify them.

By analysing the genes identified here, we reached a number of important conclusions. First, more than half the genes identified in our screen regulate GSC fates non-autonomously. Second, both the local microenvironment and signals from distant organs regulate GSCs. Third, the characterization of Tctp and Vers showed that the same protein could regulate GSC fates differentially in different cell types and through multiple pathways. Fourth, mitosis/cytokinesis proteins are especially important for male GSC maintenance. We identified 69 mitosis/ cytokinesis genes whose RNAi-mediated knockdown resulted in a strong GSC loss phenotype. In our previous screen for Drosophila ISCs, we identified only eight mitotic genes²⁶. In previous genome-wide RNAi screens for neuroblast and female GSCs, only a few mitosis/cytokinesis genes were identified^{35–37}. Collectively, these findings suggest that mitosis/cytokinesis genes are more important in GSC regulation.

To better analyse our results, we generated a gene interaction network by querying publicly available databases containing yeast–two-hybrid interactions, protein–protein interactions, text-mining data and genetic interactions between *Drosophila* genes (Supplementary Figs 4 and 5). We performed a complex-enrichment analysis using COMPLEAT⁶⁸ and identified a number of protein complexes required for GSC fate determination (Supplementary Fig. 4). However, as we have not assigned our 530 genes to GSCs, CySCs or other cell types and the protein interaction network is only meaningful in the same cells, these data may only serve as an indirect reference.

To our knowledge, this is the first genome-wide gene-knockdown screen performed in a whole animal, to identify systemic regulators of stem cells. Stem cells are regulated both intrinsically and externally by signals from the local microenvironment³⁹ and distant organs^{40,41}. Thus, it is difficult to fully appreciate the complexity of stem cell regulation if we limit our screen to intrinsic regulators. By performing a genome-wide transgenic RNAi screen by the ubiquitous knockdown of genes with *Act-Gal4*, we identified intrinsic regulators, as well as regulators from the local microenvironment and distant organs. Further characterization of these genes by knocking them down with cell-type-specific Gal4s will lead to a better understanding of GSC regulation in the whole animal. Our findings provide valuable insight and resources for studying stem cell regulation at the organismal level.

Methods

Fly stocks. Oregon R or *UAS-lacZ*^{RNAi} were used as wild type. RNAi stocks used in this study were: tum^{RNAi} (BDSC number 35007 [BL35007]), $Nedd8^{RNAi}$ (BL33881), $bab2^{RNAi}$ (VDRC Transformant ID 49042 [v49042]), $Taf8^{RNAi}$ (v27870), Cka^{RNAi} (BL34522), vn^{RNAi} (v50538), $SMC2^{RNAi}$ (BL32369), $APC10^{RNAi}$ (BL34858), $Cenp-C^{RNAi}$ (BL34692), $Tctp^{RNAi}$ (BL32911), $vers^{RNAi-1}$ was original $I(3)j2D3^{RNAi-1}$ (v22439) and $vers^{RNAi-2}$ (v101038). $FRT^{42D}SMC2^{606842}$ and $FRT^{80B}vers^{12D3}$ were obtained from Kyoto Stock Center. $FRT^{82B}Tctp^{h5p}$ was provided by Dr Kwang-Wook Choi, Korea Advanced Institute of Science and Technology.

The following Gal4 alleles were used to drive UAS lines: Nos-Gal4 (nanos-Gal4VP16)⁶⁹, Act-Gal4 and Elav-Gal4, obtained from the BDSC; upd-Gal4 and c587-Gal4, provided by Dr Ting Xie, Stowers Institute for Medical Research;

and Bam-Gal4, provided by Dr Xin Chen, Johns Hopkins University. Flies were raised on standard fly food at 25 °C and 65% humidity, unless otherwise indicated.

RNAi stocks used in the screen. *UAS-RNAi* lines were generated by VDRC and the Transgenic RNAi Project, and are available at VDRC and the BDSC. The sequences used for VDRC knockdown strains are available for each line at https://stockcenter.vdrc.at and for BDSC knockdown strains are available for each line at http://flystocks.bio.indiana.edu.

Generation of mutant GSC clones. Clones of mutant GSCs were generated as previously described⁶. To generate SMC2-mutant GSC clones, $FRT^{42D}+$ and $FRT^{42D}SMC2^{flo6842}/Cyo$ virgin females were mated with males of genotype $FRT^{42D}arm$ -lacZ/Cyo; MKRS, hs-flp/+. To generate Tctp-mutant GSC clones, $FRT^{82B}+$ and $FRT^{82B}Tctp^{hSp}/TM3$, Sb virgin females were mated with males of genotype SM6, hs-flp/+; $FRT^{80B}arm$ -lacZ/TM3, Sb. To generate vers-mutant GSC clones, $FRT^{80B}+$ and $FRT^{80B}vers^{j2D3}/TM3$, Sb virgin females were mated with males of genotype SM6, hs-flp/+; $FRT^{80B}arm$ -lacZ/TM3, Sb. One- or 2-day-old adult males carrying an arm-lacZ transgene in trans to the mutant-bearing chromosome were heat shocked four times at 37 °C for 1 h, at intervals of 8–12 h. The males were transferred to fresh food every day at 25 °C. The testes were removed 1, 2 or 7 days after the first heat-shock treatment and processed for antibody staining.

RNAi-mediated gene depletion. Male *UAS-RNAi* transgenic flies were crossed with virgin female of genotype *Act-Gal4*; *tub-Gal80*^{ts} (*Act*^{ts}), *Nos-Gal4*, *upd-Gal4* or *c587-Gal4*; *tub-Gal80*^{ts} (*c587*^{ts}). The flies were cultured at 18 °C. Three- to 5-day-old adult flies with the appropriate genotype were transferred to new vials at 29 °C for 7 days before dissection.

Immunofluorescence staining and microscopy. For immunofluorescence staining, testes were dissected in $1\times PBS$, then transferred to 4% formaldehyde in $1\times PBS$ and fixed for 30 min. The testes were then washed three times in $1\times PBST$ (PBS containing 0.1% Triton X-100) for 2 min each, then blocked with 5% goat serum in $1\times PBST$ for 1 h. Samples were then incubated with primary antibody in PBST at 4 °C overnight. Samples were then washed for 30 min (three 10 min washes) in $1\times PBST$, incubated with secondary antibody in $1\times PBST$ at room temperature for 2 h, washed for 30 min (three 10 min washes), then rinsed in $1\times PBS$ (twice, 2 min each) and mounted in VECTASHIELD with DAPI (4,6-diamidino-2-phenylindole; Vector Labs).

Confocal images were obtained using a Zeiss LSM510 system and were processed with Adobe Photoshop 7.0. GSCs were scored as Vasa-positive cells adjacent to the hub (detected using Fas 3) and containing dot spectrosome (detected using 1B1). Only images with a clear view of the complete hub were used.

The following antisera were used: rabbit polyclonal anti-Vasa antibody (1:5000; gift from R. Lehmann), rabbit polyclonal anti-β-Gal (β-galactosidase) antibody (1:1,000; Cappel), mouse monoclonal anti-β-Gal antibody (1:100; Invitrogen), mouse monoclonal anti-Hts antibody 1B1 (1:4; Developmental Studies Hybridoma Bank (DSHB)), mouse monoclonal anti-Fas3 antibody (1:10; DSHB), rabbit polyclonal anti-green fluorescent protein (GFP) antibody (1:200; Molecular Probes), mouse monoclonal anti-GFP antibody (1:100; Invitrogen), rabbit polyclonal anti-Thr3-phosphorylated histone H3 antibody (1:200; Upstate), guinea pig polyclonal anti-Zfh1 (1:2000; gift from Dr James B. Skeath, Washington University in St Louis), rabbit polyclonal anti-Cic antibody (1:1,000; gift from Dr Iswar Hariharan, University of California, Berkeley) and rabbit polyclonal anti-Smad3 antibody (1:1,000; Cell Signaling). Rabbit polyclonal anti-Stat92E antibody (1:1,000), rabbit polyclonal anti-Vers/l(3)j2D3 antibody (1:1,000) and guinea pig polyclonal anti-Tj antibody were generated in our laboratory. Secondary antibodies were goat anti-mouse, goat anti-guinea pig and goat anti-rabbit IgG conjugated to Alexa 488 or Alexa 568 (1:400; Molecular Probes). DAPI (Molecular Probes) was used to stain DNA.

Quantitative PCR. Total RNA from Act > RNAi adult fly testes was isolated using the RNeasy Mini Kit (Qiagen) with on-column DNase digestion, to remove genomic DNA. Complementary DNA was synthesized using the ThermoScript RT-PCR System (Invitrogen). Real-time PCR analysis was performed on a real-time PCR system, Mastercycler Realplex (Eppendorf), using SYBR Green PCR Master Mix (Clontech). Primers were selected using FlyPrimer Bank. The quantitative PCR primers used are shown in Supplementary Table 1.

Bioinformatics analysis. The GSC genetic network (Supplementary Fig. 5) was generated using STRING 10.0. Complex analysis (Supplementary Fig. 4) was done using COMPLEAT (http://www.flyrnai.org/compleat/), a tool that annotates protein complexes from both literature and predictions from protein–protein network, and does gene-set enrichment analysis based on protein complexes.

Statistical analysis. Statistical analyses of GSC and CySC numbers (mean ± s.e.m.) were performed using GraphPad Prism programme. Sample

sizes (n) reported reflect the individual testis number. P-values were obtained between two groups using the Student's t-test. For all statistical analysis, differences were considered to be statistically significant at values of P < 0.05.

Data availability. The authors declare that the data supporting the findings of this study are available within the article and its Supplementary Information files, and from the corresponding author upon reasonable request.

References

- de Cuevas, M. & Matunis, E. L. The stem cell niche: lessons from the *Drosophila* testis. *Development* 138, 2861–2869 (2011).
- 2. Pereira, G. & Yamashita, Y. M. Fly meets yeast: checking the correct orientation of cell division. *Trends Cell Biol.* **21**, 526–533 (2011).
- Lenhart, K. F. & DiNardo, S. Somatic cell encystment promotes abscission in germline stem cells following a regulated block in cytokinesis. *Dev. Cell* 34, 192–205 (2015).
- Hardy, R. W., Tokuyasu, K. T., Lindsley, D. L. & Garavito, M. The germinal proliferation center in the testis of *Drosophila melanogaster*. J. Ultrastruct. Res. 69, 180–190 (1979).
- Srinivasan, S., Mahowald, A. P. & Fuller, M. T. The receptor tyrosine phosphatase Lar regulates adhesion between *Drosophila* male germline stem cells and the niche. *Development* 139, 1381–1390 (2012).
- Wang, H. et al. A Rap-GEF/Rap GTPase signaling controls stem cell maintenance through regulating adherens junction positioning and cell adhesion in *Drosophila* testis. *Dev. Cell* 10, 117–126 (2006).
- Amoyel, M., Sanny, J., Burel, M. & Bach, E.A. Hedgehog is required for CySC self-renewal but does not contribute to the GSC niche in the *Drosophila* testis. *Development* 140, 56–65 (2013).
- 8. Inaba, M., Buszczak, M. & Yamashita, Y. M. Nanotubes mediate niche-stemcell signalling in the *Drosophila* testis. *Nature* **523**, 329–332 (2015).
- Michel, M., Raabe, I., Kupinski, A.P., Pérez-Palencia, R. & Bökel, C. Local BMP receptor activation at adherens junctions in the *Drosophila* germline stem cell niche. *Nat. Commun.* 2, 415 (2011).
- Issigonis, M. et al. JAK-STAT signal inhibition regulates competition in the Drosophila testis stem cell niche. Science 326, 153–156 (2009).
- Singh, S. R. et al. Competitiveness for the niche and mutual dependence of the germline and somatic stem cells in the *Drosophila* testis are regulated by the JAK/STAT signaling. J. Cell. Physiol. 223, 500–510 (2010).
- Monk, A. C. et al. HOW is required for stem cell maintenance in the *Drosophila* testis and for the onset of transit-amplifying divisions. *Cell Stem Cell* 6, 348–360 (2010).
- Zhang, Z., Lv, X., Jiang, J., Zhang, L. & Zhao, Y. Dual roles of Hh signaling in the regulation of somatic stem cell self-renewal and germline stem cell maintenance in *Drosophila* testis. *Cell Res.* 23, 573–576 (2013).
- Stine, R. R., Greenspan, L. J., Ramachandran, K. V. & Matunis, E. L. Coordinate regulation of stem cell competition by Slit-Robo and JAK-STAT signaling in the *Drosophila* testis. *PLoS Genet.* 10, e1004713 (2014).
- Singh, S. R., Liu, Y., Zhao, J., Zeng, X. & Hou, S. X. The novel tumour suppressor Madm regulates stem cell competition in the *Drosophila* testis. *Nat. Commun.* 7, 10473 (2016).
- Liu, Y., Singh, S. R., Zeng, X., Zhao, J. & Hou, S. X. The nuclear matrix protein Megator regulates stem cell asymmetric division through the mitotic checkpoint complex in *Drosophila* testes. *PLoS Genet.* 11, e1005750 (2015).
- Gonzalez, J., Qi, H., Liu, N. & Lin, H. Piwi is a key regulator of both somatic and germline stem cells in the *Drosophila* testis. Cell Rep. 12, 150–161 (2015).
- Tarayrah, L., Li, Y., Gan, Q. & Chen, X. Epigenetic regulator Lid maintains germline stem cells through regulating JAK-STAT signaling pathway activity. *Biol. Open* 4, 1518–1527 (2015).
- Amoyel, M., Anderson, J., Suisse, A., Glasner, J. & Bach, E. A. Socs36E controls niche competition by repressing MAPK signaling in the *Drosophila* testis. *PLoS Genet.* 12, e1005815 (2016).
- Morillo Prado, J. R., Srinivasan, S. & Fuller, M. T. The histone variant His2Av is required for adult stem cell maintenance in the *Drosophila* testis. *PLoS Genet.* 9, e1003903 (2013).
- Flaherty, M. S. et al. chinmo is a functional effector of the JAK/STAT pathway that regulates eye development, tumor formation, and stem cell self-renewal in Drosophila. Dev. Cell 18, 556–568 (2010).
- Leatherman, J. L. & Dinardo, S. Zfh-1 controls somatic stem cell self-renewal in the *Drosophila* testis and nonautonomously influences germline stem cell selfrenewal. *Cell Stem Cell* 3, 44–54 (2008).
- Amoyel, M., Simons, B. D. & Bach, E. A. Neutral competition of stem cells is skewed by proliferative changes downstream of Hh and Hpo. EMBO J. 33, 2295–2313 (2014).
- Kiger, A. A., White-Cooper, H. & Fuller, M. T. Somatic support cells restrict germline stem cell self-renewal and promote differentiation. *Nature* 407, 750–754 (2000).
- Tran, J., Brenner, T. J. & DiNardo, S. Somatic control over the germline stem cell lineage during *Drosophila* spermatogenesis. *Nature* 407, 754–757 (2000).

- Eun, S. H., Shi, Z., Cui, K., Zhao, K. & Chen, X. A non-cell autonomous role of E(z) to prevent germ cells from turning on a somatic cell marker. Science 343, 1513–1516 (2014).
- Cheng, J. et al. Centrosome misorientation reduces stem cell division during ageing. Nature 456, 599–604 (2008).
- Yamashita, Y. M., Mahowald, A. P., Perlin, J. R. & Fuller, M. T. Asymmetric inheritance of mother versus daughter centrosome in stem cell division. *Science* 315, 518–521 (2007).
- Yuan, H., Chiang, C. Y., Cheng, J., Salzmann, V. & Yamashita, Y. M. Regulation of cyclin A localization downstream of Par-1 function is critical for the centrosome orientation checkpoint in *Drosophila* male germline stem cells. *Dev. Biol.* 361, 57–67 (2012).
- Inaba, M., Venkei, Z. G. & Yamashita, Y. M. The polarity protein Baz forms a
 platform for the centrosome orientation during asymmetric stem cell division
 in the *Drosophila* male germline. *Elife* 4, e04960 (2015).
- Yamashita, Y. M., Jones, D. L. & Fuller, M. T. Orientation of asymmetric stem cell division by the APC tumor suppressor and centrosome. *Science* 301, 1547–1550 (2003).
- Dietzl, G. et al. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. Nature 448, 151–156 (2007).
- Ni, J. Q. et al. A Drosophila resource of transgenic RNAi lines for neurogenetics. Genetics 182, 1089–1100 (2009).
- 34. Ni, J. Q. et al. A genome-scale shRNA resource for transgenic RNAi in Drosophila. Nat. Methods 8, 405–407 (2011).
- 35. Neumüller, R. A. et al. Genome-wide analysis of self-renewal in *Drosophila* neural stem cells by transgenic RNAi. Cell Stem Cell 8, 580–593 (2011).
- Yan, D. et al. A regulatory network of Drosophila germline stem cell selfrenewal. Dev. Cell 28, 459–473 (2014).
- 37. Zeng, X. et al. Genome-wide RNAi screen identifies networks involved in intestinal stem cell regulation in *Drosophila*. Cell Rep. 10, 1226–1238 (2015).
- Sanchez, C. G. et al. Regulation of ribosome biogenesis and protein synthesis controls germline stem cell differentiation. Cell Stem Cell 18, 276–290 (2016).
- Morrison, S. J. & Spradling, A. C. Stem cells and niches: mechanisms that promote stem cell maintenance throughout life. Cell 132, 598–611 (2008).
- Li, Z., Zhang, Y., Han, L., Shi, L. & Lin, X. Trachea-derived dpp controls adult midgut homeostasis in *Drosophila*. Dev. Cell 24, 133–143 (2013).
- 41. Shim, J. et al. Olfactory control of blood progenitor maintenance. Cell 155, 1141–1153 (2013).
- Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in *Drosophila* muscles regulates organism-wide proteostasis during aging. *Cell* 143, 813–825 (2010).
- Rajan, A. & Perrimon, N. *Drosophila* cytokine unpaired 2 regulates physiological homeostasis by remotely controlling insulin secretion. *Cell* 151, 123–137 (2012).
- Owusu-Ansah, E., Song, W. & Perrimon, N. Muscle mitohormesis promotes longevity via systemic repression of insulin signaling. Cell 155, 699–712 (2013).
- Xie, J. et al. Histone H3 threonine phosphorylation regulates asymmetric histone inheritance in the Drosophila male germline. Cell 163, 920–933 (2015).
- McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K. & Davis, R. L. Spatiotemporal rescue of memory dysfunction in *Drosophila. Science* 302, 1765–1768 (2003).
- Wilson, R. J., Goodman, J. L. & Strelets, V. B.FlyBase Consortium. FlyBase: integration and improvements to query tools. *Nucleic Acids Res.* 36, D588–D593 (2008).
- Tulina, N. & Matunis, E. Control of stem cell self-renewal in *Drosophila* spermatogenesis by JAK-STAT signaling. Science 294, 2546–2549 (2001).
- Muller, P., Kuttenkeuler, D., Gesellchen, V., Zeidler, M. P. & Boutros, M. Identification of JAK/STAT signalling components by genome-wide RNA interference. *Nature* 436, 871–875 (2005).
- 50. Grönholm, J. et al. Not4 enhances JAK/STAT pathway-dependent gene expression in *Drosophila* and in human cells. FASEB J. 26, 1239–1250 (2012).
- Alekseyenko, A. A. et al. Heterochromatin-associated interactions of Drosophila HP1a with dADD1, HIPP1, and repetitive RNAs. Genes Dev. 28, 1445–1460 (2014).
- Betz, A. et al. A Drosophila PIAS homologue negatively regulates stat92E. Proc. Natl Acad. Sci. USA 98, 9563–9568 (2001).
- Shi, S., Calhoun, H. C., Xia, F., Li, J., Le, L. & Li, W. X. JAK signaling globally counteracts heterochromatic gene silencing. *Nat. Genet.* 38, 1071–1076 (2006).
- D'Avino, P. P., Giansanti, M. G. & Petronczki, M. Cytokinesis in animal cells. Cold Spring Harb. Perspect. Biol. 7, a015834 (2015).
- Fededa, J. P. & Gerlich, D. W. Molecular control of animal cell cytokinesis. Nat. Cell Biol. 14, 440–447 (2012).
- Green, R. A., Paluch, E. & Oegema, K. (2012Cytokinesis in animal cells. Annu. Rev. Cell Dev. Biol. 28, 29–58.

- Goshima, G. Genes required for mitotic spindle assembly in *Drosophila* S2 cells. Science 316, 417–421 (2007).
- Somma, M. P. et al. Identification of *Drosophila* mitotic genes by combining co-expression analysis and RNA interference. *PLoS Genet.* 4, e1000126 (2008).
- Papagiannouli, F. & Mechler, B. M. Discs large regulates somatic cyst cell survival and expansion in *Drosophila* testis. Cell Res. 19, 1139–1149 (2009).
- 60. Papagiannouli, F. & Mechler, B. M. Discs large in the *Drosophila* testis: an old player on a new task. *Fly* **4**, 294–298 (2010).
- Sarkar, A. et al. Antagonistic roles of Rac and Rho in organizing the germ cell microenvironment. Curr. Biol. 17, 1253–1258 (2007).
- 62. Hsu, Y. C., Chern, J. J., Cai, Y., Liu, M. & Choi, K. W. *Drosophila* TCTP is essential for growth and proliferation through regulation of dRheb GTPase. *Nature* 445, 785–788 (2007).
- Hong, S. T. & Choi, K.W. TCTP directly regulates ATM activity to control genome stability and organ development in *Drosophila melanogaster*. Nat. Commun. 4, 2986 (2013).
- Tseng, A. S. K. et al. Capicua regulates cell proliferation downstream of the receptor tyrosine kinase/Ras signaling pathway. Curr. Biol. 17, 728–733 (2007).
- Spradling, A. C. et al. The Berkeley Drosophila genome project gene disruption project. Single P-element insertions mutating 25% of vital Drosophila genes. Genetics 153, 135–177 (1999).
- Xu, T. & Rubin, G. M. Analysis of genetic mosaics in developing and adult Drosophila tissues. Development 117, 1223–1237 (1993).
- Michel, M., Kupinski, A. P., Raabe, I. & Bökel, C. Hh signalling is essential for somatic stem cell maintenance in the *Drosophila* testis niche. *Development* 139, 2663–2669 (2012).
- 68. Vinayagam, A. et al. Protein complex-based analysis framework for high-throughput data sets. Sci. Signal. 6, rs5 (2013).
- Van Doren, M., Williamson, A. L. & Lehmann, R. Regulation of zygotic gene expression in *Drosophila* primordial germ cells. *Curr. Biol.* 8, 243–246 (1998).

Acknowledgements

We thank Ting Xie, Kwang-Wook Choi, Xin Chen, VDRC, Kyoto, National Institute of Genetics (NIG) and the BDSC for fly stocks; Ruth Lehmann, James B. Skeath, Iswar Hariharan and the DSHB for antibodies; Dong Yan and Yanhui Hu for help analysing genes; and Dr Stephen Lockett and Kimberly Peifley of Optical Microscopy and Analysis Laboratory (OMAL) for help with the confocal microscopy. This research was supported by the Intramural Research Program of National Institutes of Health, National Cancer Institute.

Author contributions

Y.L. and S.X.H. conceived and designed the experiments. Y.L., Q.G., B.C., H. L., S.R.S., J.M., J.E., A.M.W. and S.X.H. performed the experiments. S.R.S. and B.C. performed the bioinformatics. Y.L., B.C., S.R.S. and S.X.H. analysed the data. S.X.H., S.R.S., Y.L. and B.C. prepared the manuscript. S.R.S. revised the manuscript. All authors approved the final version of the manuscript.

Additional information

Supplementary Information accompanies this paper at http://www.nature.com/

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at http://npg.nature.com/reprintsandpermissions/

How to cite this article: Liu, Y. *et al.* Whole-animal genome-wide RNAi screen identifies networks regulating male germline stem cells in *Drosophila. Nat. Commun.* 7:12149 doi: 10.1038/ncomms12149 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this

article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2016