


Welcome To BiO480 – NDDs lectures 2024-2025

EPFL Course organisation:



I. Modeling neurodegenerative disorders (NDDs) in the Lab

II. Drugs therapy in Clinical trials

III. Bench techniques applied to early diagnosis

I. Modeling neurodegenerative disorders (NDDs) in the Lab

II. Drugs therapy in Clinical trials

III. Bench techniques applied to early diagnosis




Modeling neurodegenerative disorders (NDDs) in the Lab: What does it matter?

Implications for Therapeutic Strategies



Bench to Bedside:

Translating Science from the lab to the Clinic




- The cost of developing a **new drug** that gets approval is estimated at around \$2-3 billion.
- Trials which failed at phase I or II wasted around \$6 million.
- Trial failed at phase III resulted in a loss of approximately \$77 million.
- The time it takes for an approved drug to reach the market is 10-12 years.

Having the right model that mimic human pathology is fundamental for drug discovery

Image source: www.yourgenome.org


Bench to Bedside:

Translating Science from the lab to the Clinic



Where are we in NDDs and finding a cure? ★










- Lack of approved therapies to treat or slow the progression of the NDD
- Clinical trials of drugs/antibodies targeting protein aggregation continue to fail
- No early diagnosis for NDDs

Poor understanding of the cellular mechanisms responsible for the formation of protein inclusions found in patients

EPFL Lewy bodies in PD and Synucleinopathies: Cause or Consequence?



The level of LB pathology correlates with disease development, cognitive decline or disease progression and severity




Proteins aggregates are found in the brain of healthy individuals Protein aggregates are absent in specific genetic forms of PD

Answering this question requires the development of cellular or animal models that recapitulates the process of LB formation

EPFL Understand Parkinson's disease:

Lashuel Lab

Preclinical models that accurately replicate the human pathology are essential



- 1. What triggers aSyn aggregation?
- 2. What is the mechanism of LB formation?
- 3. Are LB toxic or protective or both?


The ideal model of PD should enable to dissect all the steps leading to Lewy body formation and neurodegeneration.

PD Symposium - October 2024

EPFL Understand Parkinson's disease:



Preclinical models that accurately replicate the human pathology are essential




Such models will enable to shed light on potential therapeutic target to develop disease-modifying therapies.

PD Symposium – October 2024

Back to the basics -

In vitro studies to understand the origin of the NDDs




Guerrero-Ferreira R et al., 2018, eLife.

Before going to cellular models, we need first to understand the basic questions:

- a) What is the aggregation process?
- c) What is the kinetics of aggregation?
- d) What are the biophysics properties of the different species?
- e) Use Bioinformatics to find potential drugs/compounds that prevent aggregation or disassemble fibrils?

Back to the basics -

In vitro studies to understand the origin of the NDDs




Tau CryoEM structures - AD

Before going to cellular models, we need first to understand the basic questions:

- a) What is the aggregation process?
- c) What is the kinetics of aggregation?
- d) What are the biophysics properties of the different species?
- e) Use Bioinformatics to find potential drugs/compounds that prevent aggregation or disassemble fibrils?

So how do we make each of these species in a test tube?


How to make fibrils and co? Step #1: production of prion-like proteins at the monomeric stage






Recombinant

naked fibrils (in vitro)





How to make fibrils and co?






How to model NDDs at the bench toward therapeutics discovery


Oligomers

Fibrils

Recombinant naked fibrils (in vitro)


Step #2: aggregation of prion-like proteins – In vitro






We can produce the different types of proteins from the soluble (physiological state) to the aggregates (pathological states) in vitro


OK but what can we do with these proteins at the bench side?







At the bench side:


Understand how disease-associated mutations influence aggregation of WT proteins





- What can trigger protein aggregation (toxic insults: pesticides, lipids, other proteins, Ph, temperature.....)
- Follow the type of aggregates formed (oligomers, fibrils.....), their biophysics properties and structures (EM, CryoEM, NMR, AFM.....
- Advantage: fast (<48h), cheap (<10\$/plate), high-throughtput screening (386 well plates)





At the Drug discovery/screening side:



- Test library of:
 - chemical compounds
 - antibodies
 - venoms













- Advantage: fast (<48h), cheap (<10\$/plate), high-throughtput screening (386 well plates)
- Disadvantage: Far from the complexity of an organism or even a neuron






At the Biomarker side: New test for early diagnosis - 2023

Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL Prion-like aggregates: a new biomarker for early diagnosis *



Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL Bench side: in vitro models to study NDDs *



CSF

fluid

Can we dedect aggregates (aSyn, Tau, aß) in body fluid? And how?

Hypothesis:

If body fluids contains aggregates, then these aggregates should still retain their prion properties and be able to seed monomeric proteins

newly formed aggregates Monomers Seeds



Can we dedect aggregates (aSyn, Tau, aß) in body fluid? And how?

Hypothesis:

If body fluids contains aggregates, then these aggregates should still retain their prion properties and be able to seed monomeric proteins

newly formed aggregates Monomers Seeds **Newly formed** Monomer (aSyn, Tau, aβ) **Aggregates** (Soluble = Non-aggregated form) = seeds aggregates

Bio480 – How to model NDDs at the bench toward therapeutics discovery




CSF


fluid


Can we dedect aggregates (aSyn, Tau, aβ) in body fluid? And how?

Hypothesis:

If body fluids contains aggregates, then these aggregates should still retain their prion properties and be able to seed monomeric proteins







At the Biomarker side: New test for early diagnosis – clinical phases 2023

Patients with loss of smell

Group	CSF asyn SAA +
All PD	97% (548/567)
Sporadic PD	99% (402/407)
GBA PD	100% (59/59)
LRRK2 PD	85% (70/82)
SNCA PD	100% (11/11)
PRKN PD	50% (2/4)
НС	30% (8/26)

Patients without loss of smell


Group	CSF asyn SAA +
All PD	59% (114/193)
Sporadic PD	75% (90/120)
GBA PD	29% (2/7)
LRRK2 PD	34% (21/62)
SNCA PD	n/a
PRKN PD	50% (2/4)
НС	3% (6/179)




SAA has been acquired in 2428 PPMI participants https://www.ppmi-info.org/


- Advantage: fast (<48h), cheap (<10\$/plate), high-throughtput screening (386 well plates)
- Disadvantage: false positive level

EPFL Challenges: A growing gap between the complexity of brain pathology and simplicity of cell-free systems?★




Sio480 – How to model NDDs at the bench toward therapeutics discovery







Shahmoradian S et al., Nat. Neuroscience, 2019

EPFL To what extent can we model the human PD pathology in cellular and animal models?



Reverse engineering Lewy bodies: how far have we come and how far can we go?





Till recently, the existing cellular and animal models do not recapitulate the biochemical and ultrastructural properties of LBs

EPFL The aSyn overexpression-based models induces aSyn fibrilization but not LB formation

Fares et al.,

Bio480 - How to model NDDs at the bench toward therapeutics discovery


Induction of de novo α -synuclein fibrillization in a neuronal model for Parkinson's disease


Mohamed-Bilal Fares^a, Bohumil Maco^a, Abid Oueslati^{a,1}, Edward Rockenstein^b, Natalia Ninkina^c, Vladimir L. Buchman^c, Eliezer Masliah^b, and Hilal A. Lashuel^{a,d,2}

"Brain Mind Institute, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, CH-1015, Switzerland; "Department of Neurosciences, University of California San Diego, La Jolla, CA 92093; "School of Blosciences, Cardiff University, Cardiff, CF10 3AX, United Kingdom; and "Qatar Biomedical Research Institute, Hamad Bin Khaliff University, Doha, 5925, Oator

Edited by Thomas C. Südhof, Stanford University School of Medicine, Stanford, CA, and approved January 4, 2016 (received for review July 25, 2015)








aSyn overexpression in aSyn KO neurons induces aSyn fibrillization, but not LB formation

PNA

Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL Which features of the human pathology should the * Cellular and animal models reproduce?



EPFL Reverse Engineering Lewy bodies: it takes a team


Lashuel Lab













© Lisa clark. The Scientist://www.the-scientist.com/features/is-it-time-to-rethink-parkinsons-pathology-66449

EPFL Models of aSyn pathology formation and propagation in the absence of aSyn overexpression




Addition of exogenous α -synuclein preformed fibrils to primary neuronal cultures to seed recruitment of endogenous α -synuclein to Lewy body and Lewy neurite-like aggregates

Laura A Volpicelli-Daley¹, Kelvin C Luk² & Virginia M-Y Lee² NATURE PROTOCOLS | VOL.9 NO.9 | 2014 | 2135




EPFL The extended version of the neuronal seeding model allows the formation and maturation of LB-like inclusions






EPFLCan we bridge the gap between the complexity of brain pathology and simplicity of cell-free systems?








Shahmoradian S et al., Nat. Neuroscience, 2019

EPFL The extended version of the neuronal seeding model allows the formation and maturation of LB-like inclusions










EPFL How far does the "LB in a dish" reproduce the human pathology?

Lashuel Lab





Lewy body in a dish




Impact on Cellular dysfunctions and cell death pathways (cellular assays)

EPFL The neuronal Lewy body model enables the reproduction of inclusions similar to those found in patients






Lewy bodies in primary neurons





Mahul-Mellier at al., 2020, PNAS

Lewy body in patients




Shahmoradian et al., Nature Neurosciences, 2019








Proteomics



Proteomic comparison




Mahul-Mellier et al.

Killinger et al. Xia et al. Licket et al. Datta et al.


The Lewy body seeding model allows for the dissection of the mechanisms underlying Parkinson's disease





EPFL Application: a powerful platform for drug screening






Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL Application: a powerful platform for drug screening





Can we slow down or reduce aSyn pathology?






Bio480 - How to model NDDs at the bench toward therapeutics discovery


PFL Automated high content screening image analysis pipeline for drug screening






EPFL The neuronal seeding model: a platform for drug screening





EPFL Application: a powerful platform for drug screening





Small molecules



Immunotherapy



Venoms



Can we slow down or reduce aSyn pathology?

EPFL The Lewy body seeding model allows for the dissection of the mechanisms underlying Parkinson's disease





Limitations of the model:


- Mouse aSyn
- Human aSyn
- Does not recapitulate the full biochemical and morphological diversity of PD pathology
- Short-timeline: Not suitable to study late stages of LB formation or LB Clearance



EPFL

Bench to Bedside: *

Translating Science from the lab to the Clinic



EPFL Lewy Body in a dish:

Translating the seeding model in human iDA











Dr. Anne-Laure Mahul-Mellier

Lukas Van den Heuvel

Manel Boussouf

Yllza Jasiqi

Prof. Hilal A. Lashuel



Dr. Razan Sheta



Maxime Teixeira




Prof. Abid Oueslati

EPFL Lewy Body in a dish:

Translating the seeding model in human iDA





Human iPSCs-derived dopaminergic neurons (iDA)

Current iPSC-based models of PD have limitations:

- 1) They fail to reproduce LB pathology in both neurites and cell bodies within a reasonable time
- 2) They require the artificial overexpression of aSyn to achieve any pathology.

EPFL Establishment of the seeding model in iDA cells






Dr. Razan Sheta

Maxime Teixeira

Prof. Abid Oueslati

Combining NGN2 programming and dopaminergic patterning for a rapid and efficient generation of hiPSC-derived midbrain neurons


Razan Sheta^{1,2,11}, Maxime Teixeira^{1,2,11}, Walid Idi^{1,2,11}, Marion Pierre³,
Aurelie de Rus Jacquet^{1,4}, Vincent Emond¹, Cornelia E. Zorca^{5,6}, Benoît Vanderperre^{7,8},
Thomas M. Durcan^{5,6}, Edward A. Fon^{5,6}, Frédéric Calon^{1,9}, Mohamed Chahine^{3,10} &
Abid Oueslati^{1,2}




~80 % of the iNeurons convert in iDopaminergic neurons (iDa)

EPFL Seeded iDAs form neuritic and somatic pS129 pathology









- Formation of aSyn-pS129 pathology first in neurites and then in cell bodies
- Percentage of aggregates formed in cell bodies increased overtime

EPFL The iDA seeding model allows



The formation of seeded aggregates positives for the human LBs markers






Bona fide LBs display sequestration of various pathological markers:

- **p62** (in cell bodies only, similar to what is observed in PD brains)
- ubiquitin
- Amyloid-specific dyes (ThS)








EPFL The iDA seeding model replicates




The PTMs signature associated with the human pathology



EPFL The iDA seeding model replicates

The PTMs signature associated with the human pathology







Van den Heuvel








Stéphanie Rosset




EPFL Advancing human pathology replication with iDA seeding model



Lewy body in a dish







LBs

Why is it so important to solve the cellular mechanisms and signalling pathways behind the NDDs?

Because they are druggable




I. Modeling neurodegenerative disorders (NDDs) in the Lab

II. Drugs therapy in Clinical trials

II. Bench techniques applied to early diagnosis

EPFL Cell-to-cell transmission, internalization, and trafficking of aSyn and therapeutic targets along the pathways



Menon et al., 2022

TABLE 1 | Therapeutic targets and drugs from Figure 1.

Target	Drugs		
Inhibitors of microglial activation	Hypoestoxide		
Figure 1B	Lenalidomide		
77	Candesartan cilexetil		
Microglial toll-like receptor inhibitors	NPT520-34		
Figure 1C	CU-CPT22		
3) Gap junction blockers	CBX		
Figure 1D	Gap3211		
	Gap2409		
	Gap2605		
4) Endocytosis inhibitors	Dynasore		
Figure 1F	Sertraline		
5) Autophagy and lysosome inducers	Rapamycin		
Figure 1H	Metformin		
	Trehalose		
	Nilotinib		
	KYP-2407		
	Ambroxol AR7		
6) Misfolding inhibitors	NPT200-11		
Figures 1L,M	Clr01		
	EGCG		
	Anle1386		
	Plant extracts		
	and phytochemicals		
	SynuClean-D		
	Apigenin		
7) Lipid-induced aggregation inhibitors	ENT-01		
Figure 1N	Trodusquenine		
	NPT200-11		

EPFL Cell-to-cell transmission, internalization, and trafficking of aSyn and therapeutic targets along the pathways – small molecules

Small molecule	Origin	Mechanism of action	Model(s)	Clinical stage	References
Beta2 adreno receptor agonists 1. Metaporterenol 2. Clembuterol 3. Salbutamol	Synthetic	Agonists decrease transcription of SNCA gene.	Primary rat neurons; neuron precursors derived from IPSCs from patient with SNCA-triplication. MPTP mice: protected TH+ neurons. WT mice: reduced expression of asyn in SN.	Synucleinopathies: N/A.	(215)
4. Salmeterol	Synthetic	Inhibits transcription of pro-inflammatory cytokines	MPTP mice: protected DA neurons against induced toxicity.	N/A	(216)
Disaggregation of asyn					
1. NPT200-11 (UCB0599)	Synthetic	Inhibits lipid-dependent nucleation	PDNG78 mice: reduced retinal a-syn pathology. Line 8 mouse model: reduced cortical u-syn pathology and astrogliosis, normalized striatal dopernine transporter levels, and improved motor phenotype.	Phase 2 (NCT04658186) in PD patients	(217, 218)
2. Anie138b	Synthetic	Anti-oligomeric	MPTP mice preserved dopamnerge nigal neurors. Thy1-XSDP mice docressed pathological a-syn oligomers. Mrt 2 transgerin exer reduced damsty of a-syn aggregates, improved dopamine neuron function and gait. MSA mice rescued motor dysfunction, reduced a-syn digomers and glid typolasmic inclusions, preserved DA neurons.	Phase Ib (NCT04685265) in PD patients	(219-221)
3. ENT-01 (squalamine)	Natural	Inhibits lipid-induced asyn sggregation	SHSY5Y cells: reduced asyn oligomer-induced toxicity. C.elegans overexpressing asyn: restored motor dysfunction.	Phase 2b KARMET (NCT03781791) in PD patients. Phase I (NCT03938922) PD dementia	(222)
4. Trodusquemine (MSI-1436)	Natural	Inhibits lipid-induced and fibril-induced asyn aggregation	SHYSY cells: reduced oligomer-induced toxicity. C.elegans PD model: reduced asyn inclusions, increased longevity.	N/A	(223)
5. CLR01	Synthetic	Molecular tweezer inhibits asyn aggregation through binding to lytine residues of asyn critical for its oligomerization	Zebra fish-asyn: improved phenotype, and survey, and suppressed a-syn aggregation. Thyt-asyn mice: attenuated motor dyslunction, however, did not reduce aggregated a-syn levels. LB treated iPSCs: decreased aggregation and toxicly. LB and ptf-seeded mice: reduced asyn pathology and DA neuron rescue.	N/A	(224-226)
6. EGCG	Natural	inhibits asyn fibriliogenesis, converts fibrils into amorphous, non-toxic protein aggregates.	HEK-293 WT asyn; MPTP mice: rescued motor dysfunction, protected TH+ SN neurons, increased dopamine expression, reduced a-syn. MPTP-monkeys: reduced algomeric a-syn in striatum, increased nigral TH+	Phase 3 completed, PROMESA clinical trial of EGCG in patients with MSA, failed to show a protective effect.	(227-229)

neurons and DA levels, improved

Small molecule	Origin	Mechanism of action	Model(s)	Clinical stage	References
7. Phenylbutyrate	Natural	Histone deacetylase inhibitor, increases DJ-1 production levels.	MPTP mice: protected dopaminergic neurons and preserved motor and cognitive functions. Increased brain α-syn clearance, reduced neuroinflammation.	Completed Phase I (NCT02046434) in PD patients.	(230, 231)
8. SynuClean-D	Synthetic	Binds to fibrits and disrupts their nucleation and elongation.	C.elegans PD models: decreased asyn aggregation, rescued dopaminergic neurodegeneration, and restored motor function.	N/A	(232)
9. Apigenin	Natural	Slows secondary nucleation of asyn.	Cell free assay: reduced number of oligomers formed in asyn aggregation assay.	N/A	(233)
Modulating autophagy 10. KYP-2047	Synthetic	Prolyl oligopeptidase inhibitor	Asyn pff-SHSY5Y cells: increased degradation of HMW asyn, reduced «-syn monomers; and «-syn secretion. Cell lines over-expressing mutant a-syn: reduced a-synuclain inclusions, improved cell visibility. A3OP-asyn mice: reduced soluble «-syn; improved motor symptoms	N/A	(234-235)
11. NPT520-34	Synthetic	Toll-like receptor 2 antagonist. Facilitates clearance of misfolded asyn via upregulating autophagy	L61 mice: reduced accumulation of α-syn and neuroinflammation, preserved dopamine signaling in brain, improved motor function.	Phase 1b	(172, 237)
Iron related molecules					
12. ATH434	Synthetic	Redistributes labile iron	MPTP, 6-OHDA, A53T a-syn mice: protected SNpc neurons, reduced asyn accumulation, improved motor performance. MSA mice: preserved dopaminergic nigral neurons, reduced asyn oligomerzation.	Completed Phase I in healthy elder volunteers. Plans for Phase 2.	(238)
13. Synucleozoid	Synthetic	Downregulates SNCA mRINA translation	SHSY5Y cells: reduced asyn pff-induced toxicity, decreased asyn expression.	N/A	(239)


EPFL aSyn is not the only target to try to prevent, slow or block PD !!!!!



Parkinson's Disease Drug Therapies in the Clinical Trial Pipeline: 2023 Update

Kevin McFarthing^a, Susan Buff^b, Gary Rafaloff^c, Brian Fiske^d, Leah Mursaleen^c, Rosic Fuest^c, Richard K. Wyse^c and Simon R.W. Stott^{c,+}






Symptomatic Therapy

Parkinson's Disease Drug Therapies in the Clinical Trial Pipeline: 2023 Update

Kevin McFarthing^a, Susan Buff^b, Gary Rafaloff^c, Brian Fiske^d, Leah Mursaleen^c, Rosic Fuest^c, Richard K. Wyse^e and Simon R.W. Stott^{c,+}

139 Clinical Trial

17'000 Patients worldwide



EPFL Current clinical trial in PD ★ Symptomatic therapies

Parkinson's Disease Drug Therapies in the Clinical Trial Pipeline: 2023 Update

Kevin McFarthing^a, Susan Buff^b, Gary Rafaloff^c, Brian Fiske^d, Leah Mursaleen^c, Rosic Fuest^c, Richard K. Wyse^c and Simon R.W. Stott^{c,+}

139 Clinical Trial

17'000 Patients worldwide


Symptomatic Therapy

Definition: Symptomatic therapies aim to relieve or manage the symptoms associated with a disease without addressing the underlying cause or progression of the disease itself.

Purpose: These therapies help improve the patient's **quality of life** by reducing discomfort, pain, or other challenging symptoms.

Key Point: Symptomatic therapies **do not halt or slow the progression** of the underlying disease but instead manage the discomfort or limitations it causes.

EPFL Current clinical trial in PD * Symptomatic therapies



- 1. Compensate for dopamine deficiency due to neuronal death through medication (Levodopa, dopamine agonists) to reduce the severity of motor symptoms.
- 2. Deep Brain Stimulation: Electrical impulses in the thalamus improve neuron communication and reduce motor symptoms (effective in ~10 to 20% of patients).



Current symptomatic therapies do not prevent neuronal loss, nor do they slow, halt, or prevent disease progression. They primarily improve the daily lives of patients by alleviating motor symptoms.


EPFL Current clinical trial in PD ★ Disease modifying therapies

Definition: Disease-modifying therapies are treatments that aim to impact the underlying mechanisms or causes of a disease to alter its natural progression. **They work to slow, stop, or sometimes even reverse the progression of the disease itself.**


Purpose: These therapies focus on the underlying pathology of a disease, aiming to improve long-term outcomes and potentially change the disease course.

Key Point: Disease-modifying therapies directly affect **the disease's root causes or progression**, <u>aiming for more long-term improvement rather</u> than temporary symptom relief.





EPFL Current clinical trial in PD



EPFL PD Medications Currently in Clinical Trials – 2024 Case of drug repurposing



Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL PD Medications Currently in Clinical Trials – 2024

Case of drug repurposing

What is most famous examples of drug repurposing?

Key Points on Viagra's repurposing:

•Original purpose:

Developed as a cardiovascular drug (sildenafil) to improve blood flow and reduce angina symptoms.

•Repurposing discovery:

During clinical trials, researchers observed its unexpected effect on erectile dysfunction (ED).

•Approval for new use:

Based on these findings, Pfizer reoriented its research, and in 1998, sildenafil was approved by the FDA as a treatment for ED.

•Success and impact:

Viagra became a widely recognized drug, generating billions in revenue and highlighting the potential of drug repurposing in uncovering new therapeutic applications.

Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL PD Medications Currently in Clinical Trials – 2024 Case of drug repurposing drug ★

Drug repurposing is a medication originally developed and approved for one disease or condition but later found to be effective in treating another, different disease.

The advantages of drug repurposing include:

- 1. Reduced development time: repurposed drugs have already undergone preclinical and early-stage clinical testing, which can **save years** in development time compared to new drugs.
- 2. Lower costs: since many **safety and pharmacology studies** have already been **completed**, the overall cost is significantly lower than developing a new drug from scratch.
- 3. Known safety profile: repurposed drugs have established **safety data**, which reduces the risk of adverse side effects and allows researchers to focus more on efficacy for the new indication.
- 4. Broadening therapeutic applications: repurposing allows researchers to explore **new ways** to address complex or multi-faceted diseases, such as cancer or neurodegenerative disorders, often leading to combination therapies. E.g. GLP-1 receptor
- 7. Economic incentives: repurposing can extend the commercial life of older drugs, offering economic benefits for pharmaceutical companies, especially when **patent extensions or new intellectual property protections** can be established.

How to model NDDs at the bench toward therapeutics discovery

EPFL PD Medications Currently in Clinical Trials – 2024 Case of drug repurposing drug – GLP1 receptor (Diabetes)

156 patients 12 months study Phase II in clinical trial – April 2024

ORIGINAL ARTICLE

Trial of Lixisenatide in Early Parkinson's Disease

W.G. Meissner, P. Remy, C. Giordana, D. Maltête, P. Derkinderen, J.-L. Houéto, M. Anheim, I. Benatru, T. Boraud, C. Brefel-Courbon, N. Carrière, H. Catala, O. Colin, J.-C. Corvol, P. Damier, E. Dellapina, D. Devos, S. Drapier, M. Fabbri, V. Ferrier, A. Foubert-Samier, S. Frismand-Kryloff, A. Georget, C. Germain, S. Grimaldi, C. Hardy, L. Hopes, P. Krystkowiak, B. Laurens, R. Lefaucheur, L.-L. Mariani, A. Marques, C. Marse, F. Ory-Magne, V. Rigalleau, H. Salhi, A. Saubion, S.R.W. Stott, C. Thalamas, C. Thiriez, M. Tir, R.K. Wyse, A. Benard, and O. Rascol, for the LIXIPARK Study Group*

Lixisenatide

SCIENCES MÉDECINE

Parkinson : une étude chez l'homme suggère l'effet protecteur d'un antidiabétique

D'après un essai clinique mené chez 156 patients par un réseau d'hôpitaux français, un antidiabétique de la famille des analogues du GLP-1 semble freiner, au bout d'un an, la perte des neurones qui produisent la dopamine dans le cerveau.

Par Florence Rosins

Publié le 03 avril 2024 à 28h00, modifié hier à 09h24 . 🐧 Lecture 4 min. Read in E

Le Monde

- Encouraging results (slowing of motor symptom progression over a one-year period).
- However, this study needs to be validated in a larger cohort of patients and repeated to assess long-term benefits
- But further studies at the clinical level are block by Sanofi ©

EPFL PD Medications Currently in Clinical Trials – 2024 Case of drug repurposing drug – GLP1 receptor




Bio480 - How to model NDDs at the bench toward therapeutics discovery

EPFL PD Medications Currently in Clinical Trials – 2024 Case of drug repurposing drug – GLP1 receptor

In type 2 diabetes there

Insulin




Repurposing GLP-1 Receptor Agonists for Parkinson's Disease: Current Evidence and Future Opportunities

Daniella Balduino Victorino¹ · Mariana Nejm¹ · Marcia Guimarães-Marques¹ · Fulvio Alexandre Scorza¹ · Carla Alessandra Scorza¹


UNDER INVESTIGATION

EPFL aSyn Aggregates: a biomarker for medical imaging (PET Scan) of the Synucleinopathies

DAT(Dopamine Active Transporter) scan









EPFL




II. Drugs therapy in Clinical trials

III. Bench techniques applied to early diagnosis (exercices part)

How to model NDDs at the bench toward therapeutics discovery

EPFL New tests for early PD diagnosis – strategy of detection \bigstar



Pro and con of each assay explained in the Biomarker exercice

EPFL New tests for early PD diagnosis

See exercice in the Moodle

"Biomarkers exercice"

→ Detailled explanations of the pro and cons of each assays

(WB, ELISA, IHC, SAA)

EPFL Any questions? Or Thoughts?



