


Dynamics and Kinetics. Exercises 10: Solutions

Problem 1




Recall: O_3 is non-linear $\Rightarrow$ # of normal modes: $s = 3N - 6 = 3$

Total quanta: $j = 40$

Probability P to find $m = 10, 20, 30$ quanta in a particular mode?

We can approach this question as a problem of placing j balls (quanta) in s boxes (modes) in such a way that at least m balls are in the same box.



The first trick, illustrated in the figure, is to consider the $s - 1$ partitions (“dividing sticks”) as additional elements to be arranged, i.e., balls and partitions must both be arranged to define a single overall configuration. Thus, the number of ways w to distribute j quanta in s modes can be translated to the problem of choosing j elements (i.e., the balls) out of $j + s - 1$ elements:

$$w = w_{\text{balls}} = \binom{j + s - 1}{j} = \frac{(j + s - 1)!}{j!(s - 1)!} \quad (1)$$

or choosing $s - 1$ elements (i.e., the partitions):

$$w = w_{\text{partitions}} = \binom{j + s - 1}{s - 1} = \frac{(j + s - 1)!}{(s - 1)!j!} = w_{\text{balls}}. \quad (2)$$

On the other hand, the number of ways w' to distribute $j - m$ remaining quanta in s modes if at least m quanta are in a given mode is:

$$w' = \binom{j - m + s - 1}{j - m} = \frac{(j - m + s - 1)!}{(j - m)!(s - 1)!}. \quad (3)$$

Finally, we need to know the fraction of times that situation 3 happens among 1 total possibilities:

$$P_{\text{exact}} = \frac{w'}{w} = \frac{(j - m + s - 1)!j!}{(j - m)!(j + s - 1)!} \quad (4)$$

which can be approximated to:

$$P_{\text{approx}} \approx \left(\frac{j - m}{j} \right)^{s-1} \quad (5)$$


If we use the Sterling approximation: $n! \approx \left(\frac{n}{e}\right)^n$, and the fact that $j - m \gg s - 1$

Plugging in the numbers in Eqs. (4) and (5) gives:

	$m = 10$	$m = 20$	$m = 30$
P_{exact}	0.576	0.268	0.0767
P_{approx}	0.563	0.250	0.0625


Problem 4

- a) According to Eq. (9), if we can plot $\frac{1}{k_1}$ vs. $\frac{1}{P}$ we should obtain a straight line:



We observe linear behavior at low pressures, but deviation from theory at high pressures.

- b) The energization step:



Using the data for the three lower pressures that do not deviate from the linear regression we obtain a slope = 115.15 Torr · s.

From problem 3:

$$k_1^{\text{press}} = \frac{k_1^{\text{conc}}}{RT} = \frac{1}{\text{slope}} = 8.7 \times 10^{-3} \text{Torr}^{-1} \text{s}^{-1}$$

From collision theory:

$$k_1^{\text{coll}} = \frac{1}{2} N_{\text{Av}} \langle u \rangle \sigma e^{-\frac{E^0}{RT}}$$

Computing the different “ingredients”:

- $\langle u \rangle = \left(\frac{8k_B T}{\pi \mu} \right)^{1/2} = \left(\frac{8RT}{\pi \frac{M}{2}} \right)^{1/2} = 749 \text{ m} \cdot \text{s}^{-1}$
- $\sigma = \pi d^2 = 7.85 \times 10^{-19} \text{ m}^2$
- $\frac{E^0}{RT} = 42.5$

$$k_1^{\text{coll}} = 5.9 \times 10^{-11} \text{ m}^3 \text{ s}^{-1} \text{ mol}^{-1} \quad \times \frac{1}{RT} \quad \underline{\underline{=}} \quad 9.6 \times 10^{-15} \text{ Pa}^{-1} \text{ s}^{-1} \quad \xrightarrow{133.33 \text{ Pa} = 1 \text{ Torr}} \quad \underline{\underline{=}} \quad 1.28 \times 10^{-12} \text{ Torr}^{-1} \text{ s}^{-1}$$

which is much smaller than the constant found experimentally!

c) Rate constant for the Hinshelwood theory (k_1^H):

$$k_1^H = k_1^{\text{coll}} \cdot r \quad \Rightarrow \quad r = \frac{1}{(s-1)!} \left(\frac{E^0}{RT} \right)^{s-1}$$

For but-2-ene:

$$N = 12; \text{non-linear} \Rightarrow \quad s = 3N - 6 = 30 \Rightarrow \quad r = 1.95 \times 10^{16} \Rightarrow \quad k_1^H = 4.98 \times 10^4 \text{ Torr}^{-1} \text{ s}^{-1}$$

which is too large!