

Chemical biology CH-313

Part 1 Biological systems	1 week
Part 2 Bio-membranes	7 weeks
Part 3 Investigation of biological systems	4 weeks
Part 4 Paper presentations	2 weeks

Ruud Hovius

CH-B3-424

Ruud.Hovius@epfl.ch




Laboratory of Biophysical Chemistry of Macromolecules LCBM

Introduction

Part 1 Biological systems

- 1 • Cellular conditions
- 2 • Individual
- 3 • Labeling proteins
- 4 • Molecular interactions
- 5 • Molecular sensors
and machines

1 • Cells - Very complicated conditions



Goodsell, Nat Chem Biol 3 (2007) p.681

- Highly heterogeneous - compartments with very different contents
- Transport and separation
- Highly crowded
- Many different components – e.g. ~5' 000 different gene products

=> Very far from an ideal system in physico-chemical terms

1 • Crowding & Size-dependent hinderance




versus



1 • Effects of Crowding: Size-dependent hinderance

Diffusion in the cytosol vs in water





1 • Effects of Crowding: Excluded volume & confinement



Size matters !!

Compare a small (A) and a large (B) molecule for accessible to centre of mass (blue) and excluded (pink) volume with in a solution containing large solutes ●

÷ in bulk solution



÷ in a pore



1 • Effects of Crowding: Effective concentrations

Molar concentrations [**A**] are multiplied by an *activity coefficient* a which depends on

1


log activity coefficient

3

.

1 • Effects of Crowding: Chemical equilibria

Imagine an oligomerization reaction:



characterized by a dissociation constant

$$K_d = \frac{[A]^n}{[A_n]}$$

2 • Quantities in biology

Macroscopic world

Length 1 decimeter

Volume 1 liter




#98163335

Cellular world

1 μm

10^{-15} liter




- Consequence of scale on the ***number of molecules*** present

□

		Macroscopic	Cellular
Salts	100 mM	10^{23}	10^8
Synaptic transmitter	1 mM	10^{21}	10^6
Hormone	1 nM	10^{15}	1

2 • Poisson distribution: To be or not to be




Siméon Denis Poisson
1781 - 1840

$$p(k) = P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

**“when the average is small,
the change to observe nothing is not null !”**

In fact, there is a distribution of the average λ with different probabilities p to observe value k :



=> the lower the average λ , the longer one has to observe to be sure !

2 • Small numbers reveal individual molecular characters

Two examples :

i • Detection of molecular properties of ***N*** molecules

signal $\propto$ ***N***

standard deviation $\propto \sqrt{\mathbf{N}}$



2 • Small vs big numbers

Synapses as example :

- Synaptic contact between neurons only few active receptors present


versus

- Neuro-muscular junction thousands of receptors



2 • The “n” in biology


Escherichia coli:
Volume $\sim 10^{-15}$ L



Several important molecules are rare, almost alone
=> these molecules are deciding !!



3 • Finding a needle in a haystack



protein of interest


Advantages of fluorescence methods

- Non-invasive => *in vivo*
- Sensitive => down to a single molecule
- On line => direct info from ns to days
- High spatial & temporal resolution
- High information content

Fluorescent labelling is needed to discriminate between molecules of interest and the rest


Focus:
Bio-orthogonal
labelling of proteins

3 • How to label your Protein - of - Interest “*poi*” ?




3 • Fluorescent proteins as label

- Fusion to protein of interest
N- or C- terminal, internal




- Within single cells or living organisms



R. Y. Tsien

- Parallel labelling of several proteins with different fluorescent proteins




Which one to use ? => fpbase.org


Where to get ? => [addgene](http://addgene.org)
The nonprofit plasmid repository

3 • Self-labelling fusion tags

- Fuse tag to protein-of-interest
- Tag = a “suicide” enzyme
- Providing proteins with properties that cannot be genetically encoded
- Using a single tag for different applications
- Many different probes can react with the same tag
- Popular examples are ***Halo-tag*** and ***Snap-tag***



- **Halo-tag**
 - based on a haloalkane dehalogenase
 - provide very fast labeling



11101





Nature Communications 2021-12-6989

OPEN

Engineering of a fluorescent chemogenetic reporter with tunable color for advanced live-cell imaging

Hela Benaïssa^{1,2}, Karim Ounoughi², Isabelle Aujard², Evelyne Fischer³, Rosette Goïame³, Julie Nguyen ⁴, Alison G. Tebo ^{1,2,9}, Cheng Li^{2,10,11}, Thomas Le Saux², Giulia Bertolin ⁵, Marc Tramier ⁵, Lydia Danglot ^{4,6}, Nicolas Pietrancosta^{1,7}, Xavier Morin ³, Ludovic Jullien ² & Arnaud Gautier ^{1,2,8✉}

Photoactive protein from *Hhalophila*




a engineered protein tag pFAST is a fluorescent chemogenetic reporter with tunable color. a Structures of the chromophores of the HBO, 5-UBD, and UDBA series positioned according to the wavelengths of maximal excitation and maximal absorption of their absorbance at 280 nm.

3 • Non-natural amino acids


Introduce amino acid analogues with handles for modification by “click” chemistry

Click-ing molecules together :


÷ The original Cu⁺-catalyzed



÷ Cu-free strain-promoted





÷ Inverse Diels-Alder reaction



3 • Non-natural amino acids

An example of a “click-pair



Xue, JACS 2016



The **combination** of **non-natural amino acids** and **click chemistry** yields a **high spatial precision** for the introduction of the fluorophore

3 • Small interacting molecules



- A cell-permeable microtubule-binding conjugate of a taxol-analogue and Silicon Rhodamine




Nature Methods 2014

- A labeled serotonin-analogue of interacting with the serotonin receptor in the cell membrane

serotonin



3 • Labelling with organic dyes - Which one ?





Many available dyes :

- Alexa's** Invitrogen
- Cy dyes** GE
- Atto's** Atto-tec
- CF's** Biotium
- SiR's** Spirochrome
- Abbrerrior
-

Criteria : Brightness, stability, non-specific interactions & cell-permeability


4 • Molecular interactions - Biological networks

- The cell is extremely complex : many intertwined processes of signaling and metabolism :
 - multi-dimensional networks
 - a strong spatial-temporal organization




4 • Biological networks – Reversible molecular interactions

- Interacting neurons




4 • Proteins - Modular assembly of functional domains

- Proteins must be multi-functional
 - specific interaction
 - regulation of interaction
 - effector
- Proteins are in general not monolithic globules




but rather have a beads-on-a-string multi-domain structures, where ***each domain has a specific function***



4 • Phospho-tyrosine signalling

Growth factor-activated receptors in plasma membrane

=> Active receptors phosphorylate themselves and other proteins



Writers:

- Receptor Tyr-Kinases :
activity is ligand-dependent

Erasers:

- p-tyrosine phosphatases :
activity is generally constitutive


Readers :

- proteins with SH2-domains that
bind P-Tyr

4 • Space & time in signalling: A thought experiment

E.g. for the phosphorylation state of protein M (purple hue indicate [M-Pi].


- the phosphatase **Phos** is homogeneously distributed in the cytosol, but
- the kinase is bound to
a central structure or



=> Localized activity creates concentration gradients

4 • Imaging tyrosine phosphorylation *in vivo*

- A fusion of and SH2 domain and GFP is expressed in mammalian cells
- At $t = 0$, the epidermal growth factor (EGF) is added



SH2-GFP translocates to
plasma membrane
binding to the phosphorylated
EGF-receptor

Finally, the EGF-receptor is
internalized.

NB : Nucleus is red

=> Hormone-induced phosphorylation and re-localisation of SH2 domain and receptor

4 • Scaffolds facilitate molecular encounters

Adaptors & scaffolds: allow certain proteins to get together in a specific orientation and order at a specific site in the cell


Adaptor

Scaffold


a) Enforced proximity



(b) Combinatory use of elements




(c) Dynamic regulation



4 • Space & time in signalling: A cellular mechanism

E.g. regulation of cell division of yeast.

- Observations:
- only big cells divide
 - all components needed seem to be present continuously
 - When nuclear Cdk1 is active => cell division
 - Pom1 inhibits Cdk1 via a cascade




4 • Effective molarity

A protein with a binding site for triangles and with an “internal” ligand ◁.

Q: how does this internal ligand compete with an external one ◀ ?


Or, what is the effective concentration of ◁ ?




4 • Effective molarity

A protein with a binding site for triangles and with an “internal” ligand ◁.
Q: how does this internal ligand compete with an external one ◀ ?

Or, what is the effective concentration M_{eff} of ◁ ?




$$M_{\text{eff}} = K_d^{\text{inter}} / K_d^{\text{intra}}$$



- Effective molarity is concentration independent !!


5 • Sensors based on fluorescent proteins and molecular interaction

° Analyte detection




Frommer (2009)

° Molecular interactions



Föster resonance energy transfer (FRET):



Transfer of excitation energy from one fluorophore “Donor” to an “Acceptor”; possible only when Donor and Acceptor are within a few nm.


Semi-Up-to-date list:

codex.dpb.carnegiescience.edu/db/biosensor

5 • Snif-it's for drug monitoring : LUCID's

- **LUCID's** : **L**uciferase-based **I**ndicators of **D**rugs

e.g. for the anti-cancer drug methotrexate ● with intern analog ●






Article <https://doi.org/10.1038/s41589-023-01350-1>

A general method for the development of multicolor biosensors with large dynamic ranges

Received: 11 November 2022
Accepted: 25 April 2023
Published online: 8 June 2023



Lars Hellweg¹, Anna Edenhofer¹, Lucas Barck¹, Magnus-Carsten Huppertz¹,
Michelle. S. Frei¹, Mirosław Tarnawski², Andrea Bergner¹, Birgit Koch¹,
Kai Johnsson^{1,3} & Julien Hiblot¹✉



5 • Nanopore sequencing : From dream to reality


~2000 : The dream

Hagan Bayley (Oxford)



5 • Nanopore sequencing : From dream to reality

Today's reality Minion from Oxford Nanopore



Protein sequencing :<https://www.biorxiv.org/content/10.1101/2021.07.13.452225v1>

Reviews

- **Chemical labelling**

Schafer et al “Recent advances in Bioorthogonal reactions” *Chimia* 2019,73,308-312

Lang & Chin “Bioorthogonal Reactions for Label”

ACS Chem Biol (2014) 9, 16-20

Lang & Chin “Cellular Incorporation of Unnatural Amino Acids and Bioorthogonal Labeling of Proteins”

Chem Rev (2014) 114, 4764-4806

Resch et al “Quantum dots versus organic dyes as fluorescent labels”

Nature Methods (2008) 5, 763–775

- **Fluorescent proteins**

Wang et al “Fluorescence Proteins, Live-Cell Imaging, and Mechanobiology”

Annu. Rev. Biomed Eng. (2008) 10, 1-38.

Tsien, Campbell et al. “The Growing and Glowing Toolbox of Fluorescent and Photoactive Proteins”

doi.org/10.1016/j.tibs.2016.09.010

- **Fluorescent ligands**

Baindur & Triggie “Concepts and Progress in the Development and Utilization of

Receptor-Specific Fluorescent ligands” *Med Res Rev* (1994) 14, 591-664

Current Opinion in Chemical Biology

Cell Biology by the Numbers <http://book.bionumbers.org/>