



## **Organometallic Chemistry Basics**



**Ligand dissociation** – is the loss of a ligand from a transition metal centre which does not involve a change in the oxidation state.



**Ligand association**– is the addition of a ligand to a transition metal centre that does not involve a change in the oxidation state at the metal centre.



**Oxidative addition** – involves addition of a substrate to a complex with a **two** electron oxidation at the metal. The more reduced a metal centre, the greater its reactivity towards oxidative addition.




Which of the following will be **more** reactive towards oxidative addition?



dppe = bis(diphenylphosphino)ethane

**Reductive elimination** – is the reverse of oxidative addition, combining ligand loss with a two electron reduction at the metal centre. The groups being eliminated must be in a *cis* orientation. Reductive elimination is more likely for compounds in high a oxidation state. Reductive elimination is the **Rate Determining Step** in many catalytic reactions, especially C-C coupling reactions.



R = alkyl, aryl etc


Which of the following is **least** likely to undergo reductive elimination?



If a complex  $\text{L}_n\text{M}(\text{A})(\text{B})$  undergoes facile reductive elimination to give  $\text{AB}$  and  $\text{L}_n\text{M}$ , then what can we say about the reverse reaction, oxidative addition of  $\text{AB}$  to  $\text{L}_n\text{M}$ ?

- $\text{L}_n\text{M}$  must oxidatively add  $\text{AB}$  with the same ease.
- $\text{L}_n\text{M}$  cannot undergo oxidative addition of  $\text{AB}$ .
- Whether  $\text{L}_n\text{M}$  can undergo oxidative addition depends on both  $\text{L}_n\text{M}$  and  $\text{AB}$ .
- The oxidative addition will be slow compared to the reductive elimination reaction.


**Migratory insertion** – involves a coordinated nucleophile such as an alkyl, aryl, alkoxide or amide, *cis* to a ligand, insert into the M-ligand bond. The oxidation state of the metal does not change.




Rate of insertion depends on the strength of the metal-migratory group bond:

### $\beta$ -Hydride transfer: olefin insertion and $\beta$ -elimination

The generic example of olefin insertion is into a metal-alkyl or metal-hydride bond:



The transition state involves an **agostic interaction** of the  $\beta$ -hydrogen of the new alkyl:





The formation of **metallacycles** is a very important step in many catalysed reactions.

For  $\beta$ -hydride elimination an open coordination site on the metal complex is required.

## Nucleophilic attack on coordinated ligands

Metals withdraw electron density from a ligand making the ligand electropositive and susceptible to nucleophilic attack. Nucleophilic attack at CO is involved in many catalytic processes where carbon monoxide is present as a substrate.








When pushing arrows with carbonyls the  $\text{M=C=O}$  valence bond form should be used.

## Summary of main organometallic reactions

|                                                                                                                                                                        | $\Delta VE$ | $\Delta OS$ | $\Delta CN$ |                                                                            |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|-------------|-------------|----------------------------------------------------------------------------|
| <b>Ligand association</b>                                                                                                                                              |             |             |             |                                                                            |
| $L_nM + L' \longrightarrow L_nM-L'$                                                                                                                                    | +2          | 0           | +1          | $Pt(PPh_3)_3 + PPh_3 \longrightarrow Pt(PPh_3)_4$                          |
| <b>Ligand dissociation</b>                                                                                                                                             |             |             |             |                                                                            |
| $L_nM-L' \longrightarrow L_nM + L'$                                                                                                                                    | -2          | 0           | -1          | $Cr(CO)_6 \longrightarrow Cr(CO)_5 + CO$                                   |
| <b>Oxidative addition</b>                                                                                                                                              |             |             |             |                                                                            |
| $L_nM + \begin{array}{c} A \\   \\ B \end{array} \longrightarrow L_nM \begin{array}{c} \diagup A \\ \diagdown B \end{array}$                                           | +2          | +2          | +2          | $IrCl(CO)(PPh_3)_2 + H_2 \longrightarrow IrCl(H)_2(CO)(PPh_3)_2$           |
| <b>Reductive elimination</b>                                                                                                                                           |             |             |             |                                                                            |
| $L_nM \begin{array}{c} \diagup A \\ \diagdown B \end{array} \longrightarrow L_nM + \begin{array}{c} A \\   \\ B \end{array}$                                           | -2          | -2          | -2          | $IrCl(H)_2(CO)(PPh_3)_2 \longrightarrow IrCl(H)_2(CO)(PPh_3)_2 + H_2$      |
| <b>Migratory insertion</b>                                                                                                                                             |             |             |             |                                                                            |
| $L_nM \begin{array}{c} \diagup A \\ \diagdown B \end{array} \longrightarrow L_nM-A \begin{array}{c} \diagdown B \end{array}$                                           | -2          | 0           | -1          | $(OC)_5Mn-CH_3 \longrightarrow (OC)_4Mn-\overset{\overset{O}{  }}{C}-CH_3$ |
| <b><math>\beta</math>-Elimination</b>                                                                                                                                  |             |             |             |                                                                            |
| $L_nM \begin{array}{c} H \\ \diagup \\ A \end{array} \begin{array}{c} \diagdown \\ B \end{array} \longrightarrow L_nM-H \begin{array}{c} \diagdown \\ A=B \end{array}$ | +2          | 0           | +1          | $Cp_2Ta-CH_2-CH_3 \longrightarrow Cp_2Ta(H)(\eta^2-CH_2=CH_2)$             |

## Decomposition of alkyl complexes

Although  $\beta$ -hydride transfer is the most important process for the decomposition of alkyl complexes (see lecture notes), many other processes are also known, and they are shown below:

|                                                               |                                                                                                                                                                                                                                                               |
|---------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\alpha$ -Hydrogen transfer<br>( $\alpha$ -elimination)       |  $\text{M}-\text{CH}_2\text{CH}_3 \rightleftharpoons \text{M}=\text{CH}_2 + \text{H}-$                                                                                       |
| Cyclometallations:<br><br>$\gamma$ -hydrogen transfer         |  $\text{M}-\text{CH}_2\text{CH}_2\text{CH}_2\text{R} \rightleftharpoons \text{M}-\text{CH}_2\text{CH}(\text{R})\text{CH}_2\text{R}$                                         |
| $\delta$ -hydrogen transfer                                   |  $\text{M}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}_6\text{H}_5 \rightleftharpoons \text{M}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}_6\text{H}_5$ |
| intramolecular elimination of hydrocarbon<br>(reductive elim) |  $\text{M}-\text{CH}_2\text{CH}_2\text{R} \longrightarrow \text{M} + \text{R}-\text{CH}_3$ <p>(R can be an alkyl or H)</p>                                                |
| Binnuclear<br>(intermolecular elimination)                    |  $\text{M}-\text{R} + \text{H}-\text{M} \longrightarrow \text{M}-\text{M} + \text{R}-\text{H}$                                                                            |
| Free radical<br>(homolytic fission)                           |  $\text{M}-\text{R} \longrightarrow \text{M}\cdot + \text{R}\cdot$                                                                                                         |