




Sparse stochastic processes

RKHS, splines, and Gaussian processes


Prof. Michael Unser, LIB



March 2017 EDEE Course

Splines as a unifying mathematical concept

Functional analysis



Signal processing

Numerical analysis

CONTENT

1. Preliminaries

- Brief overview of continuous-domain operators
- Inner products vs. duality products
- Self-adjoint operators and positive definite kernels
- 2. Reproducing kernel Hilbert spaces
- 3. Variational splines and representer theorems
- 4. Gaussian processes

3

Linear operators, adjoints and kernels

Pair of vector spaces \mathcal{X}, \mathcal{Y} with topological duals $\mathcal{X}', \mathcal{Y}'$

$$\mathcal{S}(\mathbb{R}^d) \subseteq \mathcal{X}, \mathcal{Y}, \mathcal{X}', \mathcal{Y}' \subseteq \mathcal{S}'(\mathbb{R}^d)$$

Linear operator $G: \mathcal{X} \to \mathcal{Y}$ and its adjoint $G^*: \mathcal{Y}' \to \mathcal{X}'$

$$\langle f, G\varphi \rangle = \langle G^*f, \varphi \rangle$$

■ "Integral" representation: $g \in \mathcal{S}'(\mathbb{R}^d \times \mathbb{R}^d)$ = Schwartz kernel of G

$$G: \varphi \mapsto \int_{\mathbb{R}^d} g(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) d\boldsymbol{y}$$
 $G^*: \varphi \mapsto \int_{\mathbb{R}^d} g(\boldsymbol{y}, \cdot) \varphi(\boldsymbol{y}) d\boldsymbol{y}$

Self-adjoint operator

The operator H is said to be **self-adjoint** if $H\{\varphi\} = H^*\{\varphi\}$ for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$

⇔ the kernel of the operator is symmetric

Positive-definite kernel

Definition

A symmetric kernel function $h: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is said to be positive semi-definite (or **positive definite**, for short) if

$$\sum_{m=1}^{N} \sum_{n=1}^{N} z_m h(\boldsymbol{x}_m, \boldsymbol{x}_n) z_n \ge 0$$

for any $N \in \mathbb{N}$, $x_1, \ldots, x_N \in \mathbb{R}^d$, and $z_1, \ldots, z_N \in \mathbb{R}$.

Equivalent conditions

■ The square matrix $\mathbf{H} \in \mathbb{R}^{N \times N}$ with entries $[\mathbf{H}]_{m,n} = h(\boldsymbol{x}_m, \boldsymbol{x}_n)$ is positive definite for any choice of $\boldsymbol{x}_1, \dots, \boldsymbol{x}_N \in \mathbb{R}^d$; i.e.,

$$\mathbf{z}^T\mathbf{H}\mathbf{z} = \langle \mathbf{H}\mathbf{z}, \mathbf{z} \rangle \geq 0$$
 for all $\mathbf{z} \in \mathbb{R}^N \iff$ The eigenvalues of \mathbf{H} are non-negative

 $\blacksquare \ \, \text{The operator} \ \, \mathbf{H}: \varphi \mapsto \int_{\mathbb{R}^d} h(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \ \, \text{is positive-definite; i.e.,}$

$$\langle \mathbf{H}\{\varphi\}, \varphi \rangle \ge 0 \quad \text{for all } \varphi \in \mathcal{S}(\mathbb{R}^d)$$

5

Aronszajn's theorem

Theorem (Moore-Aronszajn)

The kernel function $h: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is symmetric positive (semi-)definite if and only if there exists some Hilbert space \mathcal{H} and a families of elements $\{f_x\}_{x\in\mathbb{R}^d}$ in \mathcal{H} such that

$$h(\boldsymbol{x}, \boldsymbol{y}) = \langle f_{\boldsymbol{x}}, f_{\boldsymbol{y}} \rangle_{\mathcal{H}}.$$
 (1)

In particular, there is a unique Hilbert space \mathcal{H}_{rep} —the reproducing kernel Hilbert space of h—such that (1) holds with $f_x = h(\cdot, x)$.

Sketch of proof (direct part):

$$\begin{split} \sum_{m=1}^{N} \sum_{n=1}^{N} z_m h(\boldsymbol{x}_m, \boldsymbol{x}_n) z_n &= \sum_{m=1}^{N} \sum_{n=1}^{N} z_m \langle f_{\boldsymbol{x}_m}, f_{\boldsymbol{x}_n} \rangle_{\mathcal{H}} z_n \\ &= \langle \sum_{i=1}^{n} z_m f_{\boldsymbol{x}_m}, \sum_{j=1}^{n} z_n f_{\boldsymbol{x}_n} \rangle_{\mathcal{H}} \\ &= \| \sum_{i=1}^{n} z_m f_{\boldsymbol{x}_m} \|_{\mathcal{H}}^2 \geq 0 \end{split} \tag{bilinearity of } \langle \cdot, \cdot \rangle)$$

CONTENT

- 1. Preliminaries
- 2. Reproducing kernel Hilbert spaces
 - Definition RKHS
 - Decay and continuity properties
 - RKHS associated with an invertible operator
 - Operators with non-trivial null spaces
- 3. Variational splines and representer theorems
- 4. Gaussian processes

7

2. RKHS

In essence, any Hilbert space $\mathcal{H}=\{f:\mathbb{R}^d\to\mathbb{R}:\|f\|_{\mathcal{H}}=\sqrt{\langle f,f\rangle_{\mathcal{H}}}<\infty\}\subseteq\mathcal{S}'(\mathbb{R}^d)$ whose members f are "ordinary"—but, not necessarily bounded—functions on \mathbb{R}^d is a RKHS.

Definition

The Hilbert space $\mathcal{H}\subseteq\mathcal{S}'(\mathbb{R}^d)$ is said to be a **reproducing kernel Hilbert space** (RKHS) if the shifted Dirac impulse $\delta(\cdot-r_0)\in\mathcal{H}'$ for any $r_0\in\mathbb{R}^d$.

2.1 Definition of reproducing kernel

$$\delta_{\boldsymbol{x}_0} = \delta(\cdot - \boldsymbol{x}_0) \in \mathcal{H}' \quad \Rightarrow \quad \text{existence of } \delta_{\boldsymbol{x}_0}^* = \mathrm{R}\{\delta_{\boldsymbol{x}_0}\} \in \mathcal{H} \text{ such that}$$

$$f(\boldsymbol{x}_0) = \delta_{\boldsymbol{x}_0}(f) = \langle \delta(\cdot - \boldsymbol{x}_0), f \rangle = \langle \delta_{\boldsymbol{x}_0}^*, f \rangle_{\mathcal{H}},$$
 for all $f \in \mathcal{H}$ and any $\boldsymbol{x}_0 \in \mathbb{R}^d$

Definition

The **reproducing kernel** of a RKHS on \mathbb{R}^d is the function $h: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ such that

- (i) $h(x_0, \cdot) \in \mathcal{H}$ for all $x_0 \in \mathbb{R}^d$
- (ii) $f(x_0) = \langle h(x_0, \cdot), f \rangle_{\mathcal{H}}$ for all $f \in \mathcal{H}$ and $x_0 \in \mathbb{R}^d$.

9

Reproducing kernel: fundamental properties

Proposition: Let \mathcal{H} be a RKHS on \mathbb{R}^d . Then, its reproducing kernel $h: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ has the following properties.

- 1. It is unique.
- 2. $h(\boldsymbol{x}, \boldsymbol{y}) = \langle h(\boldsymbol{x}, \cdot), h(\cdot, \boldsymbol{y}) \rangle_{\mathcal{H}}$
- 3. Symmetry: $h({m x},{m y})=h({m y},{m x})$ for all ${m x},{m y}\in\mathbb{R}^d$
- 4. Positive definiteness.
- 5. The linear span of $\{h(\boldsymbol{x},\cdot), \boldsymbol{x} \in \mathbb{R}^d\}$ is dense in \mathcal{H} .
- 6. Link with the Riesz map: The operator $R: \varphi \mapsto R\{\varphi\} = \int_{\mathbb{R}^d} h(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) d\boldsymbol{y}$ is a unitary mapping $\mathcal{H}' \to \mathcal{H}$ with the property that $\langle u, Ru \rangle = \|u\|_{\mathcal{H}'}^2$ for all $u \in \mathcal{H}'$ (the dual space of \mathcal{H}).
- 7. Invertibility: The operator R admits a unique inverse $R^{-1}: \mathcal{H} \to \mathcal{H}'$ with the property that $\langle R^{-1}f, f \rangle = \|f\|_{\mathcal{H}}^2$ for all $f \in \mathcal{H}$.

Elements for the proof:

- $\blacksquare \text{ Symmetry: } h(\boldsymbol{x},\boldsymbol{y}) = \langle h(\boldsymbol{x},\cdot),h(\boldsymbol{y},\cdot)\rangle_{\mathcal{H}} = \langle h(\boldsymbol{y},\cdot),h(\boldsymbol{x},\cdot)\rangle_{\mathcal{H}} = h(\boldsymbol{y},\boldsymbol{x})$
- lacksquare Unicity: h is the kernel of the Riesz map $\mathcal{H}' o \mathcal{H}$

$$f(\boldsymbol{x}_0) = \langle \delta(\cdot - \boldsymbol{x}_0), f \rangle = \langle R\{\delta(\cdot - \boldsymbol{x}_0)\}, f \rangle_{\mathcal{H}} = \langle h(\cdot, \boldsymbol{x}_0), f \rangle_{\mathcal{H}}$$

$$\Leftrightarrow h(\cdot, \boldsymbol{x}_0) = R\{\delta(\cdot - \boldsymbol{x}_0)\}$$

- Positive definiteness: link with inner product
- Denseness in \mathcal{H} : Consider any $g \in \mathcal{H}$ that is orthogonal to the linear span of $\{h(\boldsymbol{x},\cdot)\}_{\boldsymbol{x}\in\mathbb{R}^d}$. Then, $\langle g,h(\boldsymbol{x},\cdot)\rangle=0$ for every $\boldsymbol{x}\in\mathbb{R}^d$, which, by the reproducing kernel property, is equivalent to g=0.
- Link with Riesz map and invertibility follow from Riesz' theorem

$$\langle v,u\rangle=\langle v^*,u\rangle_{\mathcal{H}}=\langle \mathrm{R} v,u\rangle_{\mathcal{H}}=\langle v,\mathrm{R}^{-1}u\rangle_{\mathcal{H}'}=\langle v,u^*\rangle_{\mathcal{H}}$$
 for all $v\in\mathcal{H}'$ and $u\in\mathcal{H}$

11

2.2 Decay and continuity properties

Control of the rate of algebraic decay/growth: $\alpha \in \mathbb{R}$

$$\begin{split} \|f\|_{\infty,\alpha} &= \operatorname{ess \; sup}_{\boldsymbol{x} \in \mathbb{R}^d} \left(|f(\boldsymbol{x})| (1 + \|\boldsymbol{x}\|)^{\alpha} \right) \\ &= \sup_{\boldsymbol{x} \in \mathbb{R}^d} \left(|f(\boldsymbol{x})| (1 + \|\boldsymbol{x}\|)^{\alpha} \right) \quad \text{(if f is continuous)} \end{split}$$

$$L_{\infty,\alpha}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : ||f||_{\infty,\alpha} < +\infty \right\}$$

$$C_{\mathrm{b},\alpha}(\mathbb{R}^d) = \left\{ f: \mathbb{R}^d \to \mathbb{R} \text{ continuous and s.t. } \|f\|_{\infty,\alpha} < +\infty \right\}.$$

Both are Banach spaces with

$$\mathcal{S}(\mathbb{R}^d) \subseteq C_{b,\alpha+\beta}(\mathbb{R}^d) \subseteq C_{b,\alpha}(\mathbb{R}^d) \subseteq L_{\infty,\alpha}(\mathbb{R}^d) \subseteq \mathcal{S}'(\mathbb{R}^d),$$

for any $\beta \geq 0$

Continuity and decay (Cont'd)

Continuous functions of slow growth

For any continuous function $f \in \mathcal{S}'(\mathbb{R}^d)$, there exists some (critical) exponent $\alpha_0 \in \mathbb{R}$ such that $f \in C_{\mathrm{b},\alpha}(\mathbb{R}^d)$ for all $\alpha \leq \alpha_0$

$$\Leftrightarrow |f(\boldsymbol{x})| \le \frac{\|f\|_{\infty,\alpha_0}}{(1+\|\boldsymbol{x}\|)^{\alpha_0}}$$

Kernel spaces

$$C_{\mathrm{b},lpha}(\mathbb{R}^d imes\mathbb{R}^d) \subseteq L_{\infty,lpha}(\mathbb{R}^d imes\mathbb{R}^d) \subseteq \mathcal{S}'(\mathbb{R}^d imes\mathbb{R}^d)$$
 $\|h(\cdot,\cdot)\|_{\infty,lpha} = \sup_{oldsymbol{x},oldsymbol{y}\in\mathbb{R}^d}|h(oldsymbol{x},oldsymbol{y})|(1+\|oldsymbol{x}\|)^lpha(1+\|oldsymbol{y}\|)^lpha$

$$C_{\mathrm{b},\alpha}(\mathbb{R}^d\times\mathbb{R}^d) = \left\{h: \mathbb{R}^d\times\mathbb{R}^d \to \mathbb{R} \text{ (jointly) continuous and s.t. } \|h(\cdot,\cdot)\|_{\infty,\alpha} < +\infty\right\}$$

13

RKHS of continuous functions

$$\mathcal{H} \subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d) \quad \Rightarrow \quad \|f\|_{\infty,\alpha} \leq C \|f\|_{\mathcal{H}}, \text{ for all } f \in \mathcal{H} \quad \text{(continuous embedding)}$$

$$\downarrow \qquad \qquad \qquad |f(\boldsymbol{x}_0)| = |\langle f, \delta(\cdot - \boldsymbol{x}_0) \rangle| \leq (1 + \|\boldsymbol{x}_0\|)^{-\alpha} \|f\|_{\infty,\alpha} \leq C_{\boldsymbol{x}_0} \|f\|_{\mathcal{H}}$$

$$\downarrow \qquad \qquad \qquad \downarrow$$

$$\delta(\cdot - \boldsymbol{x}_0) : f \mapsto f(\boldsymbol{x}_0) \qquad \qquad \delta(\cdot - \boldsymbol{x}_0) \in \mathcal{H}' \qquad \text{(RKHS property)}$$

Theorem 1: A bivariate function $h: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is the reproducing kernel of a RKHS $\mathcal{H} \subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d)$ with $\alpha \in \mathbb{R}$ if and only if it is positive-definite, separately continuous in each variable, and such that $h(\cdot,\cdot) \in L_{\infty,\alpha}(\mathbb{R}^d \times \mathbb{R}^d)$. In particular, this implies that

1.
$$h(oldsymbol{x}_0,\cdot)\in C_{\mathrm{b},lpha}(\mathbb{R}^d)$$
 for any $oldsymbol{x}_0\in\mathbb{R}^d$

2.
$$\|h(\cdot,\cdot)\|_{\infty,\alpha} = \sup_{x,y \in \mathbb{R}^d} |h(x,y)| (1 + \|x\|)^{\alpha} (1 + \|y\|)^{\alpha} < \infty$$

3.
$$A_{\alpha,h} = \sup_{\boldsymbol{x} \in \mathbb{R}^d} h(\boldsymbol{x}, \boldsymbol{x}) (1 + \|\boldsymbol{x}\|)^{2\alpha} < \infty$$

with Conditions 2. and 3. being equivalent.

Equivalence of Conditions 2. and 3.

Key property: $h(\boldsymbol{x}, \boldsymbol{y}) = \langle h(\boldsymbol{x}, \cdot), h(\cdot, \boldsymbol{y}) \rangle_{\mathcal{H}}$

PD-kernel equivalent of Cauchy-Schwarz Inequality: $|h({m x},{m y})|^2 \leq h({m x},{m x})h({m y},{m y})$

$$\Downarrow$$

$$\begin{aligned} \|h(\cdot,\cdot)\|_{\infty,\alpha}^2 &= \sup_{\boldsymbol{x},\boldsymbol{y} \in \mathbb{R}^d} \left| (1 + \|\boldsymbol{x}\|)^{\alpha} h(\boldsymbol{x},\boldsymbol{y}) (1 + \|\boldsymbol{y}\|)^{\alpha} \right|^2 \\ &\leq \sup_{\boldsymbol{x},\boldsymbol{y} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{2\alpha} h(\boldsymbol{x},\boldsymbol{x}) (1 + \|\boldsymbol{y}\|)^{2\alpha} h(\boldsymbol{y},\boldsymbol{y}) = A_{h,\alpha}^2 \end{aligned}$$

Conversely,

$$(1 + \|\boldsymbol{x}\|)^{\alpha} h(\boldsymbol{x}, \boldsymbol{y}) (1 + \|\boldsymbol{y}\|)^{\alpha} \le \|h(\cdot, \cdot)\|_{\infty, \alpha}$$

so that

$$A_{\alpha,h} = \sup_{\boldsymbol{x} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{\alpha} h(\boldsymbol{x}, \boldsymbol{x}) (1 + \|\boldsymbol{x}\|)^{\alpha} \le \|h(\cdot, \cdot)\|_{\infty, \alpha},$$

from which we deduce that $A_{\alpha,h} = ||h(\cdot,\cdot)||_{\infty,\alpha}$.

15

Simplified version of theorem: $h \in C_{b,\alpha}(\mathbb{R}^d \times \mathbb{R}^d) \Rightarrow \mathcal{H} \subseteq C_{b,\alpha}(\mathbb{R}^d)$

■ Boundedness: $\mathcal{H} \subseteq L_{\infty,\alpha}(\mathbb{R}^d)$

$$\begin{split} |f(\boldsymbol{x})| &= |\langle h(\boldsymbol{x},\cdot), f\rangle_{\mathcal{H}}| \\ &\leq \|h(\boldsymbol{x},\cdot)\|_{\mathcal{H}} \, \|f\|_{\mathcal{H}} = \sqrt{h(\boldsymbol{x},\boldsymbol{x})} \, \|f\|_{\mathcal{H}} \end{split} \tag{Cauchy-Schwarz}$$

$$\|f\|_{\infty,\alpha}^2 = \sup_{\boldsymbol{x} \in \mathbb{R}^d} |f(\boldsymbol{x})|^2 (1 + \|\boldsymbol{x}\|)^{2\alpha} \le \sup_{\boldsymbol{x} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{2\alpha} h(\boldsymbol{x},\boldsymbol{x}) \|f\|_{\mathcal{H}}^2 = A_{h,\alpha} \|f\|_{\mathcal{H}}^2 < \infty$$

■ Continuity: $\mathcal{H} \subseteq C_{b,\alpha}(\mathbb{R}^d) \subseteq L_{\infty,\alpha}(\mathbb{R}^d)$

$$\begin{split} \left|f(\boldsymbol{x}) - f(\boldsymbol{x}_0)\right|^2 &= \left|\langle h(\boldsymbol{x},\cdot) - h(\boldsymbol{x}_0,\cdot), f\rangle_{\mathcal{H}}\right|^2 \\ &\leq \left\|h(\boldsymbol{x},\cdot) - h(\boldsymbol{x}_0,\cdot)\right\|_{\mathcal{H}}^2 \left\|f\right\|_{\mathcal{H}}^2 \qquad \text{(Cauchy-Schwarz)} \\ &= \left(h(\boldsymbol{x},\boldsymbol{x}) + h(\boldsymbol{x}_0,\boldsymbol{x}_0) - 2h(\boldsymbol{x},\boldsymbol{x}_0)\right) \left\|f\right\|_{\mathcal{H}}^2. \end{split}$$

$$h \in C_{b,\alpha}(\mathbb{R}^d \times \mathbb{R}^d) \Rightarrow \lim_{\boldsymbol{x} \to \boldsymbol{x}_0} h(\boldsymbol{x}, \boldsymbol{x}) + h(\boldsymbol{x}_0, \boldsymbol{x}_0) - 2h(\boldsymbol{x}, \boldsymbol{x}_0) = 0$$

$$\Rightarrow \lim_{\boldsymbol{x} \to \boldsymbol{x}_0} |f(\boldsymbol{x}) - f(\boldsymbol{x}_0)|^2 = 0$$

Turning $L_2(\mathbb{R}^d)$ into a RKHS

 $L_2(\mathbb{R}^d)$ is **NOT** a reproducing kernel Hilbert space ?

Candidate reproducing kernel $\delta(\cdot - x_0) : \varphi \mapsto \langle \delta(\cdot - x_0), \varphi \rangle = \varphi(x_0)$

$$lacksquare \delta(\cdot - oldsymbol{x}_0)
otin \left(L_2(\mathbb{R}^d)
ight)' = L_2(\mathbb{R}^d)$$

- $lacksquare f(oldsymbol{x}_0)$ is not well defined pointwise for all $f\in L_2(\mathbb{R}^d)$
- Solution = Smoothing

"Smoothing" operator $G: L_2(\mathbb{R}^d) \to \mathcal{H} \subseteq C_{b,\alpha}(\mathbb{R}^d)$

Proposition

Let $g: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ be a kernel such that $g(\boldsymbol{x},\cdot) \in L_2(\mathbb{R}^d)$ for any fixed $\boldsymbol{x} \in \mathbb{R}^d$. Then, the output of the linear operator $G: w \mapsto f = \int_{\mathbb{R}^d} g(\cdot, \boldsymbol{y}) w(\boldsymbol{y}) \mathrm{d} \boldsymbol{y}$ is well defined pointwise for any $w \in L_2(\mathbb{R}^d)$. If, in addition, there is some $\alpha \in \mathbb{R}$ such that

$$\sup_{\boldsymbol{x} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{\alpha} \|g(\boldsymbol{x}, \cdot)\|_{L_2(\mathbb{R}^d)} < \infty,$$

then G is bounded from $L_2(\mathbb{R}^d) \to L_{\infty,\alpha}(\mathbb{R}^d)$.

17

2.4 RKHS associated with invertible operator

Native space

$$\mathcal{H}_{\mathrm{L}} = \left\{ f: \mathbb{R}^d o \mathbb{R} \text{ s.t. } \|f\|_{\mathcal{H}_{\mathrm{L}}} \stackrel{ riangle}{=} \|\mathrm{L} f\|_{L_2(\mathbb{R}^d)} < \infty
ight\} \subseteq L_2(\mathbb{R}^d)$$

Coercive regularisation operator: $L:\mathcal{H}_L \to L_2(\mathbb{R}^d)$


$$c\|f\|_{L_2(\mathbb{R}^d)} \leq \|\mathrm{L}f\|_{L_2(\mathbb{R}^d)} \quad = \|f\|_{\mathcal{H}_\mathrm{L}}$$
 continuity \Leftrightarrow specifies a valid norm

 \Leftrightarrow Existence of (stable) inverse operator: $G = L^{-1}: L_2(\mathbb{R}^d) \to \mathcal{H}_L$

- Inner product: $\langle f, g \rangle_{\mathcal{H}_{\mathbf{L}}} = \langle \mathbf{L}f, \mathbf{L}g \rangle = \langle \underbrace{(\mathbf{L}^*\mathbf{L})f}_{f^* \in \mathcal{H}_{\mathbf{L}}'}, g \rangle$
- Riesz maps

$$\begin{split} R^{-1} &= L^*L: \mathcal{H}_L \to \mathcal{H}_L' \\ R &= G^*G = (L^*L)^{-1}: \mathcal{H}_L' \to \mathcal{H}_L \end{split}$$

Reminder: Dual of a Hilbert space



The dual of a Hilbert space \mathcal{H} is a Hilbert space \mathcal{H}' with $\mathcal{H}'=R^{-1}(\mathcal{H}).$

19

Determination of reproducing kernel

$$h(\boldsymbol{x}, \boldsymbol{y}) = \mathrm{R}\{\delta(\cdot - \boldsymbol{y})\}(\boldsymbol{x}) \quad \text{ with } \quad \mathrm{R} = (\mathrm{L}^*\mathrm{L})^{-1}: \mathcal{H}_\mathrm{L}' o \mathcal{H}_\mathrm{L}$$

Hypothesis: $h(\boldsymbol{x},\cdot) \in C_{\mathrm{b},0}(\mathbb{R}^d)$ (continuous and bounded) for any (fixed) $\boldsymbol{x} \in \mathbb{R}^d$

$$\Rightarrow$$
 $\mathcal{H}_{\mathrm{L}} \subseteq C_{\mathrm{b}}(\mathbb{R}^d)$ is a RKHS (by Theorem 1)

■ Explicit determination for LSI (=convolution) operator

 $\hat{L}(oldsymbol{\omega}) = \mathcal{F}ig\{\mathrm{L}\deltaig\}(oldsymbol{\omega})$: frequency response of operator

$$h(\boldsymbol{x},\boldsymbol{y}) = \rho_{\mathrm{L^*L}}(\boldsymbol{x} - \boldsymbol{y}) \quad \text{ where } \quad \rho_{\mathrm{L^*L}}(\boldsymbol{x}) = \mathcal{F}^{-1} \left\{ \frac{1}{|\widehat{L}(\boldsymbol{\omega})|^2} \right\}(\boldsymbol{x})$$

Necessary (and sufficient) condition for coercivity: $c = \mathrm{ess} \inf_{\pmb{\omega} \in \mathbb{R}^d} |\hat{L}(\pmb{\omega})| > 0$

$$\|\mathbf{L}f\|_{L_2}^2 = \int_{\mathbb{R}^d} |\hat{L}(\boldsymbol{\omega})\hat{f}(\boldsymbol{\omega})|^2 \frac{\mathrm{d}\boldsymbol{\omega}}{(2\pi)^d} \ge c^2 \int_{\mathbb{R}^d} |\hat{f}(\boldsymbol{\omega})|^2 \frac{\mathrm{d}\boldsymbol{\omega}}{(2\pi)^d} = c^2 \|f\|_{L_2}^2$$

2.5 Factorization of reproducing kernel

- Motivation: Extension for non-coercive regularization operator
- lacksquare Riesz map: $R = GG^* : \mathcal{H}' \to \mathcal{H}$

$$\mathcal{H}' o \mathcal{H}_0 o \mathcal{H}$$
 with $\mathcal{H}_0 \subseteq L_2(\mathbb{R}^d)$

Inverse operator: $G = L^{-1}$ with Schwartz kernel $g(x, y) = G\{\delta(\cdot - y)\}(x)$

Theorem 2

Let $\mathcal{H}\subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d)$ be a RKHS equipped with the **inner product** $\langle f,g\rangle_{\mathcal{H}}=\langle \mathrm{L} f,\mathrm{L} g\rangle$. Then, there exits a unique continuous operator $\mathrm{L}^{-1}:L_2(\mathbb{R}^d)\to L_{\infty,\alpha}(\mathbb{R}^d)$ and a Hilbert space $\mathcal{H}_0\subseteq L_2(\mathbb{R}^d)$ such that the Riesz map from $\mathcal{H}'\to\mathcal{H}$ factors though \mathcal{H}_0 as $\mathrm{R}=\mathrm{L}^{-1}\mathrm{L}^{-1*}:\mathcal{H}'\to\mathcal{H}_0\to\mathcal{H}$.

The Schwartz kernel of L^{-1} denoted by $g(\cdot,\cdot)$ satisfies the estimate

$$\sup_{\boldsymbol{x} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{\alpha} \|g(\boldsymbol{x}, \cdot)\|_{L_2(\mathbb{R}^d)} = \sqrt{A_{\alpha, h}} < \infty$$

and is linked to the reproducing kernel $h(\cdot, \cdot)$ of \mathcal{H} by

$$h(oldsymbol{x},oldsymbol{y}) = \int_{\mathbb{R}^d} g(oldsymbol{x},oldsymbol{z}) \mathrm{d}oldsymbol{z}$$

$$g(\boldsymbol{y}, \boldsymbol{x}) = L\{h(\cdot, \boldsymbol{y})\}(\boldsymbol{x}).$$

21

Example 1: RKHS associated with an ortho-basis

- \blacksquare Orthonormal system: $\{\phi_n\}_{n\in\mathbb{N}}$ with $\phi_n\in C_{\mathrm{b}}(\mathbb{R}^d)\cap L_2(\mathbb{R}^d)$
- $\qquad \text{Inner product:} \quad \langle f,g\rangle_{\mathcal{V}} = \sum_{n\in\mathbb{N}} \langle \phi_n,f\rangle \langle \phi_n,g\rangle$
- lacksquare $\mathcal{V}=\overline{\operatorname{span}\{\phi_n\}}_{n\in\mathbb{Z}}$ equipped with $\langle\cdot,\cdot
 angle_{\mathcal{V}}$ is a Hilbert space
- Generic form of reproducing kernel

$$h_{\mathcal{V}}(\boldsymbol{x}, \boldsymbol{y}) = \sum_{n \in \mathbb{N}} \phi_n(\boldsymbol{x}) \phi_n(\boldsymbol{y})$$

Simplified setting

$$\mathcal{V} = \mathcal{V}' \quad \text{with} \quad \mathcal{V} \subseteq L_2(\mathbb{R}^d)$$

lacksquare Orthogonal projector $\operatorname{Proj}_{\mathcal{V}}:L_2(\mathbb{R}^d) o \mathcal{V}$

$$\operatorname{Proj}_{\mathcal{V}}\{f\}(\boldsymbol{x}) = \langle f, h_{\mathcal{V}}(\boldsymbol{x}, \cdot) \rangle = \sum_{n \in \mathbb{N}} \phi_n(\boldsymbol{x}) \langle f, \phi_n \rangle$$

 $\blacksquare \ \, \mathsf{Factorization} \ \, (\mathsf{trivial}) \colon \ \, \mathrm{Proj}_{\mathcal{V}} \circ \mathrm{Proj}_{\mathcal{V}} = \mathrm{Proj}_{\mathcal{V}}$

$$h_{\mathcal{V}}(\boldsymbol{x}, \boldsymbol{y}) = \langle h_{\mathcal{V}}(\boldsymbol{x}, \cdot), h_{\mathcal{V}}(\cdot, \boldsymbol{y}) \rangle$$

Example 2: RKHS of bandlimitted functions

lacksquare Subspace of bandlimited functions in $L_2(\mathbb{R})$ with Nyquist frequency $\omega_{\max}=\pi$

$$\mathcal{H}_{\text{sinc}} = \{ f \in \mathcal{S}'(\mathbb{R}) : (\text{sinc} * f) \in L_2(\mathbb{R}) \}$$

■ Regularization operator

$$L_{sinc}: \varphi \mapsto sinc * \varphi$$

$$\operatorname{sinc}(x) = \frac{\sin(\pi x)}{(\pi x)} = \mathcal{F}^{-1}\{\mathbb{1}_{[-\pi,\pi]}\}(x)$$

■ Reproducing kernel

$$h_{\text{sinc}}(x,y) = \text{sinc}(x-y) = \sum_{k \in \mathbb{Z}} \text{sinc}(x-k) \text{sinc}(y-k)$$

■ Reproduction formula = Shannon's Sampling theorem

$$\forall f \in \mathcal{H}_{\text{sinc}}: \quad f(x) = (\operatorname{sinc} * f)(x) = \sum_{k \in \mathbb{Z}} \langle f, \operatorname{sinc}(\cdot - k) \rangle \operatorname{sinc}(x - k)$$
$$= \sum_{k \in \mathbb{Z}} f(k) \operatorname{sinc}(x - k)$$

23

Example 3: Subspace of polynomials

 \mathbb{Z} $\mathcal{N} = \operatorname{span}\{p_1\} \subseteq C_{\operatorname{b}}(\mathbb{R}) \text{ with } p_1(x) = 1$

Alternative solution(s)

- Biorthogonal system: (δ, p_1) with $\langle \delta, p_1 \rangle = 1$
- (ϕ_1, p_1) with $\langle \phi_1, p_1 \rangle = 1$
- Inner product: $\langle f,g\rangle_{\mathcal{N}}=f(0)g(0)=\langle \delta,f\rangle\langle \delta,g\rangle$
- $\langle f, g \rangle_{\mathcal{N}} = \langle \phi_1, f \rangle \langle \phi_1, g \rangle$
- $lacksquare \mathcal{N} = \mathrm{span}\{p_1\}$ equipped with $\langle f,g \rangle_{\mathcal{N}}$ is a RKHS
- Reproducing kernel

$$h_{\mathcal{N}}(x,y) = p_1(x)p_1(y) = 1$$

- Projection operator:
- $\operatorname{Proj}_{\mathcal{N}}\{f\} = \langle \delta, f \rangle p_1$
- $\operatorname{Proj}_{\mathcal{N}}\{f\} = \langle \phi_1, f \rangle p_1$

but, there is no factorization ... ?!?

2.6 RKHS associated with derivative operator

Key characteristics

$$L = D = \frac{\mathrm{d}}{\mathrm{d}x}$$

- \blacksquare Green's function(s) $\rho_{\rm D}$ such that ${\rm D}\{\rho_{\rm D}\}=\delta$
 - Causal solution: $\mathbb{1}_+(x)$
 - Canonical solution: $\frac{1}{2} \mathrm{sign}(x) = \mathcal{F}\left\{\frac{1}{\mathrm{j}\omega}\right\}(x) = \mathbb{1}_+(x) \frac{1}{2}$
- Null space of dimension $N_0 = 1$:

$$\mathcal{N}_{D} = \{ q \in \mathcal{S}'(\mathbb{R}) : D\{q\} = 0 \} = \operatorname{span}\{p_1\} \quad \text{with} \quad p_1(x) = 1$$

- Related operators
 - $\quad \blacksquare \ \, \mathsf{Adjoint} \colon D^* = -D$

$$\langle \mathrm{D}f, \varphi \rangle = \underbrace{f(x)\varphi(x)|_{-\infty}^{+\infty}}_{=0} - \int_{\mathbb{R}} f(x)\varphi'(x)\mathrm{d}x = \langle f, -\mathrm{D}\varphi \rangle$$

- Green's function of D*D:

$$\rho_{\mathrm{D}^*\mathrm{D}}(x) = \mathcal{F}\left\{\frac{1}{\omega^2}\right\}(x) = -\frac{1}{2}|x| \in C_{\mathrm{b},-1}(\mathbb{R})$$

25

Derivative operator (Cont'd)

Native space

$$\mathcal{H}_{\mathrm{D}} = \{f: \mathbb{R} o \mathbb{R} \text{ s.t. } \mathrm{D}f \in L_2(\mathbb{R})\} \hspace{0.5cm} \subseteq C_{\mathrm{b},-1}(\mathbb{R}^d)$$

Proposition

 \mathcal{H}_D equipped with the inner product $\langle f,g\rangle_{\mathcal{H}_D}=\langle Df,Dg\rangle+f(0)g(0)$ is a RKHS whose reproducing kernel is

$$h_{\rm D}(x,y) = \frac{1}{2}(|x| + |y| - |x - y|) + 1.$$

 $h_{\mathrm{D}}(\cdot,y)\in C_{\mathrm{b},-1}(\mathbb{R})$ for any $y\in\mathbb{R}$ (first-order of growth)

Proof by verification

- 1) \mathcal{H}_{D} is a Hilbert space
 - lacksquare $\langle \mathrm{D}f,\mathrm{D}g \rangle$ is a semi-inner product
 - \bullet $\langle \mathrm{D}f, \mathrm{D}f \rangle = 0 \quad \Leftrightarrow \quad f = b_1 p_1 \in \mathcal{N}_\mathrm{D} \text{ with } b_1 \in \mathbb{R}$
 - $\blacksquare \ \|f\|_{\mathcal{H}_{\mathrm{D}}}^2 = \langle \mathrm{D}f, \mathrm{D}f \rangle + |f(0)|^2 = 0 \qquad \Leftrightarrow \quad f \in \mathcal{N}_{\mathrm{D}} \text{ and } f(0) = 0 \quad \Leftrightarrow \quad f = 0 \quad \text{(unicity)}$
- 2) Check reproducing kernel property of $h_{\mathrm{D}}(x,y)=\frac{1}{2}\big(|x|+|y|-|x-y|\big)+1$

$$D\{h_{D}(\cdot,y)\}(x) = \frac{1}{2} (\operatorname{sign}(x) - \operatorname{sign}(x-y)) = \mathbb{1}_{[0,y)}(x) \in L_{2}(\mathbb{R}) \Rightarrow h_{D}(\cdot,y) \in \mathcal{H}_{D}$$

$$D^{*}D\{h_{D}(\cdot,y)\}(x) = -D\{\frac{1}{2} (\operatorname{sign}(\cdot) - \operatorname{sign}(\cdot-y))\}(x) = -\delta(x) + \delta(x-y)$$

$$h_{D}(0,y) = \frac{1}{2} (|0| + |y| - |0-y|) + 1 = 1$$

Hence, for all $f \in \mathcal{H}_{\mathrm{D}}$,

$$\langle f, h_{\mathcal{D}}(\cdot, y) \rangle_{\mathcal{H}_{\mathcal{D}}} = \langle \mathcal{D}f, \mathcal{D}\{h_{\mathcal{D}}(\cdot, y)\} \rangle + f(0) \times 1$$
$$= \langle f, \mathcal{D}^* \mathcal{D}\{h_{\mathcal{D}}(\cdot, y)\} \rangle + f(0)$$
$$= (-f(0) + f(y)) + f(0) = f(y).$$

27

Native space for derivative operator

 \mathcal{H}_{D} : RKHS equipped with the inner product $\langle f,g \rangle_{\mathcal{H}_{\mathrm{D}}} = \langle \mathrm{D}f,\mathrm{D}g \rangle + f(0)g(0)$.

- Direct sum structure
 - "Quotient" space $\mathcal{H}_D/\mathcal{N}_L$: $\mathcal{H}_{D,\delta} = \{f \in \mathcal{H}_D : \langle \delta, f \rangle = f(0) = 0\}$ \Rightarrow RKHS with inner product $\langle Df, Dg \rangle$
 - Null space $\mathcal{N}_D = \{f \in \mathcal{H}_D : D\{f\} = 0\} = \operatorname{span}\{p_1\}$ $\Rightarrow \mathsf{RKHS} \text{ of dim } N_0 = 1 \text{ with inner product } \langle f, g \rangle_{\mathcal{N}_L} = f(0)g(0)$

$$\Rightarrow$$
 $\mathcal{H}_{\mathrm{D}} = \mathcal{H}_{\mathrm{D},\delta} \oplus \mathcal{N}_{\mathrm{L}}$

Every function $f \in \mathcal{H}_{\mathrm{D}}$ has a unique decomposition as $f = f_1 + f_2$

$$f_1 = (\operatorname{Id} - \operatorname{Proj}_{\mathcal{N}_{\mathcal{D}}})\{f\} = f - f(0)p_1 \in \mathcal{H}_{\mathcal{D},\delta}$$

$$f_2 = \operatorname{Proj}_{\mathcal{N}_{D}} \{f\} = \langle \delta, f \rangle p_1 = f(0)p_1 \in \mathcal{N}_{D}$$

2.7 Operators with non-trivial null spaces

Specification of native space: first attempt

$$L: \mathcal{H}_L \to L_2(\mathbb{R}^d)$$

$$\mathcal{H}_{L,\text{ext}} = \{ f \in \mathcal{S}'(\mathbb{R}^d) : \|Lf\|_{L_2(\mathbb{R}^d)}^2 < \infty \}$$

"Extended" null space: $\mathcal{N}_{L,\mathrm{ext}} = \{q \in \mathcal{H}_{L,\mathrm{ext}} : L\{q\} = 0\}.$

Potential problem for d>1 as $\mathcal{N}_{\mathrm{L,ext}}$ may not be **finite-dimensional** !!!

■ Solution: Restriction of null space to $\mathcal{N}_{\mathrm{L}} = \mathrm{span}\{p_n\}_{n=1}^{N_0} \subseteq \mathcal{N}_{\mathrm{L,ext}}$

Fixed quotient space: $\mathcal{H}_Q = \mathcal{H}_{\mathrm{L,ext}}/\mathcal{N}_{\mathrm{L,ext}} = \mathcal{H}_{\mathrm{L}}/\mathcal{N}_{\mathrm{L}}$

Proposition

The native space of L, $\mathcal{H}_L = \mathcal{H}_Q \oplus \mathcal{N}_L$, where \mathcal{N}_L is a **finite-dimensional null space** of L endowed with the inner product $\langle \cdot, \cdot \rangle_{\mathcal{N}_L}$, is a **Hilbert space** for the inner product

$$\langle f, g \rangle_{\mathcal{H}_{\mathbf{L}}} = \langle \mathbf{L}f, \mathbf{L}g \rangle_{L_2} + \langle \operatorname{Proj}_{\mathcal{N}_{\mathbf{L}}} \{f\}, \operatorname{Proj}_{\mathcal{N}_{\mathbf{L}}} \{g\} \rangle_{\mathcal{N}_{\mathbf{L}}}$$

where $\mathrm{Proj}_{\mathcal{N}_L}$ is a projection operator from $\mathcal{H}_L o \mathcal{N}_L.$

29

2.7.1 Hilbert-space structure of null space

■ Motivation: $Ls = w \implies s = L^{-1}w$?

Remove the null-space ambiguity by imposing N_0 linear boundary conditions: $\phi(s)=\mathbf{0}.$

- lacksquare Admissible operator $L:\mathcal{H}_L o L_2(\mathbb{R}^d)$
- lacksquare Null space: $\mathcal{N}_{\mathrm{L}} = \mathrm{span}\{p_n\}_{n=1}^{N_0}$ equipped with a norm $\|\cdot\|_{\mathcal{N}}: \mathcal{H}_{\mathrm{L}} o \mathbb{R}^+$
- lacksquare Linear "boundary" functionals $\phi:\mathcal{H}_{\mathrm{L}} o\mathbb{R}^{N_0}$

$$f \mapsto \phi(f) = (\langle \phi_1, f \rangle, \dots, \langle \phi_{N_0}, f \rangle)$$

Definition

Let $L:\mathcal{H}_L\to L_2(\mathbb{R}^d)$ be a spline-admissible differential operator with a null space \mathcal{N}_L of dimension N_0 . The pair (L,ϕ) is said to be **admissible** if there exist two constants $B\geq A>0$ such that

$$A||f||_{\mathcal{N}} \le ||\phi(f)||_2 \le B(||Lf||_{L_2} + ||f||_{\mathcal{N}}), \quad \forall f \in \mathcal{H}_L.$$

Lower bound: $\phi(q) = \mathbf{0} \quad \Leftrightarrow \quad q = 0 \quad \forall q \in \mathcal{N}_{\mathrm{L}}$

Upper bound \Leftrightarrow $\phi_n \in \mathcal{H}'_{\mathrm{L}}$

Biorthogonal system

- lacksquare Basis for \mathcal{N}_{L} : $oldsymbol{p}=(p_1,\ldots,p_{N_0})$
- lacktriangle Analysis functionals: $\phi = (\phi_1, \dots, \phi_{N_0})$
- Biorthogonality relation

$$\langle \phi_m, p_n \rangle = \delta_{m,n} \quad \Leftrightarrow \quad \phi(p_n) = \mathbf{e}_n \quad \Leftrightarrow \quad \langle \phi, \boldsymbol{p}^T \rangle = \mathbf{I}_{N_0}$$

Projector onto $\mathcal{N}_{\mathrm{L}} = \mathrm{span}\{p_n\}_{n=1}^{N_0} \perp \mathcal{N}_{\mathrm{L}}' = \mathrm{span}\{\phi_n\}_{n=1}^{N_0}$

$$\operatorname{Proj}_{\mathcal{N}_{L}}: \mathcal{H}_{L} \to \mathcal{N}_{L}$$

$$f \mapsto \sum_{n=1}^{N_{0}} \langle \phi_{n}, f \rangle p_{n}$$

31

RKHS associated with null space

■ Biorthogonal system: $\{(\phi_n, p_n)\}_{n=1}^{N_0}$

Proposition

Let (ϕ, p) be a biorthogonal system for $\mathcal{N}_L \subseteq C_{b,\alpha}(\mathbb{R}^d)$. Then, $\mathcal{N}_L = \operatorname{span}\{p_n\}_{n=1}^{N_0}$ equipped with the inner product

$$\langle f, g \rangle_{\mathcal{N}_{L}} = \sum_{n=1}^{N_{0}} \langle \phi_{n}, f \rangle \langle \phi_{n}, g \rangle = \langle \mathcal{R}_{\phi} \{ f \}, g \rangle, \quad f, g \in \mathcal{N}_{L}$$

is a RKHS with reproducing kernel

$$R_{\boldsymbol{p}}\{\delta(\cdot - \boldsymbol{y})\}(\boldsymbol{x}) = \sum_{n=1}^{N_0} p_n(\boldsymbol{x})p_n(\boldsymbol{y}).$$

$$\Leftrightarrow \forall f \in \mathcal{N}_{L}: \|f\|_{\mathcal{N}_{L}} = \|\phi(f)\|_{2}$$

Dual of null space

■ Biorthogonal system: $\{(\phi_n, p_n)\}_{n=1}^{N_0}$

Proposition

The dual of $\mathcal{N}_L = \operatorname{span}\{p_n\}_{n=1}^{N_0}$ is the Hilbert space $\mathcal{N}_L' = \operatorname{span}\{\phi_n = p_n^*\}_{n=1}^{N_0}$ equipped with the inner product

$$\langle f^*, g^* \rangle_{\mathcal{N}'_{\mathbf{L}}} = \sum_{n=1}^{N_0} \langle p_n, f^* \rangle \langle p_n, g^* \rangle = \langle \underbrace{\mathbf{R}_{\mathbf{p}} f^*}_{=f}, g^* \rangle, \quad f^*, g^* \in \mathcal{N}'_{\mathbf{L}}.$$

$$\mathcal{N}_{\mathbf{L}}' = \operatorname{span} \big\{ \phi_n = \mathrm{R}_{\pmb{\phi}} p_n \big\}$$

$$\mathrm{R}_{\pmb{\rho}} : f \mapsto \sum_{n=1}^{N_0} p_n \langle p_n, f \rangle$$

$$\mathrm{R}_{\pmb{\rho}} : f \mapsto \sum_{n=1}^{N_0} p_n \langle p_n, f \rangle$$

$$\mathrm{Riesz\ maps}$$

$$\mathrm{Riesz\ maps}$$

$$\mathrm{Riesz\ maps}$$

$$\mathrm{R}_{\pmb{\phi}} : f \mapsto \sum_{n=1}^{N_0} \phi_n \langle \phi_n, f \rangle$$

$$\mathrm{R}_{\pmb{\phi}} : f \mapsto \sum_{n=1}^{N_0} \phi_n \langle \phi_n, f \rangle$$

Summary of operators

Description	Operator	Kernel
Riesz map $\mathcal{N}_{ m L}' o \mathcal{N}_{ m L}$	R_{p}	$\sum_{n=1}^{N_0} p_n({m x}) p_n({m y})$ reproducing kernel
Riesz map $\mathcal{N}_{\mathrm{L}} o \mathcal{N}_{\mathrm{L}}'$	$\mathrm{R}_{oldsymbol{\phi}}$	$\sum_{n=1}^{N_0} \phi_n(oldsymbol{x}) \phi_n(oldsymbol{y})$
Projector $\mathcal{H}_{\mathrm{L}} o \mathcal{N}_{\mathrm{L}}$	$\mathrm{Proj}_{\mathcal{N}_{\mathrm{L}}}$	$\sum_{n=1}^{N_0} p_n(oldsymbol{x}) \phi_n(oldsymbol{y})$
Projector $\mathcal{H}'_{\mathrm{L}} o \mathcal{N}'_{\mathrm{L}}$	$\mathrm{Proj}_{\mathcal{N}'_{\mathrm{L}}}$	$\sum_{n=1}^{N_0} \phi_n(oldsymbol{x}) p_n(oldsymbol{y})$

Relations

$$\blacksquare \ \mathbf{R}_{\phi}\{p_n\} = p_n^* = \phi_n$$

$$\blacksquare \ \mathrm{R}_{\boldsymbol{p}}\mathrm{R}_{\boldsymbol{\phi}}\{f\} = f \text{ for all } f \in \mathcal{N}_{\mathrm{L}}$$

$$Proj_{\mathcal{N}_L} = (Proj_{\mathcal{N}_r'})^*$$

$$\blacksquare \mathbf{R}_{p}\{\phi_n\} = p_n, \quad n = 1, \dots, N_0$$

$$\blacksquare \ \mathrm{R}_{\pmb{\phi}}\mathrm{R}_{\pmb{p}}\{f^*\} = f^* \text{ for all } f^* \in \mathcal{N}_\mathrm{L}'$$

34

2.7.2 Conditional positive-definiteness

Classical definition

Let $\mathcal{N}=\mathrm{span}\{p_n\}_{n=1}^{N_0}$ be a finite-dimensional subspace of $C_{\mathrm{b},\alpha}(\mathbb{R}^d)$. Then, the kernel function $h:\mathbb{R}^d\times\mathbb{R}^d\to\mathbb{R}$ is said to \mathcal{N} -conditionally positive-definite if

$$\sum_{m=1}^{N} \sum_{n=1}^{N} z_m h(\boldsymbol{x}_m, \boldsymbol{x}_n) z_n \ge 0$$

for any $N\in\mathbb{N}$, $m{x}_1,\dots,m{x}_N\in\mathbb{R}^d$, and $z_1,\dots,z_N\in\mathbb{R}$, subject to the condition

$$\sum_{m=1}^{N} z_m p_n(\boldsymbol{x}_m) = 0$$

for $n = 1, ..., N_0$. The conditional positive-definiteness is said to be **strict** if

$$\sum_{m=1}^{N} \sum_{n=1}^{N} z_m h(x_m, x_n) z_n > 0,$$

under the same conditions with $(z_1,\ldots,z_N)\in\mathbb{R}^N\setminus\{\mathbf{0}\}$ and the \boldsymbol{x}_n all being distinct.

 \Leftrightarrow **p-conditional positivity** where $\mathbf{p}=(p_1,\ldots,p_{N_0})$ represents a basis of $\mathcal N$

35

(Conditionally) positive-definite operator

Definition

Let A be a continuous operator $\mathcal{S}(\mathbb{R}^d) \to \mathcal{S}'(\mathbb{R}^d)$ and \mathcal{N} some finite-dimensional subspace of $\mathcal{S}'(\mathbb{R}^d)$ spanned by $\boldsymbol{p}=(p_1,\ldots,p_{N_0})$. The operator A is said to be:

• Symmetric or self-adjoint if, for all $\varphi_1, \varphi_2 \in \mathcal{S}(\mathbb{R}^d)$,

$$\langle A\varphi_1, \varphi_2 \rangle = \langle A\varphi_2, \varphi_1 \rangle.$$

• Positive-definite if, for any $\varphi \in \mathcal{S}(\mathbb{R}^d)$,

$$\langle A\varphi, \varphi \rangle \ge 0.$$

ullet ${\cal N} ext{-conditionally positive-definite}$ (or ${m p} ext{-conditionally positive}$), if

$$\langle A\varphi, \varphi \rangle \ge 0$$

for any $\varphi \in \mathcal{S}_{p}(\mathbb{R}^{d}) = \mathcal{S}(\mathbb{R}^{d}) \cap \mathcal{N}^{\perp} = \{ \varphi \in \mathcal{S}(\mathbb{R}^{d}) : p(\varphi) = \mathbf{0} \}.$

• Strictly \mathcal{N} -conditionally positive-definite if, for all $\varphi \in \mathcal{S}_{\boldsymbol{p}}(\mathbb{R}^d) \backslash \{0\}$,

$$\langle A\varphi, \varphi \rangle > 0.$$

Positive-definiteness: equivalences

Theorem

$$\mathrm{A}: arphi \mapsto \int_{\mathbb{R}^d} a(\cdot, oldsymbol{y}) arphi(oldsymbol{y}) \mathrm{d} oldsymbol{y} \quad ext{with} \quad a(oldsymbol{x}_0, \cdot) = a(\cdot, oldsymbol{x}_0) \in C_{\mathrm{b}, lpha}(\mathbb{R}^d) ext{ for any } oldsymbol{x}_0 \in \mathbb{R}^d$$

($m{p}$ -conditional) positive definiteness of $\mathrm{A}:\mathcal{S}(\mathbb{R}^d) o\mathcal{S}'(\mathbb{R}^d)$

(*p*-conditional) positive-definiteness of **continuous** kernel $a: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$.

Sketch of proof:

Direct part

Sequence of test functions $\varphi_k=\sum_{n=1}^N z_n u_k(\cdot-{m x}_n)\in \mathcal{S}(\mathbb{R}^d)$ such that

$$\lim_{k \to \infty} \langle f, \varphi_k(\cdot - \boldsymbol{x}_n) \rangle = \sum_{n=1}^{N} z_n f(\boldsymbol{x}_m)$$

$$u_k = k^d e^{-\frac{1}{2}||k\boldsymbol{x}||^2} \xrightarrow[k \to \infty]{} \delta$$

Indirect part

Making the link with Riemann integrals; i.e.,

$$\langle f, \varphi \rangle = \lim_{i \to \infty} \frac{1}{i^d} \sum_{\boldsymbol{m} = -(i^2, \dots, i^2)}^{(i^2, \dots, i^2)} f(\boldsymbol{x_m}) \underbrace{\varphi(\boldsymbol{x_m})}_{z_{\boldsymbol{m}}}, \quad \text{ with } \boldsymbol{x_m} = \frac{m}{i} \in \mathbb{R}^d$$

37

2.7.3 Spline-admissible operator

Definition

A linear operator $L:\mathcal{H}_L\to L_2(\mathbb{R}^d)$ is called **spline-admissible** if there exists a symmetric kernel $G_{L^*L}:\mathbb{R}^d\times\mathbb{R}^d\to\mathbb{R}$, a finite-dimensional subspace $\mathcal{N}_L=\operatorname{span}\{p_n\}_{n=1}^{N_0}$ and an order $\alpha\in\mathbb{R}$ of algebraic growth such that :

1. G_{L^*L} is a **Green's function** of (L^*L) with the property that

$$L\{G_{L^*L}(\cdot, \boldsymbol{y})\}(\boldsymbol{x}) = G_{L^*}(\boldsymbol{x}, \boldsymbol{y}) = G_L(\boldsymbol{y}, \boldsymbol{x})$$
$$L^*L\{G_{L^*L}(\cdot, \boldsymbol{y})\} = L^*\{G_{L^*}(\cdot, \boldsymbol{y})\} = \delta(\cdot - \boldsymbol{y})$$

- 2. Null-space property: $L\{q\} = 0$ for all $q \in \mathcal{N}_L \subseteq \mathcal{H}_L$.
- 3. G_{L^*L} is strictly \mathcal{N}_L -conditionally positive-definite.
- 4. Continuity and polynomial growth: $G_{\mathrm{L^*L}}(\cdot, \boldsymbol{y}_0) \in C_{\mathrm{b},\alpha}(\mathbb{R}^d)$ for any $\boldsymbol{y}_0 \in \mathbb{R}^d$ and $\mathcal{N}_{\mathrm{L}} \subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d)$.
- 5. Boundedness on the diagonal: $\sup_{\boldsymbol{x} \in \mathbb{R}^d} |G_{\mathrm{L^*L}}(\boldsymbol{x}, \boldsymbol{x})| \ (1 + \|\boldsymbol{x}\|)^{2\alpha} < \infty.$

Green's function

Definition

The kernel $G_L : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is a Green's function of L if $L\{G_L(\cdot, \boldsymbol{y})\} = \delta(\cdot - \boldsymbol{y})$.

$$\Leftrightarrow \quad \mathrm{LL^{-1}}\{\varphi\} = \varphi \text{ for any } \varphi \in \mathcal{S}(\mathbb{R}^d) \text{ where } \mathrm{L^{-1}}: \varphi \mapsto \int_{\mathbb{R}^d} G_{\mathrm{L}}(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) \mathrm{d}\boldsymbol{y}$$

lacksquare Preferred scenario: L is LSI with requency response $\widehat{L}(oldsymbol{\omega})$

$$G_{\mathrm{L}}(m{x},m{y}) =
ho_{\mathrm{L}}(m{x}-m{y}) \quad ext{with} \quad
ho_{\mathrm{L}}(m{x}) = \mathcal{F}^{-1}\left\{rac{1}{\widehat{L}(m{\omega})}
ight\}(m{x}) \quad ext{= canonical Green's function}$$

- **Example** of admissible operator: $D^m = \frac{d^m}{dx^m}$ \longleftrightarrow $(j\omega)^m$
 - \blacksquare Green's function (causal): $\mu_m(x) = \frac{x_+^{m-1}}{(m-1)!}$
 - $\blacksquare \text{ Green's function (canonical): } \rho_{\mathcal{D}^m}(x) = \mathcal{F}^{-1}\left\{\frac{1}{(\mathrm{j}\omega)^m}\right\}(x) = \tfrac{1}{2}\mathrm{sign}(x)\frac{x^{m-1}}{(m-1)!}$
 - Null space = polynomials of degree m-1 : $\mathcal{N}_{\mathrm{D}^m} = \mathrm{span}\{\frac{x^{n-1}}{(n-1)!}\}_{n=1}^{N_0}$

$$\mathcal{N}_{\mathbb{D}^m}$$
-conditionally-positive kernel: $G(x,y) = \frac{(-1)^m}{2} \frac{|x-y|^{2m-1}}{(2m-1)!}$

39

2.7.4 RKHS associated with admissible operator

lacksquare "Orthogonal" complement of \mathcal{N}_L in \mathcal{H}_L

$$\mathcal{H}_{\mathrm{L},oldsymbol{\phi}} = \big\{ f \in \mathcal{H}_{\mathrm{L}} : oldsymbol{\phi}(f) = \mathbf{0} \big\},$$

Theorem 3

Let L be an admissible operator and (ϕ, p) a corresponding biorthogonal system for \mathcal{N}_{L} . Then, $\mathcal{H}_{\mathrm{L},\phi}$ is a **Hilbert space** equipped with the **inner product** $\langle f,g\rangle_{\mathrm{L}}=\langle\mathrm{L}\{f\},\mathrm{L}\{g\}\rangle$. Moreover, there exists an isometric map $\mathrm{L}_{\phi}^{-1}:L_2(\mathbb{R}^d)\to\mathcal{H}_{\mathrm{L},\phi}$ such that

$$\mathcal{H}_{L,\phi} = \{ f = L_{\phi}^{-1} w : w \in L_2(\mathbb{R}^d) \}.$$

The operator ${\bf L}_\phi^{-1}$ (see next slide) is uniquely specified through the following properties


- 1. Right-inverse property: $\mathrm{LL}_{m{\phi}}^{-1}w=w$ for all $w\in L_2(\mathbb{R}^d)$
- 2. Boundary conditions: $\phi(\mathbf{L}_{\phi}^{-1}w)=\mathbf{0}$ for all $w\in L_2(\mathbb{R}^d)$.

Construction of stable right-inverse

- Ingredients for construction of $L_{\phi}^{-1}: L_2(\mathbb{R}^d) \to \mathcal{H}_{L,\phi} \subseteq C_{b,\alpha_0}(\mathbb{R}^d)$
 - lacksquare Basis for \mathcal{N}_{L} : $oldsymbol{p}=(p_1,\ldots,p_{N_0})$
 - lacksquare Biorthogonal analysis functionals: $\phi = (\phi_1, \dots, \phi_{N_0})$
 - lacksquare Green's function $G_{\mathrm{L}}:\mathbb{R}^d imes\mathbb{R}^d o\mathbb{R}$ of L $\qquad\Leftrightarrow\qquad \mathrm{L}\{G_{\mathrm{L}}(\cdot,oldsymbol{y})\}=\delta(\cdot-oldsymbol{y})$

$$egin{aligned} \mathrm{L}_{m{\phi}}^{-1}: arphi \mapsto \int_{\mathbb{R}^d} g_{m{\phi}}(\cdot, m{y}) arphi(m{y}) \ g_{m{\phi}}(m{x}, m{y}) = G_{\mathrm{L}}(m{x}, m{y}) - \sum_{n=1}^{N_0} p_n(m{x}) q_n(m{y}) \end{aligned}$$

$$\mathcal{H}_{\mathrm{L}, \phi} \subseteq C_{\mathrm{b}, \alpha_0}(\mathbb{R}^d) \quad \Leftrightarrow \quad \sup_{\boldsymbol{x} \in \mathbb{R}^d} (1 + \|\boldsymbol{x}\|)^{\alpha_0} \|g_{\phi}(\boldsymbol{x}, \cdot)\|_{L_2(\mathbb{R}^d)} < \infty$$



2.7.5 Determination of reproducing kernel

Generalized impulse response of $A_{\phi} = L_{\phi}^{-1} L_{\phi}^{-1*}$

Theorem 4

The **reproducing kernel** of the Hilbert space $\mathcal{H}_{\mathrm{L},\phi}$ specified in Theorem 3 is

$$a_{oldsymbol{\phi}}(oldsymbol{x},oldsymbol{y}) = \int_{\mathbb{R}^d} g_{oldsymbol{\phi}}(oldsymbol{x},oldsymbol{z}) g_{oldsymbol{\phi}}(oldsymbol{y},oldsymbol{z}) \mathrm{d}oldsymbol{z}$$

where $g_{\phi}(\boldsymbol{x}, \boldsymbol{y}) = L_{\phi}^{-1} \{\delta(\cdot - \boldsymbol{y})\}(\boldsymbol{x})$. Moreover, if $G_{L^*L}(\boldsymbol{x}, \boldsymbol{y})$ is the symmetric Green's function of L^*L , then the reproducing kernel can be expressed as

$$a_{\phi}(\boldsymbol{x}, \boldsymbol{y}) = G_{L^*L}(\boldsymbol{x}, \boldsymbol{y}) - \sum_{n=1}^{N_0} p_n(\boldsymbol{x}) v_n(\boldsymbol{y}) - \sum_{n=1}^{N_0} v_n(\boldsymbol{x}) p_n(\boldsymbol{y}) + \sum_{m=1}^{N_0} \sum_{n=1}^{N_0} r_{m,n} p_m(\boldsymbol{x}) p_n(\boldsymbol{y})$$

with

$$v_n(\boldsymbol{y}) = \langle \phi_n, G_{\mathrm{L^*L}}(\cdot, \boldsymbol{y}) \rangle = \int_{\mathbb{R}^d} \phi_n(\boldsymbol{z}) G_{\mathrm{L^*L}}(\boldsymbol{z}, \boldsymbol{y}) \mathrm{d}\boldsymbol{z}$$
$$r_{m,n} = \langle \phi_m \otimes \phi_n, G_{\mathrm{L^*L}} \rangle = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \phi_m(\boldsymbol{x}) \phi_n(\boldsymbol{y}) G_{\mathrm{L^*L}}(\boldsymbol{x}, \boldsymbol{y}) \mathrm{d}\boldsymbol{x} \mathrm{d}\boldsymbol{y}.$$

43

Reproducing kernel and Riesz map

$$a_{\phi}(\boldsymbol{x}, \boldsymbol{y}) = G_{L^*L}(\boldsymbol{x}, \boldsymbol{y}) - \sum_{n=1}^{N_0} p_n(\boldsymbol{x}) v_n(\boldsymbol{y}) - \sum_{n=1}^{N_0} v_n(\boldsymbol{x}) p_n(\boldsymbol{y}) + \sum_{m=1}^{N_0} \sum_{n=1}^{N_0} r_{m,n} p_m(\boldsymbol{x}) p_n(\boldsymbol{y})$$

lacksquare Riesz map $\mathcal{H}'_{\mathrm{L},oldsymbol{\phi}} o\mathcal{H}_{\mathrm{L},oldsymbol{\phi}}$

$$\mathbf{A}_{\phi} = \mathbf{L}_{\phi}^{-1} \mathbf{L}_{\phi}^{-1*} : \varphi \mapsto \int_{\mathbb{D}^d} a_{\phi}(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) d\boldsymbol{y}$$

- Link with unregularized inverse $A: \quad \varphi \mapsto \int_{\mathbb{R}^d} G_{\mathrm{L^*L}}({m x},{m y}) \varphi({m y}) \mathrm{d}{m y}$

 - $r_{m,n} = \langle \phi_m \otimes \phi_n, G_{L^*L} \rangle = \langle A\{\phi_m\}, \phi_n \rangle$

$$(\mathrm{L}^*\mathrm{L})\mathrm{A}\{\varphi\} = (\mathrm{L}^*\mathrm{L})\mathrm{A}_{\pmb{\phi}}\{\varphi\} = \varphi \quad \text{for all } \varphi \in \mathcal{S}(\mathbb{R}^d)$$

but A (unlike A_{ϕ}) does not admit a stable extension $\mathcal{H}'_{L} o \mathcal{H}_{L}$

Characterization of dual space

Proposition

Let L be an admissible operator and (ϕ, p) a corresponding biorthogonal system for \mathcal{N}_L . Then, the continuous dual of the RKHS $\mathcal{H}_{L,\phi}$ is the Hilbert space

$$\mathcal{H}'_{\mathbf{L},\boldsymbol{\phi}} = \{ f^* = \mathbf{L}^* w : w \in L_2(\mathbb{R}^d) \}$$


equipped with the inner product

$$\langle f, g \rangle_{\mathcal{H}'} = \langle \mathcal{L}_{\phi}^{-1*} f, \mathcal{L}_{\phi}^{-1*} g \rangle_{L_2(\mathbb{R}^d)} = \langle \mathcal{A} f^*, g^* \rangle$$

where the operators L_{ϕ}^{-1} and A are as previously defined. The operator (L^*L) is the **Riesz map** $\mathcal{H}_{L,\phi} \to \mathcal{H}'_{L,\phi}$ so that any $f^* = L^*L\{f\} \in \mathcal{H}'_{L,\phi}$ can be viewed as the Riesz conjugate of some corresponding $f \in \mathcal{H}_{L,\phi}$. Finally, we have the "orthogonality" property

$$p(f^*) = 0 \Leftrightarrow p(L^*L\{f\}) = 0$$

for any $f^* \in \mathcal{H}'_{\mathrm{L}, \phi}$ and/or $f \in \mathcal{H}_{\mathrm{L}, \phi}$.



Properties of positive operator $A_{\phi} = L_{\phi}^{-1} L_{\phi}^{-1*}$

Proposition

The operator $A_{\phi}: \varphi \mapsto \int_{\mathbb{R}^d} a_{\phi}(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) d\boldsymbol{y}$, where $a_{\phi}(\cdot, \boldsymbol{y})$ is given in Theorem 3, has the following properties:

- 1. It is the Riesz map $\mathcal{H}'_{\mathrm{L}, \phi} \to \mathcal{H}_{\mathrm{L}, \phi} = \{ f \in \mathcal{H}_{\mathrm{L}} : \phi(f) = \mathbf{0} \}.$
- 2. It is bounded $\mathcal{H}'_{L} \to \mathcal{H}_{L} = \mathcal{H}_{L,\phi} \oplus \mathcal{N}_{L}$.
- 3. It has a finite-dimensional null space $\operatorname{span}\{\phi_n\}_{n=1}^{N_0}=\mathcal{N}'_{\operatorname{L}}.$
- 4. It imposes the boundary conditions: $\phi(A_{\phi}\{f^*\}) = 0$ for all $f^* \in \mathcal{H}'_L$. In other words, A_{ϕ} continuously maps $\mathcal{H}'_L \to \mathcal{H}_{L,\phi}$.
- 5. Relation with unregularized operator $A: \varphi \mapsto \int_{\mathbb{R}^d} G_{L^*L}(\cdot, \boldsymbol{y}) \varphi(\boldsymbol{y}) d\boldsymbol{y}$

$$\forall f^* \in \mathcal{H}'_{\mathrm{L},\phi}: \qquad \mathbf{A}_{\phi}\{f^*\} = \mathbf{A}\{f^*\} - \sum_{n=1}^{N_0} \langle \mathbf{A}\{\phi_n\}, f^* \rangle p_n$$

 $\forall \phi = p^* \in \mathcal{N}_{\mathrm{L}}': \quad \mathrm{A}_{oldsymbol{\phi}}\{\phi\} = 0 \quad \text{ and } \quad \mathrm{A}\{\phi\} \in \mathcal{H}_{\mathrm{L}}$

$$\mathcal{H}'_{L} = \mathcal{H}'_{L,\phi} \oplus \mathcal{N}'_{L} \qquad \mathcal{H}_{L} = \mathcal{H}_{L,\phi} \oplus \mathcal{N}_{L}$$

$$\mathcal{H}'_{L,\phi} \qquad \mathcal{H}_{L,\phi}$$

$$\{0\} \qquad \qquad (L^*L) \qquad \qquad \mathcal{H}_{L,\phi}$$

$$\text{span}\{\phi_n\}_{n=1}^{N_0} \qquad \text{span}\{p_n\}_{n=1}^{N_0}$$

RHKS structure of native space \mathcal{H}_{L}

Theorem 5

Any $f \in \mathcal{H}_{\mathrm{L}} = \mathcal{H}_{\mathrm{L}, oldsymbol{\phi}} \oplus \mathcal{N}_{\mathrm{L}}$ has a unique representation as

$$f = L_{\phi}^{-1} w + q$$

where $w = Lf \in L_2(\mathbb{R}^d)$ and $q = \operatorname{Proj}_{\mathcal{N}_L}\{f\} = \sum_{n=1}^{N_0} \langle f, \phi_n \rangle p_n \in \mathcal{N}_L$. Moreover, \mathcal{H}_L equipped with the inner product

$$\langle f, g \rangle_{\mathcal{H}_{L}} = \langle Lf, Lg \rangle + \sum_{n=1}^{N_0} \langle \phi_n, f \rangle \langle \phi_n, g \rangle$$

is a RKHS whose reproducing kernel is

$$h_{\boldsymbol{\phi}}(\boldsymbol{x}, \boldsymbol{y}) = a_{\boldsymbol{\phi}}(\boldsymbol{x}, \boldsymbol{y}) + \sum_{n=1}^{N_0} p_n(\boldsymbol{x}) p_n(\boldsymbol{y}).$$

Finally, if the operator A meets the admissibility conditions (strict conditional positivity and α -boundedness) and the ϕ_n are such that $A\{\phi_n\} \in C_{b,\alpha}(\mathbb{R}^d)$, then $\mathcal{S}(\mathbb{R}^d) \subseteq \mathcal{H}_L \subseteq C_{b,\alpha}(\mathbb{R}^d) \subseteq \mathcal{S}'(\mathbb{R}^d)$ with the insurance that \mathcal{H}_L is **dense** in $\mathcal{S}'(\mathbb{R}^d)$.

49

Summary of operators

Description	Operator	Kernel	
Right-inverse of ${f L}$	G	$G_{\mathrm{L}}(\boldsymbol{x}, \boldsymbol{y}) = \mathrm{L}\{G_{\mathrm{L^*L}}(\cdot, \boldsymbol{y})\}$	$\overline{(x)}(y)$???
Right-inverse of $\left(L^*L\right)$	$A=GG^*$	$G_{\mathrm{L^*L}}(oldsymbol{x},oldsymbol{y})$	
Stable inverse of \boldsymbol{L}	$\mathcal{L}_{m{\phi}}^{-1}$	$G_{\mathrm{L}}(oldsymbol{x},oldsymbol{y}) - \sum_{n=1}^{N_0} p_n(oldsymbol{x}) \mathrm{G}^*\{\phi_n\}(oldsymbol{y})$	
Riesz map $\mathcal{H}'_{\mathrm{L},oldsymbol{\phi}} o \mathcal{H}_{\mathrm{L},oldsymbol{\phi}}$	${ m A}_{m{\phi}}$	$a_{oldsymbol{\phi}}(oldsymbol{x},oldsymbol{y})$ rep	roducing kernel
Riesz map $ {\cal H}_{{ m L},\phi} o {\cal H}'_{{ m L},\phi} $	(L^*L)		
Riesz map $ {\cal H}'_{ m L} ightarrow {\cal H}_{ m L} $	$\mathrm{A}_{m{\phi}} + \mathrm{R}_{m{p}}$	$a_{oldsymbol{\phi}}(oldsymbol{x},oldsymbol{y}) + \sum_{n=1}^{N_0} p_n(oldsymbol{x}) p_n(oldsymbol{y})$	reproducing kernel
Riesz map $ {\cal H}_{ m L} ightarrow {\cal H}_{ m L}'$	$(L^*L) + R_{\phi}$		

■ Direct sum decomposition

$$\quad \blacksquare \ \forall \phi = p^* \in \mathcal{N}'_{\mathbf{L}}: \quad \mathbf{A}_{\phi}\{\phi\} = 0$$

$$\forall p \in \mathcal{N}_{L}: \qquad (L^*L)\{p\} = 0$$

$$\forall \tilde{f}^* \in \mathcal{H}'_{L, \phi}: \qquad \mathbf{R}_{p}\{\tilde{f}^*\} = 0$$

$$\forall \tilde{f} \in \mathcal{H}_{L,\phi} : \quad R_{\phi} \{ \tilde{f} \} = 0$$

Summary of key relations

$$\mathcal{L}_{\phi}^{-1}: \mathcal{L}_{2}(\mathbb{R}^{d}) \to \mathcal{H}_{\mathcal{L},\phi} \subseteq \mathcal{H}_{\mathcal{L}}$$

$$\forall w \in L_2(\mathbb{R}^d): LL_{\phi}^{-1}\{w\} = w,$$

$$\forall w \in L_2(\mathbb{R}^d): \quad \phi(\mathbf{L}_{\phi}^{-1}w) = \mathbf{0}$$

$$L_{\phi}^{-1*}: \mathcal{H}'_{L,\phi} \to L_2(\mathbb{R}^d)$$

$$\blacksquare L_{\phi}^{-1*} = G^*(\operatorname{Id} - \operatorname{Proj}_{\mathcal{N}_{r}'})$$

$$\forall w \in L_2(\mathbb{R}^d): \quad L_{\boldsymbol{\phi}}^{-1*}L^*\{w\} = w$$

$$\forall \phi \in \mathcal{N}'_{L}: \qquad L_{\phi}^{-1*} \{\phi\} = 0$$

Riesz map
$$A_{\phi}: \mathcal{H}'_{L,\phi} o \mathcal{H}_{L,\phi} \subseteq \mathcal{H}_{L}$$

$$\blacksquare A_{\phi} = (\operatorname{Id} - \operatorname{Proj}_{\mathcal{N}_{L}})(GG^{*})(\operatorname{Id} - \operatorname{Proj}_{\mathcal{N}_{L}'}) = L_{\phi}^{-1}L_{\phi}^{-1*}$$

$$\forall \phi \in \mathcal{N}'_{L}: \quad A_{\phi}\{\phi\} = 0$$

$$\forall f^* \in \mathcal{H}'_{\mathrm{L}}: \quad \phi(\mathrm{A}_{\phi}\{f^*\}) = \mathbf{0}$$

$$\forall \tilde{f}^* \in \mathcal{H}'_{L,\phi} : (L^*L)A_{\phi}\{\tilde{f}^*\} = \tilde{f}^*$$

$$\forall \tilde{f}^* \in \mathcal{H}'_{L,\phi} : \quad A_{\phi}\{\tilde{f}^*\} = A\{\tilde{f}^*\} + \sum_{n=1}^{N_0} \langle A\{\phi_n\}, \tilde{f}^* \rangle p_n$$

51

Native space: the bottom line

■ The native space of L is a **Hilbert space** $\mathcal{H}_L \subseteq C_{b,\alpha}(\mathbb{R}^d)$ equipped with the inner product

$$\langle f, g \rangle_{\mathcal{H}_{L}} = \langle Lf, Lg \rangle + \sum_{n=1}^{N_{0}} \langle \phi_{n}, f \rangle \langle \phi_{n}, g \rangle$$

with $\phi_n \in \mathcal{H}_L'$ where $\left(\phi, \boldsymbol{p}\right) = \left((\phi_n), (p_n)\right)$ is a biothonormal system for $\mathcal{N}_L = \operatorname{span}\{p_n\}_{n=1}^{N_0} \subseteq \mathcal{H}_L$.

- Direct sum structure: $\mathcal{H}_L = \mathcal{H}_{L,\phi} \oplus \mathcal{N}_L = \mathcal{H}_{L,\tilde{\phi}} \oplus \mathcal{N}_L$ where $\mathcal{H}_{L,\phi} = \{f \in \mathcal{H}_L : \phi(f) = \mathbf{0}\}$ with the norms induced by the **biorthogonal systems** (ϕ, p) and $(\tilde{\phi}, \tilde{p})$ being equivalent.
- If there exists a symmetric \mathcal{N}_L -conditionally-positive operator $A: \mathcal{S}(\mathbb{R}^d) \to \mathcal{S}'(\mathbb{R}^d)$ such that $(L^*L)A\{\varphi\} = \varphi$ for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$, then:
 - lacksquare \mathcal{H}_{L} can be defined as the completion of $\mathcal{S}(\mathbb{R}^d)$ is the $\|\cdot\|_{\mathcal{H}_{\mathrm{L}}}$ -norm.
 - lacksquare \mathcal{H}_L is **dense** in $\mathcal{S}'(\mathbb{R}^d)$.
 - Any function $f \in C_{b,\alpha}(\mathbb{R}^d)$ can be approximated as closely as desired by a **kernel** expansion of the form

$$f_K = \sum_{k=1}^{K} a_k G_{\mathrm{L^*L}}(\cdot, \boldsymbol{x}_k) + \sum_{n=1}^{N_0} b_n p_n$$

with a finite number $(K+N_0)$ of terms and some (possibly adaptive) centers $oldsymbol{x}_k \in \mathbb{R}^d$.

CONTENT

- 1. Preliminaries
- 2. Reproducing kernel Hilbert spaces
- 3. Variational splines and representer theorems
 - Regularization induced by an inner product
 - Non-coercive regularization functionals
 - Discretization and numerical solutions
- 4. Gaussian processes

53

3. VARIATIONAL SPLINES

■ Generalized interpolation problem

Recover $f \in \mathcal{H}_{\mathrm{L}}$ from the measurements $y_m = \langle \nu_m, f \rangle, (m = 1, \dots, M)$

Traditional setting: $y_m = f(x_m) = \langle \delta(\cdot - x_m), f \rangle$

■ Variational (=minimum energy) solution

$$f_0 = \arg\min_{f \in \mathcal{H}_L} \|Lf\|_{L_2}^2$$
 s.t. $(f(x_m) = y_m)_{m=1}^M$

■ Minimization of least-squares functional

Perturbed measurements model: $y_m = \langle \nu_m, f \rangle + \epsilon_m \quad (m=1,\dots,M)$

$$J_{\mathrm{LS}}(f|\mathbf{y},\lambda) = \sum_{m=1}^{M} (y_m - \langle \nu_m, f \rangle)^2 + \lambda ||\mathbf{L}f||_{L_2}^2$$

3.1 Regularization induced by an inner product

Optimization is performed over a Hilbert space ${\cal H}$

Effect of measurement functional $\nu_m \in \mathcal{H}'$: $\langle \nu_m, f \rangle = \langle \nu_m^*, f \rangle_{\mathcal{H}}$

$$J_{LS}(f|\mathbf{y},\lambda) = \sum_{m=1}^{M} (y_m - \langle \nu_m^*, f \rangle_{\mathcal{H}})^2 + \lambda \|f\|_{\mathcal{H}}^2 = \frac{1}{2}a(f,f) - v(f) + C_0$$

Linear functional:
$$v(f) = 2\sum_{m=1}^{M} y_m \langle \nu_m^*, f \rangle_{\mathcal{H}}$$

Bilinear form:
$$a(f_1, f_2) = 2\lambda \langle f_1, f_2 \rangle_{\mathcal{H}} + 2\sum_{m=1}^{M} \langle \nu_m^*, f_1 \rangle_{\mathcal{H}} \langle \nu_m^*, f_2 \rangle_{\mathcal{H}}$$

$$f_0 = \arg\min_{f \in \mathcal{H}} J(f|\mathbf{y}, \lambda) \quad \Leftrightarrow \quad v(f) = a(f_0, f) \quad \text{(Condition of optimality)}$$

$$\Rightarrow \qquad f_0 \in \operatorname{span}\{\nu_m^*\}_{m=1}^M$$

with guarantee of unicity (Hilbert's projection theorem)

55

Regularized least-squares fit: discretization

 \mathcal{H} : RKHS with reproducing kernel $r_{\mathcal{H}}: \mathbb{R}^d \times \mathbb{R}^d o \mathbb{R}$

Solution space parametrized by $\mathbf{a} = (a_1, \dots, a_M) \in \mathbb{R}^M$

$$f(\boldsymbol{x}) = \sum_{m=1}^{M} a_m \nu_m^*(\boldsymbol{x}), \qquad \nu_m^*(\boldsymbol{x}) = \int_{\mathbb{R}^d} r_{\mathcal{H}}(\boldsymbol{x}, \boldsymbol{y}) \nu_m(\boldsymbol{y}) d\boldsymbol{y}$$

$$\Rightarrow J_{LS}(f|\mathbf{y}, \lambda) = J(\mathbf{a}|\mathbf{y}, \lambda) = \|\mathbf{y} - \mathbf{G}\mathbf{a}\|_{2}^{2} + \lambda \mathbf{a}^{T}\mathbf{G}\mathbf{a}$$

Condition for optimality:
$$\frac{\partial J(\mathbf{a}|\mathbf{y},\lambda)}{\partial \mathbf{a}} = 2\mathbf{G}\big(\mathbf{y} - (\mathbf{G} + \lambda \mathbf{I}_M)\mathbf{a}\big) = 0$$
 + unicity

$$\Rightarrow$$
 $\mathbf{y} = (\mathbf{G} + \lambda \mathbf{I}_M)\mathbf{a} \Leftrightarrow \mathbf{a} = (\mathbf{G} + \lambda \mathbf{I}_M)^{-1}\mathbf{y}$

Generalization: Abstract representer theorem

- lacksquare \mathcal{H} : Hilbert space with inner product $\langle\cdot,\cdot\rangle_{\mathcal{H}}$ and Riesz map $R:\mathcal{H}'\to\mathcal{H}$
- $\mathbf{v}: f \mapsto \boldsymbol{\nu}(f) = (\langle \nu_1, f \rangle, \dots, \langle \nu_M, f \rangle)$ is a continuous linear operator $\mathcal{H}_L \to \mathbb{R}^M$ that extracts M measurements from the signal f;
- lacktriangleright \mathcal{C} : closed convex subset of \mathbb{R}^M such that its preimage in \mathcal{H} is nonempty.

Theorem

The solution of the general convex optimization problem

$$\arg\min_{f\in\mathcal{H}}\|f\|_{\mathcal{H}}^2$$
 s.t. $\boldsymbol{
u}(f)\in\mathcal{C}$

is unique and of the form

$$f_0 = \sum_{m=1}^M a_m \nu_m^*$$

with $\nu_m^* = \mathbb{R}\{\nu_m\} \in \mathcal{H}$ and suitable linear weights $\mathbf{a} = (a_m) \in \mathbb{R}^M$.

Special case: $\mathcal{C}_{\mathrm{LS}} = \{\mathbf{z} \in \mathbb{R}^d : \|\mathbf{z} - \mathbf{y}\|_2^2 \leq \sigma^2\}$

57

Proof (step by step)

- Unicity of the solution (by Hilbert's projection theorem) \mathcal{C} convex & closed, $\boldsymbol{\nu}$ linear & continuous \Rightarrow $\mathcal{U} = \boldsymbol{\nu}^{-1}(\mathcal{C})$ convex & closed
- $\blacksquare \ \ \text{Riesz' representation theorem:} \quad \langle \nu_m, f \rangle = \langle \nu_m^*, f \rangle_{\mathcal{H}} \ \ \text{with} \ \nu_m^* = \mathrm{R}\{\nu_m\} \in \mathcal{H}$
- Identification of reconstruction space: $\mathcal{V} = \operatorname{span}\{\nu_m^*\}_{m=1}^M \subseteq \mathcal{H}$
- Optimization space: $\mathcal{H} = \mathcal{V} \oplus \mathcal{V}^{\perp}$ $\Rightarrow \quad f = u + u^{\perp} \text{ with } u \in \mathcal{V} \text{ and } u^{\perp} \in \mathcal{V}^{\perp}$
- Null space of $\boldsymbol{\nu}$: $\mathcal{V}^{\perp}=\{f\in\mathcal{H}:\langle f,\nu_m\rangle=\langle f,\nu_m^*\rangle_{\mathcal{H}}=0,\ m=1,\ldots,M\}$ $\Rightarrow \quad f_0=u_0+u_0^{\perp} \text{ with } \boldsymbol{\nu}(f_0)=\boldsymbol{\nu}(u_0) \text{ and } \boldsymbol{\nu}(u_0^{\perp})=0$
- f_0 is the minimal-norm solution $\|f_0\|_{\mathcal{H}}^2 = \|u_0 + u_0^\perp\|_{\mathcal{H}}^2 \le \|u_0\|_{\mathcal{H}}^2 \quad \Leftrightarrow \quad u_0^\perp = 0 \qquad \qquad \Leftrightarrow \quad f_0 \in \mathcal{V} \qquad \qquad \mathsf{QED}$

Smoothing splines and/or ridge regression

 \mathcal{H} : RKHS with reproducing kernel $r_{\mathcal{H}}: \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$

"Ideal" sampling functionals: $\nu_m = \delta(\cdot - \boldsymbol{x}_m), \quad (m = 1, \dots, M)$

$$f_{\text{LS}} = \arg\min_{f \in \mathcal{H}} \left(\sum_{m=1}^{M} (y_m - f(\boldsymbol{x}_m))^2 + \lambda \|f\|_{\mathcal{H}}^2 \right)$$

Solution space parametrized by $\mathbf{a} = (a_1, \dots, a_M) \in \mathbb{R}^M$

Regularized least-squares fit

autocorrelation ridge \Rightarrow $\mathbf{a}_{LS} = (\mathbf{R} + \lambda \mathbf{I}_M)^{-1} \mathbf{y}$

Statistical linear regression: $\lambda \propto \sigma^2$

with
$$[\mathbf{R}]_{m,n} = \langle r_{\mathcal{H}}(\cdot, \boldsymbol{x}_m), r_{\mathcal{H}}(\cdot, \boldsymbol{x}_n) \rangle_{\mathcal{H}} = r_{\mathcal{H}}(\boldsymbol{x}_m, \boldsymbol{x}_n)$$

59

Application to machine learning

- The "machine learning" problem
 - Find a function $f: \mathbb{R}^d \to \mathbb{R}$ such that $f(x_m) \approx y_m$ (with a **training set** of size M)
 - Proximity between $\mathbf{z}(f) = (f(x_1), \dots, f(x_M))$ and $\mathbf{y} = (y_1, \dots, y_M)$ is measured by some loss function $F: \mathbb{R}^M \times \mathbb{R}^M \to \mathbb{R}$

Definition

A multivariate function $g: \mathbb{R}^M \to \mathbb{R}$ is **convex** if

$$g(\tau \mathbf{z}_1 + (1-\tau)\mathbf{z}_2) \le \tau g(\mathbf{z}_1) + (1-\tau)g(\mathbf{z}_2, \mathbf{y})$$

for all $\mathbf{z}_1, \mathbf{z}_2 \in \mathbb{R}^M$ and all $\tau \in [0,1]$. It is **strictly convex** if the order relation holds with a strict inequality for $\mathbf{z}_1 \neq \mathbf{z}_2$ and $\tau \in (0,1)$.

A multivariate function $g: \mathbb{R}^M \to \mathbb{R}$ is said to be **coercive** if $\lim_{\|\mathbf{z}\| \to \infty} g(\mathbf{z}) = \infty$.

The loss function $F: \mathbb{R}^M \times \mathbb{R}^M \to \mathbb{R}$ is said to be convex (resp., coercive) if $F(\cdot, \mathbf{y}) : \mathbb{R}^M \to \mathbb{R}$ is convex (resp., coercive) for any fixed $\mathbf{y} \in \mathbb{R}^d$

Examples of loss functions

61

Representer theorem for machine learning

- $\blacksquare \ \mathcal{H} \subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d) \text{: RKHS with inner product } \langle \cdot, \cdot \rangle_{\mathcal{H}} \text{ and reproducing kernel } r_{\mathcal{H}} : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$
- \blacksquare Sampling operator $f\mapsto \mathbf{z}(f)=ig(f(m{x}_1),\dots,f(m{x}_M)ig)$ with $m{x}_1,\dots,m{x}_M\in\mathbb{R}^d$
- lacksquare Corresponding data values $\mathbf{y}=(y_1,\ldots,y_M)\in\mathbb{R}^M$

Theorem

The solution of the generic minimization problem

$$\arg\min_{f\in\mathcal{H}} \left(F(\mathbf{z}(f), \mathbf{y}) + \lambda ||f||_{\mathcal{H}}^{2} \right),$$

where the cost function F is strictly convex, is unique and of the form

$$f(oldsymbol{x}) = \sum_{m=1}^{M} a_m r_{\mathcal{H}}(oldsymbol{x}, oldsymbol{x}_m)$$

with suitable linear weights $\mathbf{a} = (a_m) \in \mathbb{R}^M$.

Theoretical justification for kernel methods

(Schoenberg 1964, Kimeldorf-Wahba 1971) (Schölkopf-Smola 2001)

$$F(\mathbf{z}(f), \mathbf{y}) = \|\mathbf{z}(f) - \mathbf{y}\|_2^2$$

Proof

- $lackbox{}{} f \mapsto \mathbf{z}(f)$: linear and continuous on \mathcal{H}
- $f \mapsto F(\mathbf{z}(f), \mathbf{y})$: strictly convex and continuous on \mathcal{H}
- $lacksquare\ f\mapsto \lambda \|f\|_{\mathcal{H}}^2$: strictly convex, continuous and trivially coercive on \mathcal{H}

Theorem (Classical existence result from convex analysis)

Let \mathcal{X} be a reflexive Banach space and $J: \mathcal{X} \to \mathbb{R}$ a **convex**, (lower semi-) continuous, and coercive functional on \mathcal{X} . Then, the problem

$$S = \arg\inf_{x \in \mathcal{X}} J(x)$$

admits at least one solution. Moreover, the solution set S is a convex subset of $\mathcal X$ which reduces to a single point (unique solution) when J is **strictly convex**.

 \Rightarrow Unique solution f_0 with "optimal" data term $F(\mathbf{y}_0,\mathbf{y})$ where $\mathbf{y}_0=\mathbf{z}(f_0)\in\mathbb{R}^M$

Since f_0 achieves the minimum, it is also the solution of the interpolation problem

$$f_0 = \min_{f \in \mathcal{H}} \|f\|_{\mathcal{H}}^2$$
 s.t. $\mathbf{z}(f) = \mathbf{y}_0$

$$\Rightarrow f_0 \in \operatorname{span}\{r_{\mathcal{H}}(\cdot, \boldsymbol{x}_m)\}_{m=1}^M$$

63

3.2 Non-coercive regularization functionals

- Functional context
 - \blacksquare Non-trivial null space $\mathcal{N}_{\mathrm{L}} = \{f \in \mathcal{H}_{\mathrm{L}} : \mathrm{L}\{f\} = 0\}$
 - \blacksquare Biorthogonal system (ϕ, p) for $\mathcal{N}_{\mathrm{L}} = \mathrm{span}\{p_n\}_{n=1}^{N_0}$ with $\phi_n \in \mathcal{H}_{\mathrm{L}}'$
 - Native space $\mathcal{H}_{\mathrm{L}}\subseteq C_{\mathrm{b},\alpha}(\mathbb{R}^d)$ is the Hilbert space equipped with the composite norm $\|f\|_{\mathcal{H}_{\mathrm{L}}}=\sqrt{\|\mathrm{L}f\|_{L_2}^2+\|\phi(f)\|_2^2}$
- Variational formulation of linear inverse problems

$$\arg\min_{f\in\mathcal{H}_{\mathrm{L}}}\left(F\left(\boldsymbol{\nu}(f),\mathbf{y}\right)+\lambda\|\mathrm{L}f\|_{L_{2}(\mathbb{R}^{d})}^{2}\right)$$

- Motivation: favors ("smooth") solution with large null-space component
- \blacksquare Theoretical challenge: $\|\mathcal{L}f\|_{L_2}^2$ is only a semi-norm !

Proper regularization operator

Definition

The operator $L:\mathcal{H}_L\to L_2(\mathbb{R}^d)$ with null space $\mathcal{N}_L=\operatorname{span}\{p_n\}_{n=1}^{N_0}\subseteq\mathcal{H}_L$ is a proper regularization operator for the measurement operator $\boldsymbol{\nu}:f\mapsto \boldsymbol{\nu}(f)=(\langle \nu_1,f\rangle,\ldots,\langle \nu_M,f\rangle)$ if the following technical conditions are met:

- 1. L is spline-admissible
- 2. $\nu_1, \ldots, \nu_M \in \mathcal{H}'_{L}$
- 3. For all $q \in \mathcal{N}_L$, $\|\boldsymbol{\nu}(q)\|_2 \geq 0$ with equality if and only if q = 0.

$$\Leftrightarrow \quad \exists c>0 \quad \text{s.t.} \quad \|q\|_{\mathcal{N}_{\mathbf{L}}}^2 = \|\phi(q)\|_2^2 \leq \tfrac{1}{c} \|\boldsymbol{\nu}(q)\|_2^2.$$

Criterion

The singular values of $\mathbf{P} = [\boldsymbol{\nu}(p_1) \cdots \boldsymbol{\nu}(p_{N_0})]$ are strictly positive and bounded

65

Representer theorem for linear inverse problems

- ν : $f \mapsto \nu(f) = (\langle \nu_1, f \rangle, \dots, \langle \nu_M, f \rangle)$ is a continuous linear operator $\mathcal{H}_L \to \mathbb{R}^M$ that extracts M measurements from the signal f;
- ${f L}:{\cal H}_{
 m L} o L_2({\Bbb R}^d)$ is a proper regularization operator;
- $\P \{p_n\}_{n=1}^{N_0}$ is a basis of the null space of the regularization operator;
- lacksquare $F: \mathbb{R}^M imes \mathbb{R}^M o \mathbb{R}$ is a strictly convex and coercive loss function;
- $\mathbf{y} \in \mathbb{R}^M$ is a given data vector and $\lambda \in \mathbb{R}^+$ an adjustable regularization parameter.

Theorem

The solution of the general minimization problem

$$\arg\min_{f\in\mathcal{H}_{\mathrm{L}}}F(\boldsymbol{\nu}(f),\mathbf{y})+\lambda\|\mathrm{L}f\|_{L_{2}(\mathbb{R}^{d})}^{2}$$

is unique and has the generic linear parametric form

$$f_{(\lambda)} = \sum_{m=1}^{M} a_m \varphi_m + \sum_{n=1}^{N_0} b_n p_n$$

with $\varphi_m = A\{\nu_m\} = \int_{\mathbb{R}^d} G_{L^*L}(\cdot, \boldsymbol{y}) \nu_m(\boldsymbol{y}) d\boldsymbol{y}, \quad \mathbf{a} = (a_m) \in \mathbb{R}^M, \quad \mathbf{b} = (b_n) \in \mathbb{R}^{N_0},$ subject to the "orthogonality" constraint: $\langle \mathbf{a}, \boldsymbol{\nu}(p_n) \rangle = 0$ for $n = 1, \dots, N_0$.

Proof

Minimization of
$$J(f|\mathbf{y},\lambda) = F(\boldsymbol{\nu}(f),\mathbf{y}) + \lambda \|\mathbf{L}f\|_{L_2(\mathbb{R}^d)}^2$$

$$\mathcal{H}_{\mathbf{L}} = \mathcal{H}_{\mathbf{L},\boldsymbol{\phi}} \oplus \mathcal{H}_{\mathbf{L}} \quad \Leftrightarrow \quad f = \mathbf{L}_{\boldsymbol{\phi}}^{-1}w + q \quad \text{with} \quad w = \mathbf{L}f, \, q = \mathrm{Proj}_{\mathcal{N}_{\mathbf{L}}}\{f\}$$

- Step 1: Establish existence and unicity
 - Continuity of J: directly follows from continuity of $f\mapsto \|\mathrm{L} f\|_{L_2}^2$, $m{
 u}$, and $F(\cdot,\mathbf{y})$.
 - Strict-convexity of J: follows from strict convexity of J and $F(\cdot, \mathbf{y})$, and linearity of ν .

Key relations: $\|f\|_{\mathcal{H}_1}^2 = \|w\|_{L_2}^2 + \|\phi(q)\|_2^2$ and $c\|\phi(q)\|_2 \leq \|\nu(q)\|_2$

67

Proof (Cont.)

Step 2: Generic form of solution

Minimizer:
$$f_0$$
 such that $\boldsymbol{\nu}(f_0) = \mathbf{y}_0 \in \mathbb{R}^M$ and $\boldsymbol{\phi}(f_0) = \mathbf{c}_0 = (c_1, \dots, c_{N_0}) \in \mathbb{R}^{N_0}$

Direct sum decomposition in $\mathcal{H}_{\mathrm{L}}=\mathcal{H}_{\mathrm{L},\phi}\oplus\mathcal{H}_{\mathrm{L}}$

$$\begin{split} f_0 &= \tilde{f}_0 + q_0 \quad \text{with} \quad \tilde{f}_0 = f_0 - q_0 \in \mathcal{H}_{\mathrm{L}, \phi}, \quad q_0 = \mathrm{Proj}_{\mathcal{N}_{\mathrm{L}}} \{ f_0 \} \in \mathcal{N}_{\mathrm{L}} \\ \text{subject to } \phi(\tilde{f}_0) &= \mathbf{0} \quad \Rightarrow \quad \phi(q_0) = \mathbf{c}_0, \quad \text{and} \quad \boldsymbol{\nu}(\tilde{f}_0) = \mathbf{y}_0 - \boldsymbol{\nu}(q_0) \end{split}$$

$$\Rightarrow \quad q_0 = \sum_{n=1}^{N_0} c_n p_n \quad \text{ and } \quad \tilde{f}_0 = \arg\min_{\tilde{f} \in \mathcal{H}_{\mathrm{L}, \phi}} \|\tilde{f}\|_{\mathcal{H}_{\mathrm{L}, \phi}}^2 \quad \text{s.t. } \quad \boldsymbol{\nu}(\tilde{f}) = \mathbf{y}_0 - \boldsymbol{\nu}(q_0)$$

$$\Rightarrow \quad f_0 = \sum_{m=1}^M a_m \mathbf{A}_{\pmb{\phi}} \{ \nu_m \} + \sum_{n=1}^{N_0} c_n p_n \qquad \qquad \text{(by Abstract representer theorem)}$$

$$lacksquare$$
 Orthogonality property: $ilde{f}_0^* = (\mathrm{L}^*\mathrm{L})\{ ilde{f}_0\} = \sum_{m=1}^M a_m \nu_m \in \mathcal{H}'_{\mathrm{L},oldsymbol{\phi}} \perp \mathcal{N}_{\mathrm{L}}$

$$f_0 = A_{\phi} \left\{ \sum_{m=1}^{M} a_m \nu_m \right\} + \sum_{n=1}^{N_0} c_n p_n = A \left\{ \sum_{m=1}^{M} a_m \nu_m \right\} + \sum_{n=1}^{N_0} b_n p_n$$

(using relation between A_{ϕ} and A for $f^* \in \mathcal{H}'_{\mathrm{L},\phi}$)

3.3 Discretization and numerical solution

Finite-dimensional solution space

$$\begin{split} \mathcal{V}_{\mathrm{L},\nu} &= \{ f = \boldsymbol{\varphi}^T \mathbf{a} + \boldsymbol{p}^T \mathbf{b} : \mathbf{a} \in \mathbb{R}^M, \mathbf{b} \in \mathbb{R}^{N_0} \} \subseteq \mathcal{H}_{\mathrm{L}} \\ \text{where } \boldsymbol{\varphi} &= (\varphi_1, \dots, \varphi_M) \text{ with } \varphi_n = \mathrm{A}\{\nu_n\} \quad \text{ and } \quad \boldsymbol{p} = (p_1, \dots, p_{N_0}) \\ \Rightarrow \quad J(f|\mathbf{y}, \lambda) &= J(\mathbf{a}, \mathbf{b}|\mathbf{y}, \lambda) = F(\mathbf{G}\mathbf{a} + \mathbf{P}\mathbf{b}, \mathbf{y}) + \lambda \mathbf{a}^T \mathbf{G}\mathbf{a} \end{split}$$

Exact discretization

$$\begin{aligned} \text{System/Gram matrix } \mathbf{G} \in \mathbb{R}^{M \times M} \colon \quad [\mathbf{G}]_{m,n} &= \langle \nu_m, \varphi_n \rangle = \langle \nu_m, \mathbf{A}\{\nu_n\} \rangle = \langle \varphi_m, \varphi_n \rangle_{\mathcal{H}} \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \nu_m(\boldsymbol{x}) G_{\mathrm{L^*L}}(\boldsymbol{x}, \boldsymbol{y}) \nu_n(\boldsymbol{y}) \mathrm{d}\boldsymbol{x} \mathrm{d}\boldsymbol{y} \end{aligned}$$

$$\text{Additional system matrix } \mathbf{P} \in \mathbb{R}^{M \times N_0} \colon \quad [\mathbf{P}]_{m,n} = \langle \nu_m, p_n \rangle = \int_{\mathbb{R}^d} \nu_m(\boldsymbol{x}) p_n(\boldsymbol{x}) \mathrm{d}\boldsymbol{x}$$

69

Discrete optimization problem

$$\arg\min_{\mathbf{a} \in \mathbb{R}^{M}, \, \mathbf{b} \in \mathbb{R}^{N_{0}}} \left\{ F(\mathbf{G}\mathbf{a} + \mathbf{P}\mathbf{b}, \mathbf{y}) + \lambda \, \mathbf{a}^{T}\mathbf{G}\mathbf{a} \right\}$$

Steepest descent algorithm

Initialization: $(\mathbf{a}_0, \mathbf{b}_0)$

$$\mathbf{a}_{k+1} = \mathbf{a}_k - \tau \mathbf{G} \left(\nabla F(\mathbf{G} \mathbf{a}_k + \mathbf{P} \mathbf{b}_k, \mathbf{y}) + 2\lambda \mathbf{a}_k \right)$$
$$\mathbf{b}_{k+1} = \mathbf{b}_k - \tau \mathbf{P}^T \nabla F(\mathbf{G} \mathbf{a}_k + \mathbf{P} \mathbf{b}_k, \mathbf{y})$$

Until stop criterion

Least-squares approximation problem

$$\begin{split} J(\mathbf{a},\mathbf{b}|\mathbf{y},\lambda) &= \|\mathbf{G}\mathbf{a} + \mathbf{P}\mathbf{b} - \mathbf{y}\|_2^2 + \lambda \, \mathbf{a}^T \mathbf{G}\mathbf{a} \\ &\frac{\partial J(\mathbf{a},\mathbf{b}|\mathbf{y},\lambda)}{\partial \mathbf{a}} = 2\mathbf{G}\Big((\mathbf{G} + \lambda \mathbf{I})\mathbf{a} + \mathbf{P}\mathbf{b} - \mathbf{y}\Big) = \mathbf{0} \\ &\frac{\partial J(\mathbf{a},\mathbf{b}|\mathbf{y},\lambda)}{\partial \mathbf{b}} = 2\mathbf{P}^T(\mathbf{G}\mathbf{a} + \mathbf{P}\mathbf{b} - \mathbf{y}) = 0 \\ &\Rightarrow \quad \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} = \begin{pmatrix} (\mathbf{G} + \lambda \mathbf{I}) & \mathbf{P} \\ \mathbf{P}^T & \mathbf{0} \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix} \end{split}$$

Extreme case: $\lambda \to \infty$

$$\mathbf{a}_{(\infty)} = \mathbf{0} \quad \text{ and } \quad \mathbf{b}_{(\infty)} = (\mathbf{P}^T\mathbf{P})^{-1}\mathbf{P}^T\mathbf{y} \qquad = \arg\min_{\mathbf{b}} \|\mathbf{P}\mathbf{b} - \mathbf{y}\|_2^2$$

71

CONTENT

- 1. Preliminaries
- 2. Reproducing kernel Hilbert spaces
- 3. Variational splines and representer theorems
- 4. Gaussian processes
 - Generalized stochastic processes (GSP)
 - Mean and covariance forms
 - The characteristic functional
 - Characterization of Gaussian processes
 - MMSE solution of inverse problems