


# Biogeochemistry is the scientific discipline that studies the physical, chemical and biological processes that govern the composition and health of the natural environment

#### **William Schlesinger**

"the chemistry of the arena of life— that is Earth's biogeochemistry— will be at the center of how well we do, and all biogeochemists should strive to articulate that message clearly and forcefully to the public and to leaders of society, who must know our message to do their job well."

"... the premier scientific discipline to examine human impacts on the global environment."



**Vladimir Vernadsky** 



Biogeochemistry and ecosystem science are inherently linked by the study of the flow of energy and nutrients.

They strive to understand basic mechanistic processes at scales from within ecosystems to the entire globe and provide a framework for investigating aspects of global change





#### **Rivers**

- Natural channels configured to ensure transportation of water and sediments
- Flood events (current: warm Mediterranean, impermeabilisation, channelisation)

# Perception of hydrologists long shaped our understanding of rivers as 'pipelines'

[Physical transport overwhelms biogeochemical transformations]







# The superlatives of rivers

Largest biogeochemical nexus between land, the oceans and atmosphere


**Span multiple catchments and biomes** 

**High biodiversity** 

Provide critical ecosystem services (e.g., drinking water, proteins, hydropower)



## Largest biogeochemical nexus between land, the oceans and atmosphere



#### River ecosystem metabolism

Respiration and Gross Primary Production

- Sum of metabolic activities of all organisms
- Controls carbon and nutrient cycling
- Regulates greenhouse (CO<sub>2</sub>, CH<sub>4</sub>) production


Metabolism = ecosystem process

Carbon and nutrient cycling = biogeochemical process





## Largest biogeochemical nexus between land, the oceans and atmosphere



1 Gigatonne (Gt) = 1 billion tonnes =  $1 \times 10^{15}$ g = 1 Petagram (Pg)

- Rivers figure among the most heterotrophic ecosystems (that is, they respire more than they produce)
- The world's rivers are major CO<sub>2</sub> sources to the atmosphere [...this is a natural process]
- They receive OC from the terrestrial promary production, which they respire, store or transport downstream
- They receive CO<sub>2</sub> from terrestrial respiration and rock weathering, which they outgas to the atmosphere
- These processes shift the ratio of organic-toinorganic carbon en route from small streams to the coastal ocean




River ecosystem metabolism and carbon biogeochemistry in a changing world



# The world's rivers are now part of the IPCC Global Carbon Budget

Chapter 5

**Global Carbon and Other Biogeochemical Cycles and Feedbacks** 



Plumbing the Global Carbon Cycle:
Integrating Inland Waters into the
Terrestrial Carbon Budget

J. J. Cole, Y. T. Prairie, 2.\* N. F. Caraco, W. H. McDowell, 3 L. J. Tranvik, 4
R. G. Striegl, 5 C. M. Duarte, 6 P. Kortelainen, 7 J. A. Downing, 8
J. J. Middelburg, 9 and J. Melack, 10

ECO2 evasion

b

0.75

Land
0.9 Inland waters 
0.9 Ocean

CO2 evasion

b

1.19

Inland waters 
0.9 Ocean

CO3 evasion

b

0.75

Land
1.9

Inland waters 
0.9 Ocean

CO3 evasion

b

0.75

Land
1.9

Inland waters 
0.9 Ocean

CO3 evasion

b

0.75

Land
1.9

Inland waters 
0.9 Ocean

CO3 evasion

b

0.75

Land
1.9


Inland waters 
0.9 Ocean

CO3 evasion


Figure 5.12 | Global carbon (CO<sub>2</sub>) budget (2010–2019). Yellow arrows represent annual carbon fluxes (in PqC yr<sup>-1</sup>) associated with the natural carbon cycle, estimated for the time prior to the industrial era, around 1750. Pink arrows represent anthropogenic fluxes averaged over the period 2010-2019. The rate of carbon accumulation in the atmosphere is equal to net land-use change emissions, including land management (called LULUCF in the main text) plus fossil fuel emissions, minus land and ocean net sinks (plus a small budget imbalance, Table 5.1). Circles with yellow numbers represent pre-industrial carbon stocks in PgC. Circles with pink numbers represent anthropogenic changes to these stocks (cumulative anthropogenic fluxes) since 1750. Anthropogenic net fluxes are reproduced from Friedlingstein et al. (2020). The relative change of gross photosynthesis since pre-industrial times is based on 15 DGVMs used in Friedlingstein et al. (2020). The corresponding emissions by total respiration and fire are those required to match the net land flux, exclusive of net land-use change emissions which are accounted for separately. The cumulative change of anthropogenic carbon in the terrestrial reservoir is the sum of carbon cumulatively lost by net land-use change emissions, and net carbon accumulated since 1750 in response to environmental drivers (warming, rising CO<sub>2</sub>, nitrogen deposition). The adjusted gross natural ocean-atmosphere CO<sub>2</sub> flux was derived by rescaling the value in Figure 1 of Sarmiento and Gruber (2002) of 70 PqC yr<sup>-1</sup> by the revised estimate of the bomb radiocarbon (14C) inventory in the ocean. The original bomb 14C inventory yielded an average global gas transfer velocity of 22 cm hr<sup>-1</sup>; the revised estimate is 17cm hr<sup>-1</sup> leading to 17/22\*70=54. Dissolved organic carbon reservoir and fluxes from Hansell et al. (2009). Dissolved inorganic carbon exchanges between surface and deep ocean, subduction and obduction from Levy et al. (2013). Export production and flux from (Boyd et al., 2019). Net primary production (NPP) and remineralization in surface layer of the ocean from Kwiatkowski et al. (2020); Séférian et al. (2020). Deep ocean reservoir from Keppler et al. (2020). Anthropogenic carbon reservoir in the ocean is from Gruber et al. (2019b) extrapolated to 2015. Fossil fuel reserves are from BGR (2020); fossil fuel resources are 11,490 PgC for coal, 6,780 PgC for oil and 365 PgC for natural gas. Permafrost region stores are from Hugelius et al. (2014); Strauss et al. (2017); Mishra et al. (2021) (see also Box 5.1) and soil carbon stocks outside of permafrost region from Batjes (2016); Jackson et al. (2017). Biomass stocks (range of seven estimates) are from Erb et al. (2018). Sources for the fluxes of the land—ocean continuum are provided in main text and adjusted within the ranges of the various assessment to balance the budget (Section 5.2.1.5).

Canadell, J.G., P.M.S. Monteiro, M.H. Costa, L. Cotrim da Cunha, P.M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C. Koven, A. Lohila, P.K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld, 2021: Global Carbon and other Biogeochemical Cycles and Feedbacks. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, doi: 10.1017/9781009157896.007.





## Global change, river ecosystem processes and biogeochemistry




- Global-change impacts on river ecosystem processes and related biogeochemistry complex and poorly understood.
- Intricate connection with the terrestrial environment and coupling of physical, chemical and biological processes.

Review

River ecosystem metabolism and carbon biogeochemistry in a changing world





Mountains account for 25% of Earth's land surface and their streams contribute more than 30% to global runoff.

Mountains are highly vulnerable to climate change.



#### Mountains are highly vulnerable to climate change






Climate warming and glacier melt (a) Alaska (b) Iceland (c) Scandinavia (d) North Asia A = 86,725 km<sup>2</sup> V = 43.3±11.2 mm SLE A = 11,060 km<sup>2</sup> V = 9.1±2.4 mm SLE A = 2,949 km<sup>2</sup> V = 0.7±0.2 mm SLE  $A = 2,410 \text{ km}^2$ V = 0.3±0.1 mm SLE A= Glacier area e) Western Canada and USA (f) High Mountain Asia V= Volume; SLE= Sea Level Equivalent -2000  $A = 14,524 \text{ km}^2$ V = 2.6±0.7 mm SLE A = 97,605 km<sup>2</sup> V = 16.9±2.7 mm SLE (g) Low Latitudes (h) New Zealand Mass budget 2006-2015 Zemp et al., 2019 Wouters et al., 2019 Gardner et al., 2013 Other regional estimates A = 2,341 km<sup>2</sup> V = 0.2±0.1 mm SLE A = 1,162 km<sup>2</sup> V = 0.2±0.0 mm SLE (Appendix 2.A) (i) Southern Andes (k) Caucasus and Middle East All 11 mountain regions (i) Central Europe 1 mm SLE yr A = 29,429 km<sup>2</sup> V = 12.8±3.3 mm SLE  $A = 2,092 \text{ km}^2$ V = 0.3±0.1 mm SLE A = 1,307 km<sup>2</sup> V = 0.2±0.0 mm SLE A = 251,604 km<sup>2</sup> V = 87±15 mm SLE 1960 1960

### **Consequences of melting mountain glaciers**

Vanishing water towers







Millions of people rely on water from the world's glaciers






### **Cruising the roof of our planet**





#### **Cruising the roof of our planet**

171 glacier-fed streams worldwide





#### **Glacier-fed streams**

Interface between the cryosphere and hydrosphere

- o Freezing cold
- b Long darkness in winter
- o High UV-radiation
- o High turbulence
- o Unstable sediments
- o Ultra-oligotrophic (like the deep sea)





## An amazing 'field team'

Mike Styllas, Martina Schön, Matteo Tolosano, Vincent de Staercke






### **Cruising the roof of our planet**





# 1347 km 87 km










## A team with outstanding qualities















## **Expedition logistics**

Keeping the cryopreservation chain unbroken — from the field to Lausanne











## **Expedition logistics**

Food, hiring porters and guides













Thanks to the more than 150 porters, guides and helpers in Nepal, Uganda, Kyrgyzstan, New Zealand, Norway, Greenland, Chile, Ecuador, Russian Caucasus, and Alaska









#### **Microbial biofilms**

Dominate life in glacier-fed streams












#### **Microbial biofilms**


Most successful form of life over 3.2 billion years

















#### Why microbial biofilms?

Microorganisms are the 'conductors' of Earth's biogeochemical cycles



Understanding the link between microbes, ecosystem processes and biogeochemistry






#### The scientific approach

- Meltwater runoff and downstream water flow descrease
- Less sediments produced and transported downstream
- Streamwater temperature increases

Glacier size, water flow, temperature and suspended sediments are master variables of the glacier-fed stream environment

Space-for-time substitutions




Zhang et al. 2023 Nature Reviews Earth and Environment



#### Scandinavian Mountains Greenland GFS38-GFS41 Central Asia Kyrgystan July-August 2021 GFS120-GFS125 GFS126-GFS129 GFS130-GFS139 Caucaus Russia June-July 2020 September-October 2019 GFS22-GFS29 Andes Ecuador GFS42-GFS46 GFS30-GFS31, GFS76 February-March 2020 Himalaya Nepa GFS47-GFS51 GFS77-GFS81 GFS61-GFS64 GFS52-GFS55 GFS82-GFS85 GFS65-GFS68 GFS56-GFS60 GFS96-GFS102 GFS69-GFS72 GFS108-GFS109 GFS110-GFS114 GFS73-GFS75 Mt. Rwenzori Uganda Andes Chile January-March 2022 Alps New Zealand -February 2019 GFS141-GFS14 GFS143-GFS147 GFS148-GFS150 GFS151-GFS155


#### **Nested sampling design**




- o Microbial biomass
- o 16S &18S rRNA, metagenomics
- o Extracellular enzymes, bacterial carbon production, etc...
- o Glaciology, geology, mineral composition
- o GFS water physico-chemistry



# Resource limitation and microbial starvation








- Microbial metabolism limitated by carbon and phosphorus
- Low microbial carbon use efficiency (median: 0.15)
- Starvation and little growth



A space-for-time substitution approach









#### Global trends indicate

- Streamwater turbidity decreases as glacier
- More light reaches the stream bottom
- Algal biomass is stimulated and streams become greener



Modelling climate-change scenarios





Average for the 2070–2100 period Sampling year 2019–2022

Median chlorophyll-a increase 340%


#### Global trends indicate

- Streamwater turbidity decreases as glaciers shrink
- More light reaches the stream bottom
- Algal biomass is stimulated and streams become greener
- Greening promotes bacterial growth







- Algae provide organic carbon as a resource to the microbial metabolism
- Relieves microbes from energy limitation and starvation
- Increases carbon use efficiencies relatively more organic carbon allocated to biomass growth




Genomic evidence for shifting energetics







Will microbial energetics transit from chemolithotrophy to photoautotrophy, and further to heterotrophy?







Genomic evidence for shifting energetics







Will microbial energetics transit from chemolithotrophy to photoautotrophy, and further to heterotrophy?






#### The 'green transition'

Functional consequences


- Glacier shrinkage stimulates the organic matter decomposition rate
- Modulated by chlorophyll a and temperature
- Decomposition rate partially drives the pace of carbon cycling
- Entails a shift of the glacier-fed stream microbiome





The 'green transition' is only part

of the story



- Glacierized catchments are mineral landscapes
- Chemical weathering (e.g., via carbonic acid, sufuric acid) of minerals
- Drawdown and consumption of atmospheric CO<sub>2</sub>
- Glacier-fed streams as sources of CO<sub>2</sub>
- Likely in equilibrium with the atmosphere before the industrialisation




# Glacier-fed streams are transitional ecosystems

- At the interface to the cryosphere, glacierfed streams are highly vulnerable to climate change
- As glaciers melt, the environment of glacier-fed streams changes dramatically
- Impacts on the microbiome (see lecture by Hannes Peter) – losing biodiversity
- Changing microbial energetics and ecosystem functioning





#### From the 'world' to Valais





**EPFL ALPOLE** 



#### Taking the 'heartbeat' of glacier-fed streams


https://metalp.epfl.ch/







A unique sensor network for monitoring impacts of glacier shrinkage on stream biogeochemistry








#### Taking the 'heartbeat' of glacier-fed streams

https://metalp.epfl.ch/


#### **Our Study Sites**

Explore our stations and their catchments



#### Ferret catchment

	DRANSE DE FERRET DOWNSTREAM	DRANSE DE FERRET UPSTREAM	LA PEULE
Station ID	FED	FEU	PEU
Latitude [°]	45.905091	45.883084	45.893655
Longitude [°]	7.115597	7.130949	7.107973
Station altitude [m]	1773	1996	2024
Catchment avg alt. [m]	2421	2528	2384
Area [km²]	20.24	9.33	3.97
Vegetated area [%]	50.07	39.60	61.19
Glacier coverage [%]	1.78	3.86	0







- Streams and rivers matter for global biogeochemical cycles despite their small contributions by areal extent
- High-mountain streams are highly vulnerable to climate change — yet poorly studied
- Glacier-fed streams are emblematic of climate change
- Transitional ecosystems undergoing major environmental and biogeochemical shifts



https://www.glacierstreams.ch/ https://www.epfl.ch/labs/river/

