



Ens: Z. Patakfalvi
 Analysis I - (n/a)
 November 2019
 65 minutes

n/a

n/a

SCIPER: 999999

Do not turn the page before the start of the exam. This document is double-sided, has 4 pages, the last ones possibly blank. Do not unstaple.

- Place your student card on your table.
- **No other paper materials** are allowed to be used during the exam.
- Using a **calculator** or any electronic device is not permitted during the exam.
- For the **multiple choice** questions, we give :
 - +3 points if your answer is correct,
 - 0 points if you give no answer or more than one,
 - −1 points if your answer is incorrect.
- For the **true/false** questions, we give :
 - +1 points if your answer is correct,
 - 0 points if you give no answer or more than one,
 - −1 points if your answer is incorrect.
- Use a **black or dark blue ballpen** and clearly erase with **correction fluid** if necessary.
- If a question is wrong, the teacher may decide to nullify it.

This exam is printed on recycled paper.

Respectez les consignes suivantes Observe this guidelines Beachten Sie bitte die unten stehenden Richtlinien		
choisir une réponse select an answer Antwort auswählen	ne PAS choisir une réponse NOT select an answer NICHT Antwort auswählen	Corriger une réponse Correct an answer Antwort korrigieren
ce qu'il ne faut PAS faire what should NOT be done was man NICHT tun sollte		

CATALOG

First part: multiple choice questions

For each question, mark the box corresponding to the correct answer. Each question has **exactly one** correct answer.

Question [QCM-inf-sup-E] : Let A be the subset of $\mathbb{R}$ defined by $A = \left\{x > 0 : \cos\left(\frac{1}{x}\right) > 0\right\}$. Then

- ☒ $\inf A = 0$
☐ $\sup A = \frac{\pi}{2}$
☐ $\sup A = 0$
☐ $\inf A = \frac{2}{\pi}$

Question [QCM-complexes-B] : Let $S \subseteq \mathbb{C}$ be the set of solutions of the equation $\bar{z}^2 = z^2$ among the complex numbers. Then:

- ☐ $S = \{-1, +1, -i, +i\}$
☐ $S = \emptyset$
☒ $S = \{z \in \mathbb{C} : \operatorname{Re}(z) = 0 \text{ or } \operatorname{Im}(z) = 0\}$
☐ $S = \mathbb{R}$

Question [QCM-limsup-liminf-B] : Let $(x_n)_{n \geq 1}$ be the sequence defined by $x_n = \sqrt[n]{7}$ if n is even and by $x_n = \frac{1}{n^7}$ if n is odd. Then:

- ☒ $\limsup_{n \rightarrow \infty} x_n = 1$, and $\liminf_{n \rightarrow \infty} x_n = 0$
☐ $\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n = 0$
☐ $\limsup_{n \rightarrow \infty} x_n = 0$, and $\liminf_{n \rightarrow \infty} x_n = 1$
☐ $\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n = 1$

Question [QCM-serie-B] : Set $\lambda := -\frac{1}{6}$. Find the convergent one out of the following series:

- ☒ $\sum_{n=1}^{\infty} \left(\frac{\lambda+1}{\lambda-1}\right)^n$
☐ $\sum_{n=1}^{\infty} \left(\frac{1}{1-\lambda^2}\right)^n$
☐ $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\lambda^n}$
☐ $\sum_{n=1}^{\infty} \frac{1}{n^{1+\lambda}}$

Question [QCM-serie-parametre-B] : Let s be a real number, and let $(b_n)_{n \geq 1}$ be the sequence defined by $b_n = \frac{1}{n^s}$ if n is even, and $b_n = \frac{1}{n^{2s}}$ if n is odd. Then the series $\sum_{n=1}^{\infty} b_n$ converges if and only if

- ☒ $s > 1$
☐ $s > \frac{1}{2}$
☐ $s > 0$
☐ $s > 2$

Question [QCM-suites-recurrence-B] : Let $(a_n)_{n \geq 0}$ be the sequence defined by $a_0 = \frac{3}{2}$, and $a_{n+1} = \frac{1}{2} + \frac{1}{2}\sqrt{8a_n - 7}$, for every $n \geq 0$. Then:

- ☐ $\lim_{n \rightarrow \infty} a_n = 1$
☐ $\lim_{n \rightarrow \infty} a_n = +\infty$
☒ $\lim_{n \rightarrow \infty} a_n = 2$
☐ the sequence is divergent

CATALOG

Question [QCM-limite-prolongmt-B] : Let $m \in \mathbb{R}$ be a real number, and let $f : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by

$$f(x) = \begin{cases} \frac{\sin^2(x)}{\ln(1+2x^2)} & \text{if } x < 0, \\ m & \text{if } x = 0, \\ \frac{x+1}{x^2+3x+1} & \text{if } x > 0. \end{cases}$$

- ☒ If $m = \frac{1}{2}$, then f is left continuous but not right continuous at $x = 0$.
- ☐ If $m = \frac{1}{3}$, then f is right continuous but not left continuous at $x = 0$.
- ☐ If $m = 1$, then f is continuous at $x = 0$.
- ☐ If $m = \frac{1}{2}$, then f is continuous at $x = 0$.

Second part: true/false questions

For each question, mark the box (without erasing) TRUE if the statement is **always true** and the box FALSE if it is **not always true** (i.e., it is sometimes false).

Question [TF-inf-sup-B] : Let $A \subset \mathbb{R}$ be a bounded set, and let $B := \{x \in \mathbb{R} : x \text{ is an upper bound of } A\}$. Then $\inf B \in B$.

☒ TRUE ☐ FALSE

Question [TF-complexes-C] : Let $z \neq 0$ be a complex number with argument $\frac{\pi}{4}$. Then, the argument of the complex number $\frac{1}{z^2}$ is $-\frac{\pi}{2}$.

☒ TRUE ☐ FALSE

Question [TF-fonction-etc-B] : Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a bijective function. Then f is strictly monotone.

☐ TRUE ☒ FALSE

Question [TF-induction-suites-limites-A] : Let $(x_n)_{n \geq 0}$ be the sequence defined by $x_0 = 2$, and $x_n = x_{n-1} - \frac{1}{n}$, for every $n \geq 1$. Then $(x_n)_{n \geq 0}$ is convergent.

☐ TRUE ☒ FALSE