

MATH-111(en)

Linear Algebra

Fall 2024 Annina Iseli

### MINI SOLUTIONS for Homework 11

### Ex 11.2 (Diagonalization of a matrix)

Diagonalize the following matrix.

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix}$$

If not stated otherwise, this means finding a diagonal matrix D and an invertible matrix P such that  $A = PDP^{-1}$ . In particular, you do not need to compute  $P^{-1}$ .

#### **Solution:**

$$D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}, P = \begin{pmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

## Ex 11.3 (Déjà vu?)

Diagonalize the following matrix.

$$\begin{pmatrix}
-2 & 0 & 0 & 0 \\
0 & -2 & 0 & 0 \\
2 & -4 & 2 & 0 \\
0 & 0 & 0 & 2
\end{pmatrix}$$

**Solution:** 

$$P = \begin{pmatrix} 2 & -2 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, D = \begin{pmatrix} -2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$

#### Ex 11.4 (More diagonalizability examples)

Consider

$$A = \begin{pmatrix} 3 & -1 \\ 1 & 5 \end{pmatrix}, B = \begin{pmatrix} 4 & 0 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 5 \end{pmatrix} C = \begin{pmatrix} -1 & 4 & -2 \\ -3 & 4 & 0 \\ -3 & 1 & 3 \end{pmatrix}, D = \begin{pmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 4 \end{pmatrix} \text{ and } E = \begin{pmatrix} 5 & 1 \\ 0 & 5 \end{pmatrix}.$$

- 1. For each matrix find out the eigenvalues and the corresponding eigenvectors.

  Hint: For C and D the rational root theorem (see Ex. 10.8) helps to find the eigenvalues.
- 2. Find out which ones are diagonalizable.

#### Solution:

- A has unique eigenvalue  $\lambda = 4$  with eigenvector  $\begin{pmatrix} -1\\1 \end{pmatrix}$ .
- B has eigenvalues are  $\lambda_1 = 4$  and  $\lambda_2 = 5$  with corresponding eigenvectors  $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ .
- C has the following eigenvalue/eigenvector pairs

$$\begin{pmatrix} 1\\1\\1 \end{pmatrix}$$
 for  $\lambda = 1$ ,  $\begin{pmatrix} 2/3\\1\\1 \end{pmatrix}$  for  $\lambda = 2$ ,  $\begin{pmatrix} 1/4\\3/4\\1 \end{pmatrix}$  for  $\lambda = 3$ .

• D has the following eigenvalue/eigenvector pairs

$$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$
 for  $\lambda = 8$ .  $\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$  for  $\lambda = 2$ .

• E has one eigenvalue 5 with eigenvector  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ .

# Ex 11.6 (Matrix representation of linear maps)

Let  $T: \mathbb{R}^3 \to \mathbb{R}^3$  be the linear transformation given by

$$T\left(\begin{pmatrix} x \\ y \\ z \end{pmatrix}\right) = \begin{pmatrix} 4z \\ 3x + 5y - 2z \\ x + y + 4z \end{pmatrix}.$$

Consider the ordered basis of  $\mathbb{R}^3$  given by

$$\mathcal{B} = \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\}.$$

Find the matrix M = [T] that represents T in the basis  $\mathcal{B}$ .

**Solution:** 

$$[T] = \begin{pmatrix} 6 & 2 & 1\\ 0 & 6 & 2\\ -2 & -8 & -3 \end{pmatrix}$$

# Ex 11.7 (Another matrix representation)

Let the linear transformation  $T: \mathbb{P}_2 \to \mathbb{R}^3$  be defined by:

$$T(\mathbf{p}) = \begin{pmatrix} p(0) \\ p(0) \\ p(2) \end{pmatrix}$$
 for any polynomial  $\mathbf{p} \in \mathbb{P}_2$ .

- a) Find the matrix A of the linear transformation T in terms of the standard bases of  $\mathbb{P}_2$  and  $\mathbb{R}^3$ .
- b) Using the matrix A, determine the kernel and image of T.

## **Solution:**

a) 
$$A = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 2 & 4 \end{pmatrix}.$$

b) 
$$\operatorname{Ker} T=\{\mathbf{p}\in\mathbb{P}_2:\quad p(t)=\alpha\cdot(-2t+t^2)\quad \text{with}\quad \alpha\in\mathbb{R}\}$$
 and

$$\operatorname{Ran} T = \left\{ \mathbf{x} \in \mathbb{R}^3 : \quad \mathbf{x} = \alpha \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ 0 \\ 2 \end{pmatrix} \quad \text{with} \quad \alpha \in \mathbb{R} \quad \text{and} \quad \beta \in \mathbb{R} \right\}.$$