

Analysis III - 203(d)

Winter Semester 2024

Session 3: September 26, 2024

Exercise 1 Consider the scalar field $f(x_1, x_2) = x_1x_2$.

- Describe its level sets for the values $-2, -1, 0, 1, 2$.
- Draw the gradient at a few points, such as

$$(1, 0), (0, 1), (1, 1), (-1, -1), (\sqrt{2}, \sqrt{2}), (\sqrt{2}, -\sqrt{2}) \quad (1)$$

- Recall the hyperbolic sine and hyperbolic cosine functions $\sinh(t)$ and $\cosh(t)$. Sketch the curve

$$\gamma : (-\infty, \infty) \rightarrow \mathbb{R}^2, \quad t \mapsto (\cosh(t), \sinh(t)) \quad (2)$$

and compute its tangent vector $\dot{\gamma}$.

Solution 1 The tangent vector $\dot{\gamma}$ is given by

$$\dot{\gamma}(t) = \begin{pmatrix} \sinh(t) \\ \cosh(t) \end{pmatrix}.$$

x The figure in total has got the following shape:

Exercise 2 Consider the curves

$$\gamma : [-1, 1] \rightarrow \mathbb{R}^2, \quad t \mapsto (t^3, t^2)$$

$$\delta : [-1, 1] \rightarrow \mathbb{R}^2, \quad t \mapsto (\cos(t), \sin(2t))$$

Draw them. Are they simple, closed, differentiable, or regular? What are their tangents? Show that γ is the graph of a function in the first coordinate.

Solution 2 For the first curve the tangent is given by $\dot{\gamma}(t) = (3t^2, 2t)$. Moreover the curve is simple (it does not intersect itself) and differentiable. However, it is not regular since $\dot{\gamma} = \vec{0}$ at $t = 0$.

For the second curve we have the tangent is given by $\dot{\delta}(t) = (-\sin(t), 2\cos(2t))$. Moreover the curve is simple (it does not intersect itself), differentiable and regular (since $\dot{\delta} \neq \vec{0}$).




Figure 1: The dotted arrows show the gradients at the points $(0, 1)$ (a), $(1, 0)$ (b), $(1, 1)$ (c), and $(-1, -1)$ (d). The solid black line is the curve γ .

For the first curve γ , we can define another coordinate $s(t) := t^3$, such that $s([-1, 1]) = [-1, 1]$. Then, the curve can be expressed in s as $\gamma(s) = (s, \sqrt[3]{s^2})$, and thus is a graph in the first coordinate. Note here, that we cannot apply the same method to the second variable. Indeed, defining $\tilde{s}(t) := t^2$ would not work, as $\tilde{s}([-1, 1]) = [0, 1]$. Therefore, the corresponding change of variables would not be bijective.

Exercise 3 Compute the line integral of

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x_1, x_2) \mapsto 1 + x_1^2 + x_2^2$$

along the two curves γ and δ , given by

$$\gamma : [0, \pi] \rightarrow \mathbb{R}^2, \quad t \mapsto (-\cos(t), \sin(t))$$

$$\delta : [-1, 1] \rightarrow \mathbb{R}^2, \quad t \mapsto (t, 0)$$

Compare the results. What are the endpoints of the two curves?

Solution 3

$$\int_{\Gamma} f(x_1, x_2) d\ell = \int_0^{\pi} (1 + \cos(t)^2 + \sin(t)^2) |\dot{\gamma}(t)| dt$$

$$\begin{aligned}
&= \int_0^\pi (1 + \cos(t)^2 + \sin(t)^2) \sqrt{\cos(t)^2 + \sin(t)^2} dt \\
&= \int_0^\pi (1 + 1) \sqrt{1} dt \\
&= \int_0^\pi 2 dt \\
&= 2\pi.
\end{aligned}$$

The endpoints of the first curve are: $\gamma(0) = (-1, 0)$ and $\gamma(\pi) = (1, 0)$.
Let us now compute the second integral. We see

$$\begin{aligned}
\int_\Gamma f(x_1, x_2) dl &= \int_{-1}^1 (f \circ \delta)(t) |\dot{\delta}(t)| dt \\
&= \int_{-1}^1 (1 + t^2) \left| \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right| dt \\
&= \int_{-1}^1 (1 + t^2) dt \\
&= 2 \int_0^1 (1 + t^2) dt \\
&= 2 \left[t + \frac{t^3}{3} \right]_0^1 \\
&= \frac{8}{3}.
\end{aligned}$$

Note that from the third to fourth line, we have used that the integrand was even. The endpoints of the second curve are: $\delta(-1) = (-1, 0)$ and $\delta(1) = (1, 0)$.

The exercise also shows that integrating a function f along different curves may give different results, even if the curves share the same starting and endpoints.

Exercise 4 Compute the line integral of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ along the circle C with radius 3 centered at the origin, where

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x_1, x_2) \mapsto 3x_2^2 + x_2^3$$

Hint: you must first find a parameterization of C .

Solution 4 We may choose the following parameterization for the circle C :

$$\gamma : [-\pi, \pi] \rightarrow \mathbb{R}^2, \quad t \mapsto (3 \cos(t), 3 \sin(t))$$

$$\int_\Gamma f(x_1, x_2) dl = \int_{-\pi}^\pi (f \circ \gamma)(t) |\dot{\gamma}(t)| dt$$

$$\begin{aligned}
&= \int_{-\pi}^{\pi} [3(3 \sin(t))^2 + (3 \sin(t))^3] \left\| \begin{pmatrix} -3 \sin(t) \\ 3 \cos(t) \end{pmatrix} \right\| dt \\
&= \int_{-\pi}^{\pi} 3 [27 \sin^2(t) + 27 \sin^3(t)] dt \\
&= 81 \int_{-\pi}^{\pi} \sin^2(t) dt + 81 \int_{-\pi}^{\pi} \sin^3(t) dt
\end{aligned}$$

The second integral is 0 since the integrand is uneven. For the first integral we use the cosine half-angle formula ($1 - 2 \sin^2(\alpha) = \cos(2\alpha)$).

$$81 \int_{-\pi}^{\pi} \sin^2(t) dt = 81 \int_{-\pi}^{\pi} \frac{1}{2} - \frac{1}{2} \cos(2t) dt = 81 \left[\frac{1}{2}t - \frac{1}{4} \sin(2t) \right]_{-\pi}^{\pi} = 81\pi$$

Exercise 5 You are excavating a tunnel deep beneath surface through rock from a point $A = (-1, 0)$ to a point $B = (1, 1)$ along a curve Γ . Geological observations show that the rock mass density of the region can be modeled by

$$f(x_1, x_2) = \exp(x_1 + x_2).$$

The costs and abbrasion of the tools are therefore proportional to the curve integral

$$\int_{\Gamma} f \, dl$$

Compute the integral along the straight line

$$\gamma : [0, 1] \rightarrow \mathbb{R}^2, \quad t \mapsto (-1 + 2t, t)$$

Solution 5 We set up the first curve integral and compute:

$$\begin{aligned}
\int_{\Gamma} f \cdot dl &= \int_0^1 \exp(-1 + 2t + t) \|(2, 1)\| dt \\
&= \int_0^1 \exp(-1 + 2t + t) \sqrt{5} dt \\
&= \sqrt{5} \int_0^1 \exp(3t - 1) dt \\
&= \frac{\sqrt{5}}{3} \int_0^1 3 \exp(3t - 1) dt \\
&= \frac{\sqrt{5}}{3} \left[\exp(3t - 1) \right]_0^1 = \frac{\sqrt{5}}{3} (\exp(2) - \exp(-1)) = \frac{\sqrt{5}}{3} \left(e^2 - \frac{1}{e} \right).
\end{aligned}$$

Exercise 6 A matrix $A \in \mathbb{R}^{n \times n}$ is called symmetric if $A = A^t$ and antisymmetric if $A = -A^t$.

1. Show that a matrix that is both symmetric and antisymmetric must be zero.

2. Show that every matrix $B \in \mathbb{R}^{n \times n}$ can be written as a sum $A = A_1 + A_2$ where $A_1 \in \mathbb{R}^{n \times n}$ is symmetric and $A_2 \in \mathbb{R}^{n \times n}$ is antisymmetric.
3. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a scalar field with continuous partial derivatives up to second order. Show that its Hessian is a symmetric matrix.
4. The trace of a matrix is the sum of its diagonal elements. Find a scalar field in \mathbb{R}^2 whose Hessian is not zero but has zero trace.
5. Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a differentiable vector field. What is the antisymmetric part of its Jacobian?
6. Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a differentiable vector field. What is the antisymmetric part of its Jacobian?
7. What do you notice in the answers to the last two questions?

Solution 6 1. Assume $A = (a_{ij})_{ij} \in \mathbb{R}^{n \times n}$ is both symmetric and antisymmetric. From the symmetry of A we deduce for an entry a_{ij} that $a_{ij} = a_{ji}$. From the antisymmetry of A we deduce that $a_{ij} = -a_{ji}$. Therefore $a_{ij} = -a_{ij}$ for all i, j and hence $a_{ij} = 0$ for all i, j .

2. We can set $A_1 := \frac{1}{2}(B + B^t)$ and $A_2 := \frac{1}{2}(B - B^t)$.
3. By Schwartz' theorem, it holds that $\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$ for all i, j . Hence, the Hessian is symmetric.
4. We can use $f(x_1, x_2) = x_1 x_2$.
5. Writing $F = [F_x, F_y]^T$, we find that the antisymmetric part of the Jacobian is given by

$$\frac{1}{2} \begin{pmatrix} 0 & \frac{\partial F_x}{\partial y} - \frac{\partial F_y}{\partial x} \\ \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} & 0 \end{pmatrix}$$

6. Similarly to before, we find by writing $G = [G_x, G_y, G_z]^T$ that the antisymmetric part of the Jacobian is given by

$$\frac{1}{2} \begin{pmatrix} 0 & \frac{\partial G_x}{\partial y} - \frac{\partial G_y}{\partial x} & \frac{\partial G_x}{\partial z} - \frac{\partial G_z}{\partial x} \\ \frac{\partial G_y}{\partial x} - \frac{\partial G_x}{\partial y} & 0 & \frac{\partial G_y}{\partial z} - \frac{\partial G_z}{\partial y} \\ \frac{\partial G_z}{\partial x} - \frac{\partial G_x}{\partial z} & \frac{\partial G_z}{\partial y} - \frac{\partial G_y}{\partial z} & 0 \end{pmatrix}.$$

7. We notice that the entries in the antisymmetric parts of the Jacobians contain already all the information for the curl of the respective vector fields (in 2D and 3D). More generally, the antisymmetric part of the Jacobian of a vector field $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is sometimes identified as the "curl" of that vector field in any dimension.