

Graph Theory - Problem Set 9

November 14, 2024

Exercises

1. Deduce the undirected version of Menger's theorem from the directed version.
2. Let G be a k -connected graph. Show using the definitions that if G' is obtained from G by adding a new vertex V adjacent to at least k vertices of G , then G' is k -connected.
3. Prove that a graph G on at least $k+1$ vertices is k -connected if and only if $G-X$ is connected for every vertex set X of size $k-1$.

Problems

4. Prove the following variants of Menger's theorem. Let G be a graph and let S, T be disjoint vertex sets. An S - T path is a path with one endpoint in S and the other in T . Then:
 - (a) The maximum number of edge-disjoint S - T paths equals the min size of an S - T edge separator.
 - (b) If $|S|, |T| \geq k$ and there is no S - T separator of size $k-1$, then G contains k internally vertex disjoint S - T paths.
(An S - T separator $X \subseteq V(G)$ is a set such that $G-X$ has no path between $S \setminus X$ and $T \setminus X$.)
5. Find a graph G with $\kappa(G) = 10$ and $\kappa'(G) \geq 50$.
6. Let G be a connected graph with all degrees even. Show that G is 2-edge-connected.
7. Show that if G is a graph with $|V(G)| = n \geq k+1$ and $\delta(G) \geq (n+k-2)/2$ then G is k -connected.
8. Prove that G is 2-connected if and only if for any three vertices x, y, z there is a path in G from x to z containing y .