

Series 6 - October 30, 2024

Exercise 1.

(The maximum principle) Consider a d -dimensional SDE driven by an m -dimensional Brownian motion with $m \geq d$.

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t, \quad t \geq 0$$

and its corresponding generator

$$L = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(x) \frac{\partial}{\partial x_i}$$

with $a_{ij} = \sum_{l=1}^m \sigma_{il} \sigma_{jl}$. Let $D \subset \mathbb{R}^d$ be a bounded domain and suppose b, σ globally Lipschitz and satisfying the linear growth bound. Moreover assume a uniformly elliptic on $\bar{D}$ (i.e. $\exists \lambda > 0$ such that $v^T a(x) v \geq \lambda |v|^2$ for all $x \in \bar{D}$, $v \in \mathbb{R}^d$). Let $u \in C^2(D) \cap C(\bar{D})$. Using the Feynman-Kac formula, show the maximum principle, i.e.

- if $Lu \geq 0$, then $u(x) \leq \max_{\partial D} u$ for all $x \in D$
- if $Lu = 0$, then $\min_{\partial D} u \leq u(x) \leq \max_{\partial D} u$ for all $x \in D$

Exercise 2.

Consider a d -dimensional SDE driven by an m -dimensional Brownian motion with $m \geq d$.

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t, \quad t \geq 0 \tag{2.1}$$

and its corresponding generator

$$L = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(x) \frac{\partial}{\partial x_i}$$

with $a_{ij} = \sum_{l=1}^m \sigma_{il} \sigma_{jl}$. Let $D \subset \mathbb{R}^d$ be a bounded open domain and suppose b, σ globally Lipschitz and satisfying the linear growth bound. Moreover assume a uniformly elliptic on $\bar{D}$ (i.e. $\exists \lambda > 0$ such that $v^T a(x) v \geq \lambda |v|^2$ for all $x \in \bar{D}$, $v \in \mathbb{R}^d$).

- 1) Show that if $u \in C^2(D) \cap C(\bar{D})$ is the solution of

$$\begin{cases} Lu = -1 & \text{on } D \\ u|_{\partial D} = 0, \end{cases} \tag{2.2}$$

then u has the characterization $u(x) = \mathbb{E}^x(\tau)$ where $\tau = \inf\{t : X_t \notin D\}$ and $\mathbb{E}^x$ denotes expectation when the process (2.1) starts in x at time zero.

- 2) Let $\{B_t\}_t$ be a one dimensional Brownian motion and $\tau = \inf\{t : B_t \notin (-1, 1)\}$. Compute $\mathbb{E}[\tau]$.

3) Let

$$\begin{aligned} dX_t &= \sigma(X_t)dB_t \\ X_0 &= x \end{aligned} \tag{2.3}$$

with σ globally Lipschitz on $\mathbb{R}$ and strictly positive in $[0, 1]$. Let $\tau = \inf\{t : X_t \notin (0, 1)\}$ be the exit time from $(0, 1)$. Show that $\tau < \infty$ a.s. and compute $P(X_\tau = 1)$ (notice that this probability does not depend on σ).

Exercise 3.

Let $\lambda = 2$, $\mu = 1$ and consider the stochastic differential equation

$$\begin{aligned} dX(t) &= \lambda X(t)dt + \mu X(t)dW(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0, \end{aligned} \tag{3.1}$$

and the Euler–Maruyama (EM) method for (3.1)

$$X_{n+1} = X_n + \lambda X_n \Delta t + \mu X_n (W(t_{n+1}) - W(t_n)).$$

The exact solution of (3.1) is given by $X(t) = X_0 \exp((\lambda - \frac{1}{2}\mu^2)t + \mu W(t))$.

Compute a discretized Brownian path over $[0, 1]$ with $\delta t = 2^{-8}$ and compare the exact solution (on the discretized path) with the EM method (using the same Brownian path) with $\Delta t = 2^4 \delta t, 2^2 \delta t$.

Exercise 4.

A numerical method for an SDE

$$\begin{aligned} dX(t) &= f(X(t))dt + g(X(t))dW(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0, \end{aligned} \tag{4.1}$$

is said to have a *strong order of convergence equals to r* if there exists a constant C such that

$$\mathbb{E} \left[\sup_{0 \leq n \leq N} |X_n - X(t_n)| \right] \leq C(\Delta t)^r,$$

with $N = T/\Delta t$. Consider the following one dimensional SDE

$$\begin{aligned} dX(t) &= \lambda X(t)dt + \mu X(t)dW(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0, \end{aligned} \tag{4.2}$$

and set $t_n = T = 1$. Consider $\{X_n\}_{n=0}^N$ the Euler-Maruyama approximation and set $e_{\Delta t}^s := \mathbb{E}|X_N - X(T)|$. Verify numerically that the Euler-Maruyama method satisfies $e_{\Delta t}^s \leq C(\Delta t)^{1/2}$. To evaluate $\mathbb{E}|X_N - X(T)|$ you need to compute $\frac{1}{M} \sum_{i=1}^M |X_N^i - X^i(T)|$, i.e., the average over M realizations of the random variables at time $T = 1$. For that:

- i) take $M = 10^5$ independent discretized Brownian path over $[0, 1]$ with $\delta t = 2^{-10}$,
- ii) for each path apply EM with $\Delta t = 2^p \delta t$, $1 \leq p \leq 5$ and store the endpoint error (at $t = T$),
- iii) take the mean over the error and then report the result (Δt versus $e_{\Delta t}^s$) in a loglog plot.

Exercise 5.

A numerical method for an SDE

$$\begin{aligned} dX(t) &= f(X(t))dt + g(X(t))dW(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0, \end{aligned} \tag{5.1}$$

is said to have a *weak order of convergence equals to r* if there exists a constant C such that

$$\sup_{0 \leq n \leq N} |\mathbb{E}p(X_n) - \mathbb{E}p(X(t_n))| \leq C(\Delta t)^r,$$

with $N = T/\Delta t$ and all sufficiently smooth function p . Consider the following one dimensional SDE

$$\begin{aligned} dX(t) &= \lambda X(t)dt + \mu X(t)dW(t), \quad 0 \leq t \leq T, \\ X(0) &= X_0, \end{aligned} \tag{5.2}$$

with $\lambda = 2$, $\mu = 1$, and $e_{\Delta t}^w := |\mathbb{E}(X_N) - \mathbb{E}(X(T))|$ and verify numerically that the Euler-Maruyama method satisfies $e_{\Delta t}^w \leq C\Delta t$.