

## Exercises – week 12

**Exercise 1.** *Homotopy invariance of class groups.* Let  $X$  be integral, Noetherian, separated and regular in codimension 1.

- (1) Show that  $X \times \mathbb{A}^1$  is also integral, Noetherian, separated and regular in codimension 1.
- (2) Show that the projection  $\pi$  to the first component induces a morphism  $\pi^*: \text{Cl}(X) \rightarrow \text{Cl}(X \times \mathbb{A}^1)$ .
- (3) Show that  $\pi^*$  is an isomorphism.

*You may look at II.6.6 in Hartshorne.*

**Exercise 2.** *A Künneth formula for class groups.* Let  $X$  be an integral, separated, Noetherian and locally factorial scheme. Let  $n \geq 1$ . Show that  $\mathbb{P}_X^n = X \times \mathbb{P}_{\mathbb{Z}}^n$  also satisfies the above and that

$$\text{Cl}(X \times \mathbb{P}_{\mathbb{Z}}^n) \cong \text{Cl}(X) \times \mathbb{Z}.$$

*Hint: Consider  $\phi: \mathbb{P}_{K(X)}^n \rightarrow X \times \mathbb{P}_{\mathbb{Z}}^n$ . Show that  $\phi^*: \text{Pic}(\mathbb{P}_{K(X)}^n) \rightarrow \text{Pic}(\mathbb{P}_{K(X)}^n) \cong \mathbb{Z}$  gives a retraction of the first arrow in the exact sequence (exercise 8, week 9)*

$$\mathbb{Z} \rightarrow \text{Cl}(\mathbb{P}_X^n) \rightarrow \text{Cl}(\mathbb{A}_X^n) \rightarrow 0$$

*coming from the divisor  $V_+(X_0)$  in  $\text{Cl}(\mathbb{P}_X^n)$ .*

**Exercise 3.** *Very ample divisors.* Let  $k$  be a field. Let  $S$  be a  $\mathbb{N}$ -graded ring finitely generated in degree 1 with  $S_0 = k$ . Denote by  $X = \text{Proj}(S)$ . Suppose that  $X$  is integral and  $\mathcal{O}_X(X) = k$ .<sup>1</sup>

- (1) Show that  $\mathcal{O}_X(1)$  is  $k$ -very ample.
- (2) If  $\dim(X) \geq 1$ , show that  $\mathbb{Z} \xrightarrow{\mathcal{O}_X(1)} \text{Pic}(X)$  is injective.  
*Hint: If  $\mathcal{O}_X(1)$  is torsion, it would imply that  $\mathcal{O}_X$  is  $k$ -very ample.*
- (3) If  $X$  is normal, deduce that if  $0 \neq s \in \mathcal{O}_X(1)(X)$ , then  $\text{div}(s) \in \text{Cl}(X)$  has infinite order.

**Exercise 4.** *Projective Cone.* This exercise is a follow-up to exercise 2, week 7.

Let  $S$  be a  $\mathbb{N}$ -graded ring finitely generated in degree 1 over  $S_0$ . Consider the  $\mathbb{N}$ -graded ring  $S[t]$  with elements of  $S$  keeping their grading and with  $t$  placed in degree 1. We call  $\text{Proj}(S[t])$  with this grading the *projective cone*.

---

<sup>1</sup>This condition follows from previous assumptions if  $k$  is algebraically closed.

- (1) Show that this grading comes from the product action

$$\mathbb{G}_{m,S_0} \times_{S_0} \text{Spec}(S) \times_{S_0} \mathbb{A}_{S_0}^1 \xrightarrow{(\mu_S \text{ pr}_{12}, \mu_{\mathbb{A}^1} \text{ pr}_{13})} \text{Spec}(S) \times_{S_0} \mathbb{A}_{S_0}^1$$

where  $\text{pr}_{ij}$  denote projections,  $\mu_S$  the action on  $\text{Spec}(S)$  and  $\mu_{\mathbb{A}^1}$  the usual  $\mathbb{G}_m$ -action on  $\mathbb{A}^1$ .

- (2) Show that there are natural identifications  $V_+(t) = \text{Proj}(S)$  and  $D_+(t) = \text{Spec}(S)$ . Show furthermore that  $V_+(S_+)$  (taken in  $\text{Proj}(S[t])$ ) identifies to the vertex (see exercise 6, week 10) in  $\text{Spec}(S)$ . We therefore denote this closed subscheme by  $v$ .
- (3) Let  $s_0, \dots, s_n$  be generators of  $S$  in degree 1. Show that  $\text{Proj}(S[t]) \setminus v$  is covered by the open sets  $D_+(s_i)$  and that each open set is isomorphic to  $\text{Spec}(S_{(s_i)}[t])$ . Deduce that we have a natural map

$$p: \text{Proj}(S[t]) \setminus v \rightarrow \text{Proj}(S).$$

- (4) Let  $k$  be an algebraically closed field, and suppose  $S_0 = k$ . Suppose that  $S$  is integral, Noetherian and normal. Suppose that  $X = \text{Proj}(S)$  is of dimension  $\geq 1$ . Show that  $p^*$  induces an isomorphism on class groups. Deduce that, if  $C = \text{Spec}(S)$  denotes the cone of  $X$  then we have an exact sequence

$$1 \rightarrow \mathbb{Z} \rightarrow \text{Cl}(X) \rightarrow \text{Cl}(C) \rightarrow 1$$

where the first morphism sends 1 to the class of  $\mathcal{O}_X(1)$ , and the second is the composition of  $p^*$  and the restriction to  $C$ .

**Exercise 5.** *Computations of class groups on quadric hypersurfaces.* Suppose that  $k$  is algebraically closed and  $\text{char}(k) \neq 2$ . Let  $2 \leq r \leq n$ . Consider the ring (equipped with the standard grading)

$$S_r = k[x_0, \dots, x_n] / (x_0^2 + \dots + x_r^2).$$

You can assume that this ring is normal. (See Hartshorne, exercise 6.4 for a proof).

- (1) Show that up to a linear change of variable, we can suppose that

$$S_r = k[x_0, \dots, x_n] / (x_0x_1 + x_2^2 \dots + x_r^2).$$

Denote by  $C_r = \text{Spec}(S_r)$  and  $X_r = \text{Proj}(S_r)$ .

- (2) Show that  $\text{Cl}(C_r)$  is cyclic when  $r \neq 3$   
*Hint: Consider the prime divisor  $V(\sqrt{(x_1)})$  and the exact sequence of week 9, exercise 8.*
- (3) Show that  $\text{Cl}(C_2) \cong \mathbb{Z}/2\mathbb{Z}$ .  
*Hint: Consider the same exact sequence. See Hartshorne example 6.5.2.*
- (4) Show that  $\text{Cl}(C_3) \cong \mathbb{Z}$ .  
*Hint: show that after a suitable change of variable we see that  $X_r \cong \mathbb{P}_k^1 \times \mathbb{P}_k^1$ . Then use exercise 1 and the exact sequence of the last point of the above exercise*
- (5) Show that  $\text{Cl}(C_r) \cong 0$  if  $r \geq 4$ . In particular,  $S_r$  is factorial.  
*Hint: show that  $(x_1)$  is prime in this case and conclude.*

- (6) Use the exact sequence of the last point of the above exercise to compute  $\text{Cl}(X_r)$  for all  $r \geq 2$ .

**Exercise to hand in.** *Morphisms between projective spaces, again.* (Due 16 December, 18:00) Please write your solution in  $\text{T}_E\text{X}$ .

Let  $k$  be a field. You may find part (1), (2) of the exercise useful while solving part (3) of the exercise.

- (1) (**Graded Nakayama**) Let  $R = \bigoplus_{n \in \mathbb{N}} R_n$  be a graded ring and  $M = \bigoplus_{n \in \mathbb{N}} M_n$  be a graded  $R$  module,  $J \subseteq R_+$  be a homogeneous ideal of  $R$ . If  $M = JM$ , then show that  $M = 0$ . Recall that  $R_+$  is the homogeneous ideal  $\bigoplus_{n > 0} R_n$  of  $R$ .
- (2) Let  $R$  be a Noetherian ring generated in degree 1 as an  $R_0$ -algebra.<sup>2</sup> Let  $F_0, \dots, F_m$  be homogeneous elements of strictly positive degrees of  $R$ . Assume that the radical of  $(F_0, \dots, F_r)$  is  $R_+$ . Show that the graded ring inclusion  $R_0[F_0, \dots, F_r] \rightarrow R$  is a finite ring map. You may want to use part (1).
- (3) Given a morphism of  $k$ -schemes  $f: \mathbb{P}_k^n \rightarrow \mathbb{P}_k^m$ , show that the image is either a point; and if the image is not a point, then  $m \geq n$  and the image has dimension  $n$ . In the second case, show that the morphism is finite. *Hint: we have  $f^* \mathcal{O}_{\mathbb{P}_k^m}(1) \cong \mathcal{O}_{\mathbb{P}_k^n}(d)$  for some  $d \geq 0$ . Break the study into two cases:  $d = 0$  and  $d \geq 1$ . In this last case show that the polynomials  $F_0, \dots, F_m$  homogeneous of degree  $d$  which defines the map generates an ideal which radical is  $k[X_0, \dots, X_n]_+$ .*
- (4) Identify  $\mathbb{P}_k^n = \text{Proj}(k[x_0, \dots, x_n])$ . Show that the projection map from  $\mathbb{P}_k^n - [0 : 0 : \dots : 1]$  to  $\mathbb{P}_k^{n-1}$  given by the sections  $x_0, x_1, \dots, x_{n-1} \in \mathcal{O}_{\mathbb{P}_k^n}(1)$  cannot be extended to  $\mathbb{P}_k^n$ .

**Remark.** When the image of a map  $\mathbb{P}_k^n \rightarrow \mathbb{P}_k^m$  is not a point, the map  $\mathbb{P}_k^n \rightarrow \mathbb{P}_k^m$  can be written as a composition of (1) a Veronese embedding  $\mathbb{P}_k^n \rightarrow \mathbb{P}_k^N$ , for some  $N$ , (2) an automorphism of  $\mathbb{P}_k^N$ , (3) a projection map

$$\mathbb{P}_k^N - V(X_0, \dots, X_{m'}) \rightarrow \mathbb{P}_k^{m'},$$

sending  $(x_0 : \dots : x_N)$  to  $(x_0 : \dots : x_{m'})$ , for some  $m' \leq m$ , (4) a linear embedding  $\mathbb{P}_k^{m'} \rightarrow \mathbb{P}_k^m$  – here, a linear embedding is map that sends  $\underline{x} = (x_0 : \dots : x_{m'})$  to  $(x_0 : \dots : x_{m'} : L_{m'+1}(\underline{x}) : \dots : L_m(\underline{x}))$ , where the  $L_j$ 's are some linear polynomials in  $m' + 1$  variable with  $k$ -coefficients – and (5) an automorphism which is a permutation of variables  $\mathbb{P}_k^m$ .

Indeed, a map  $\mathbb{P}_k^n \rightarrow \mathbb{P}_k^m$  is given by homogeneous  $m + 1$  polynomials of some fixed degree  $d$ , that we denote by  $F_0, \dots, F_m$ . By functoriality of  $\text{Proj}$  we can express this map (where the first map is given by  $X_i \mapsto F_i$ )

$$k[X_0, \dots, X_m] \rightarrow (k[X_0, \dots, X_n])_d \subset k[X_0, \dots, X_n].$$

So we can factor by a Veronese embedding

$$k[X_0, \dots, X_m] \rightarrow k[Y_j]_{j=0}^N \rightarrow (k[X_0, \dots, X_n])_d$$

<sup>2</sup>It is enough to assume that  $R$  is  $\mathbb{N}$ -graded Noetherian.

sending  $X_i$  to lifts of  $F_i$ 's that we denote by  $G_i$ . Up to permuting variables of  $\mathbb{P}_k^m$  we can suppose that  $G_0, \dots, G_{m'}$  for some  $0 \leq m' \leq m$  form a basis of  $\text{span}(G_0, \dots, G_m)$  in  $\bigoplus_{j=0}^N kY_j$ . Using an automorphism of  $\mathbb{P}_k^N$  we can suppose that  $G_0, \dots, G_{m'}$  are equal to  $Y_0, \dots, Y_{m'}$  and that  $G_{m'+1}, \dots, G_m$  are  $k$ -linear combination of the former, say  $G_l = L_l(Y_0, \dots, Y_{m'})$  for  $m'+1 \leq l \leq m$ . Say  $k[X_0, \dots, X_m] \xrightarrow{\varphi} k[Y_0, \dots, Y_{m'}]$  is given sending  $X_i \rightarrow Y_i$  if  $i \leq m'$  and  $X_i \rightarrow L_i(Y_0, \dots, Y_{m'})$ . Now, all in all the composition

$$\begin{aligned} k[X_0, \dots, X_m] &\xrightarrow{\text{permutation}} k[X_0, \dots, X_m] \xrightarrow{\varphi} k[Y_0, \dots, Y_{m'}] \\ &\xrightarrow{\subseteq} k[Y_0, \dots, Y_N] \xrightarrow{\text{automorphism}} k[Y_0, \dots, Y_N] \rightarrow (k[X_0, \dots, X_n])_d \end{aligned}$$

induces by functoriality of Proj the map we started with.

- (5) Explicitly decompose the map  $\mathbb{P}_k^1 \rightarrow \mathbb{P}_k^2$  sending  $[x : y]$  to  $[x^2 : x^2 + y^2 + xy : 2x^2 + y^2 + xy]$ , into the steps mentioned in the previous remark.