


**ME-445 AERODYNAMICS**  
**Exercise 06**  
**Week 5**



# Formula sheet

## Cylindrical coordinates

$$\nabla \vec{u} = \left( \frac{\partial v_r}{\partial r}, \frac{1}{r} \frac{\partial v_\theta}{\partial \theta}, 0 \right)$$

$$\nabla \cdot \vec{u} = \frac{1}{r} \frac{\partial(rv_r)}{\partial r} + \frac{1}{r} \frac{\partial v_\theta}{\partial \theta}$$

$$\nabla \times \vec{u} = \left( 0, 0, \frac{1}{r} \left[ \frac{\partial(rv_\theta)}{\partial r} - \frac{\partial v_r}{\partial \theta} \right] \right)$$

## Potential flow

$$v_r = \frac{\partial \phi}{\partial r} = \frac{1}{r} \frac{\partial \psi}{\partial \theta}, \quad v_\theta = \frac{1}{r} \frac{\partial \phi}{\partial \theta} = -\frac{\partial \psi}{\partial r}$$

Uniform parallel flow  $w = \phi + i\psi = U_\infty e^{-i\alpha} z$

Potential vortex in  $z_0$   $w = -\frac{i\gamma}{2\pi} \ln(z - z_0)$

Point source or sink in  $z_0$   $w = \frac{Q}{2\pi} \ln(z - z_0)$

Source-sink doublet in  $z_0$   $w = \frac{\mu}{2\pi(z - z_0)}$

$$\frac{dw}{dz} = u - iv$$

Milne-Thomson circle theorem:

$$g(z) = w(z) + \overline{w\left(\frac{a^2}{\bar{z}}\right)}$$

## Thin airfoil theory

For a camber line with:

$$\frac{dy_c}{dx} = A_0 + \sum_{n=1}^{\infty} A_n \cos n\theta$$

$$\frac{x}{c} = \frac{(1 - \cos \theta)}{2}$$

we know:

$$k = 2U_\infty \left[ (\alpha - A_0) \frac{\cos \theta + 1}{\sin \theta} + \sum_{n=1}^{\infty} A_n \sin n\theta \right]$$

$$A_0 = \frac{1}{\pi} \int_0^\pi \frac{dy_c}{dx} d\theta$$

$$A_n = \frac{2}{\pi} \int_0^\pi \frac{dy_c}{dx} \cos n\theta d\theta$$

$$C_l = 2\pi\alpha + \pi(A_1 - 2A_0)$$

$$C_{m,1/4} = -\frac{\pi}{4}(A_1 - A_2)$$

$$x_{cp} = \frac{1}{4} + \frac{\pi}{4C_l}(A_1 - A_2)$$

## Finite wings with $AR=b^2/S$

**Sign convention:**

if induced velocity points downward:  $w(y) > 0$ ,  $\alpha_i(y) > 0$

if induced velocity points upward:  $w < 0$ ,  $\alpha_i < 0$

Prandtl's lifting-line theory

$$U_\infty \alpha_i(y_0) = w(y_0) = -\frac{1}{4\pi} \int_{-b/2}^{b/2} \frac{(d\Gamma/dy)}{y - y_0} dy$$


$$\alpha(y_0) = \alpha_{eff}(y_0) + \alpha_i(y_0)$$

**Elliptical loading**  $\Gamma(y) = \Gamma_0 \sqrt{1 - \left(\frac{2y}{b}\right)^2}$

$$w = \frac{\Gamma_0}{2b}$$

$$\alpha_i = \frac{C_L}{\pi AR}$$

$$C_{D,i} = \frac{C_L^2}{\pi AR}$$



**General loading**  $\Gamma(\theta) = 2bU_\infty \sum_{n=1}^{\infty} A_n \sin n\theta$

$$w(\theta) = U_\infty \sum_{n=1}^{\infty} n A_n \frac{\sin n\theta}{\sin \theta}$$

$$C_L = \pi A_1 AR$$

$$C_{D,i} = \frac{C_L^2}{\pi AR} (1 + \delta) \text{ with } \delta = \sum_{n=2}^{\infty} n (A_n/A_1)^2$$

## Boundary Layer

Flat plate **laminar** boundary layer

$$\frac{\delta}{x} = \frac{5}{\sqrt{Re_x}} \quad \text{boundary layer growth}$$

$$C_f = \frac{1.328}{\sqrt{Re_x}} \quad \text{skin friction drag coefficient}$$

Flat plate **turbulent** boundary layer

$$\frac{\delta}{x} = \frac{0.37}{Re_x^{1/5}} \quad \text{boundary layer growth}$$

$$C_f = \frac{0.074}{Re_x^{1/5}} \quad \text{skin friction drag coefficient}$$

## Miscellaneous

| $\theta$      | $0^\circ$ | $30^\circ$           | $45^\circ$           | $60^\circ$           | $90^\circ$ |
|---------------|-----------|----------------------|----------------------|----------------------|------------|
| $\sin \theta$ | 0         | $\frac{1}{2}$        | $\frac{\sqrt{2}}{2}$ | $\frac{\sqrt{3}}{2}$ | 1          |
| $\cos \theta$ | 1         | $\frac{\sqrt{3}}{2}$ | $\frac{\sqrt{2}}{2}$ | $\frac{1}{2}$        | 0          |

**water**

kinematic viscosity  $\nu = 1 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$

density  $\rho = 1000 \text{ kg m}^{-3}$

**air**

kinematic viscosity  $\nu = 1.5 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$

density  $\rho = 1.2 \text{ kg m}^{-3}$

$$\sin (x \pm y)=\sin x \cos y \pm \cos x \sin y$$

$$\cos (x \pm y)=\cos x \cos y \mp \sin x \sin y$$

$$\cos 2 \theta=2 \cos ^2 \theta-1$$

$$\sin 2 \theta=2 \sin \theta \cos \theta$$

$$\sin 3 \theta=3 \sin \theta-4 \sin ^3 \theta$$

$$\cos 3 \theta=4 \cos ^3 \theta-3 \cos \theta$$

$$\int_0^{\pi} \cos \theta \mathrm{d} \theta=0$$


$$\int_0^{\pi} \sin \theta \mathrm{d} \theta=2$$

$$\int_0^{\pi} \cos ^2 \theta \mathrm{d} \theta=\int_0^{\pi} \sin ^2 \theta \mathrm{d} \theta=\frac{\pi}{2}$$


$$\int_0^{\pi} \frac{\cos n \theta}{\cos \theta-\cos \theta_1} \mathrm{d} \theta=\pi \frac{\sin n \theta_1}{\sin \theta_1} \quad n=0,1,2, \ldots$$

$$\int_0^{\pi} \frac{\sin n \theta \sin \theta}{\cos \theta-\cos \theta_1} \mathrm{d} \theta=-\pi \cos n \theta_1 \quad n=1,2,3, \ldots$$

1. Consider a clockwise and a counterclockwise potential vortex of strength  $G$  along  $y$  separated by a distance  $D$ .




- (a) Find the resultant complex potential  $w(z)$ .




- (b) Sketch the streamlines and equipotential lines in the above graph. Use a marker or colours (not red) to clearly distinguish between both and do not forget to add a legend.


- (c) Find the magnitude and direction of the velocity  $\vec{u}_q$  at point q in between the two vortices in function of the parameters  $G$  and  $D$ .




- (d) Now, add a horizontal uniform free stream flow with velocity  $u_\infty = \frac{1}{2}|\vec{u}_q|$  to the vortex pair, with  $\vec{u}_q$  the velocity found in the previous part. Find the new resultant complex potential  $w(z)$ .




- (e) Find the location of all stagnation points. Write your solution(s) in function of the parameters  $G$  and  $D$ .




(f) Sketch the new streamline pattern in the graph below.



(g) Find the pressure coefficient in point  $q$



(h) Find the total circulation  $\Gamma$  in the area enclosed by the large rectangle in the figure below.



(i) Find the total circulation  $\Gamma$  in the area enclosed by the circle in the figure below.

