



Most flows are unstable...



1. Intro-definitions
2. Rayleigh-Taylor
3. Rayleigh Plateau (destabilization through surface tension)
4. Rayleigh-Benard (convection)
5. Taylor Couette-Centrifugal instability
6. Kelvin-Helmholtz
7. Inflection point theorem Rayleigh ! Orr sommerfeld
8. transient growth
9. Spatial growth

SPATIALLY DEVELOPING SHEAR FLOWS



Flat plate boundary layer





Mixing layer





Cylinder wake



Plane channel flow



2D jet



2D PARALLEL FLOW CONCEPTS

Dispersion relation

2D vorticity equation

$$\left(\frac{\partial}{\partial t} + \frac{\partial \Psi}{\partial y} \frac{\partial}{\partial x} - \frac{\partial \Psi}{\partial x} \frac{\partial}{\partial y} \right) \nabla^2 \Psi = \frac{1}{Re} \nabla^4 \Psi$$

Basic flow + perturbation

$$\Psi(x, t) = \int U(y) dy + \psi(x, y, t)$$

Linear vorticity equation

$$\left(\frac{\partial}{\partial t} + U(y) \frac{\partial}{\partial x} \right) \nabla^2 \psi - U''(y) \frac{\partial \psi}{\partial x} = \frac{1}{Re} \nabla^4 \psi$$

Dispersion relation

$$D(k, \omega) = 0$$

Temporal approach:

k is real; ω is complex

Perturbation grow and decay in time!

Spatial approach:

ω is real; k is complex

Perturbations grow and decay in space!

Shear layer is inviscidly unstable!


Hyperbolic tangent mixing layer

$$U(y) = \bar{U} + \frac{\Delta U}{2} \tanh\left(\frac{2y}{\delta_\omega}\right)$$

$$\delta_\omega(x) \equiv \frac{(U_1 - U_2)}{(dU/dy)_{\max}}$$


Velocity ratio

$$R \equiv \frac{U_1 - U_2}{U_1 + U_2} = \frac{\Delta U}{2\bar{U}}$$



Why?

Only a necessary condition for instability!
Remember: Influence of confinement



$$R = 1$$

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

$$U(y; R) = 1 + R \tanh y$$


$$U_1(y) = \tanh y$$

Dispersion relation


$$\omega(k; R) = k + R \omega_1(k)$$

PARALLEL FLOW CONCEPTS


Effect of velocity ratio




$$R = 0$$



$$0 < R < 1$$



$$R = 1$$



$$R > 1$$

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Temporal approach

$$\omega_1(k) = i \omega_{1,i}(k)$$

$$\omega_i(k; R) = R \omega_{1,i}(k)$$

$$c_r = \omega_r / k = 1$$

Temporal approach: k is real; ω is complex

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Spatial approach

$$k + R \omega_1(k) = \omega$$

$$R \ll 1$$

$$-k_i(\omega, R) \sim R \omega_{1,i}(\omega)$$

Gaster transformation

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer


$$U(y) = \begin{cases} U_1, & y > \delta_\omega/2 \\ (U_1 + U_2)/2 + (U_1 - U_2)y/\delta_\omega, & |y| < \delta_\omega/2 \\ U_2, & y < -\delta_\omega/2 \end{cases}$$

$$\phi'' - k^2 \phi = 0$$

$$\phi_1(y) = A_1 e^{-ky}, \quad y > \delta_\omega/2,$$

$$\phi_2(y) = B_2 e^{ky}, \quad y < -\delta_\omega/2,$$

$$\phi_0(y) = A_0 e^{-ky} + B_0 e^{ky}, \quad |y| < \delta_\omega/2$$



2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$A_1 e^{-k\delta_\omega/2} = A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2},$$

$$B_2 e^{-k\delta_\omega/2} = A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2},$$

$$\begin{aligned} -k(U_1 - c)A_1 e^{-k\delta_\omega/2} &= k(U_1 - c)(-A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2}) \\ &\quad - \frac{\Delta U}{\delta_\omega}(A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2}), \end{aligned}$$

$$\begin{aligned} k(U_2 - c)B_2 e^{-k\delta_\omega/2} &= k(U_2 - c)(-A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2}) \\ &\quad - \frac{\Delta U}{\delta_\omega}(A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2}). \end{aligned}$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$\begin{aligned}-\frac{\Delta U}{\delta_\omega} A_0 e^{-k\delta_\omega/2} + \left[2k(U_1 - c) - \frac{\Delta U}{\delta_\omega} \right] B_0 e^{k\delta_\omega/2} &= 0 \\ \left[2k(U_2 - c) + \frac{\Delta U}{\delta_\omega} \right] A_0 e^{k\delta_\omega/2} + \frac{\Delta U}{\delta_\omega} B_0 e^{-k\delta_\omega/2} &= 0\end{aligned}$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$4(k\delta_\omega)^2(c - \bar{U})^2 - \left[(k\delta_\omega - 1)^2 - e^{-2k\delta_\omega} \right] \Delta U^2 = 0$$

$$k\delta_\omega \mapsto 2k, c/\bar{U} \mapsto c$$

$$4k^2(c - 1)^2 - R^2 \left[(2k - 1)^2 - e^{-4k} \right] = 0$$



$$c \equiv \frac{\omega}{k} = 1 \pm \frac{R}{2k} \left[(2k - 1)^2 - e^{-4k} \right]^{1/2}$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer



Temporal approach

$$2k_n - 1 = e^{-2k_n}$$



Spatial approach



$$R = 0.5$$





2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Temporal



Spatial



Michalke (1964, 65)

Solving a spatial instability problem

ex: Rayleigh equation

Back to temporal stability analysis!

How to solve Rayleigh equation for real k and complex ω ?

We fix k , we need to find all ω and ψ such that

$$k \left(U \left(\frac{d^2}{dy^2} - k^2 \right) - U''(y) \right) \psi = \omega \left(\frac{d^2}{dy^2} - k^2 \right) \psi$$
$$\psi(-L) = \psi(L) = 0$$

Formally,

$$\mathcal{A}\psi = c\mathcal{E}\psi$$

$$c = \omega/k$$


Discretize

$$\mathbf{A}\Psi = c\mathbf{E}\Psi$$

Generalized eigenvalue problem

How to solve Rayleigh equation for real k and complex ω ?

Finite differences of order 1



$$\Psi = \begin{pmatrix} \psi(y_1) \\ \psi(y_2) \\ \vdots \\ \psi(y_N) \\ \psi(y_{N+1}) \end{pmatrix} = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \vdots \\ \psi_N \\ \psi_{N+1} \end{pmatrix} \quad \Psi'' = \begin{pmatrix} \psi''(y_1) \\ \psi''(y_2) \\ \vdots \\ \psi''(y_N) \\ \psi''(y_{N+1}) \end{pmatrix}$$

How to solve Rayleigh equation for real k and complex ω ?

Finite differences

$$\begin{pmatrix} \psi_2'' \\ \psi_3'' \\ \psi_4'' \\ \vdots \\ \psi_{N-3}'' \\ \psi_{N-2}'' \\ \psi_{N-1}'' \end{pmatrix} = \begin{pmatrix} -2 & 1 & 0 & \dots & \dots & \dots & 0 \\ 1 & -2 & 1 & 0 & \dots & \dots & 0 \\ 0 & 1 & -2 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 & -2 & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 & -2 & 1 \\ 0 & \dots & \dots & \dots & 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} \psi_2 \\ \psi_3 \\ \psi_4 \\ \vdots \\ \psi_{N-3} \\ \psi_{N-2} \\ \psi_{N-1} \end{pmatrix}$$

Sparse matrix but low order!

How to solve Rayleigh equation for complex k and real ω ?

We fix ω , we need to find all k and ψ such that

$$k \left(U \left(\frac{d^2}{dy^2} - k^2 \right) - U''(y) \right) \psi = \omega \left(\frac{d^2}{dy^2} - k^2 \right) \psi$$
$$\psi(-L) = \psi(L) = 0$$


Formally,

$$(A_0(\omega, y) + kA_1(\omega, y) + k^2A_2(\omega, y) + k^3A_3(\omega, y)) \psi = 0$$

Polynomial eigenvalue problem

Many more eigenvalues (for Rayleigh equation: 3 x more!)

$$U=1+0.9*\tanh(y); \omega=0.4; L=5$$



Which of these waves are unstable?


$\text{Im}(k) < 0?$

$\text{Im}(k) > 0?$

Recall : $\exp(i(kx - \omega t))$

The stability of a spatial wave can be only determined if one knows in which direction it propagates!

k^+ waves propagate towards positive x



k^- waves propagate towards negative x




However, determining this direction of propagation is particularly difficult, except in the case of a temporally stable flow.


The addition of a positive imaginary offset to the frequency makes the temporal problem stable!

This separates the spatial waves into k^+ and k^- waves.

offset spatial stability




spatial stability: $\text{Im}(\omega)=0$



The branches are then followed by continuity

Validity of Gaster transformation?

$R=0.4$




$$R \ll 1$$

$$-k_i(\omega, R) \sim R \omega_{1,i}(\omega)$$

Validity of Gaster transformation?

R=0.9


x: Gaster transformation




$$-k_i(\omega, R) \sim R \omega_{1,i}(\omega)$$

Spatio-temporal instability theory


👉 Convective instability




✚ Absolute instability




👉 Convective instability




F Absolute instability



Spectrum



👉 amplifier




👉 oscillator


Mixing layer experiments by Strikowsky and Niccum (1991)

Spatio-temporal instability theory

👉 Convective instability



F Absolute instability



We need to generalize the concept of group velocity since ω (and why not k) is complex

For neutral waves, the group velocity is $d\omega/dk$

Here this quantity is the derivative of a complex function with respect to a complex variable. Cauchy-Rieman conditions apply.

Spatio-temporal spectral analysis

Inverse Fourier Transform

$$u(x,t) = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \hat{u}(k, \omega) e^{i(kx - \omega t)} dk d\omega$$

$$\hat{u}(k, \omega) = (2\pi)^2 \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} u(x, t) e^{-i(kx - \omega t)} dx dt$$

Direct Fourier Transform

Spatio-temporal spectral analysis

Inverse Fourier Transform

$$u(x,t) = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \hat{u}(k, \omega) e^{i(kx - \omega t)} dk d\omega$$

Use dispersion relation $\omega(k)$!

Dispersion relation

$$\omega^2 = \tanh(kH) \left(\frac{\gamma k^3}{\rho} + gk \right)$$

Capillary wavenumber: $k_c = \sqrt{\rho g / \gamma}$

Length scale: $\tilde{k} = k / k_c$

Time scale: $\tilde{\omega} = \omega / \sqrt{g k_c}$

One single non-dimensional parameter $\tilde{H} = H k_c$


$$\tilde{\omega}^2 = \tanh(\tilde{k} \tilde{H}) \left(\tilde{k}^3 + \tilde{k} \right)$$

Dispersion relation


$$\tilde{\omega}^2 = \tanh(\tilde{k}\tilde{H}) \left(\tilde{k}^3 + \tilde{k} \right)$$

	gravity $\tilde{k} \ll 1$	capillary $\tilde{k} \gg 1$
shallow water $\tilde{k} \ll 1/\tilde{H}$	$\pm \tilde{k} \sqrt{\tilde{H}}$	$\pm \tilde{k}^2 \sqrt{\tilde{H}}$
Deep water $\tilde{k} \gg 1/\tilde{H}$	$\pm \sqrt{\tilde{k}}$	$\pm \tilde{k} \sqrt{\tilde{k}}$


Difference between group velocity v and phase velocity c




Spatio-temporel diagram



Spatio-temporal diagram



Spatio-temporal diagram




$$v_g = \frac{3}{2} v_\phi$$

Dispersion relation

	gravity $\tilde{k} \ll 1$	capillary $\tilde{k} \gg 1$
shallow water $\tilde{k} \ll 1/\tilde{H}$	$\omega_{\text{shallow/gravity}} \sim \pm k \sqrt{gH}$ $c_{\text{shallow/gravity}} \sim \pm \sqrt{gH}$ $v_{\text{shallow/gravity}} \sim \pm \sqrt{gH}$	$\omega_{\text{shallow/capillary}} \sim \pm k^2 \sqrt{\gamma H / \rho}$ $c_{\text{shallow/capillary}} \sim \pm k \sqrt{\gamma H / \rho}$ $v_{\text{shallow/capillary}} \sim \pm 2k \sqrt{\gamma H / \rho}$
Deep water $\tilde{k} \gg 1/\tilde{H}$	$\omega_{\text{deep/gravity}} \sim \pm \sqrt{gk}$ $c_{\text{deep/gravity}} \sim \pm \sqrt{\frac{g}{k}}$ $v_{\text{deep/gravity}} \sim \pm \frac{1}{2} \sqrt{\frac{g}{k}}$	$\omega_{\text{deep/capillary}} \sim \pm k^{3/2} \sqrt{\gamma / \rho}$ $c_{\text{deep/capillary}} \sim \pm k^{1/2} \sqrt{\gamma / \rho}$ $v_{\text{deep/capillary}} \sim \pm 3/2 k^{1/2} \sqrt{\gamma / \rho}$

Dispersion relation



Spectral analysis

Fourier transform:

$$u(x, t) = \frac{1}{2} \int_0^{+\infty} \hat{u}(k) e^{i(kx - \omega(k)t)} dk + c.c.$$

Carrier/enveloppe

Carrier/enveloppe :

$$u(x, t) = \frac{1}{2} A(x, t) e^{i(k_0 x - \omega_0 t)} + c.c.$$

Spectral analysis

Fourier transform:

$$u(x, t) = \frac{1}{2} \int_0^{+\infty} \hat{u}(k) e^{i(kx - \omega(k)t)} dk + c.c.$$

Carrier/enveloppe :

$$u(x, t) = \frac{1}{2} A(x, t) e^{i(k_0 x - \omega_0 t)} + c.c.$$

Spectral analysis

Fourier transform:

$$u(x, t) = \frac{1}{2} \int_0^{+\infty} \hat{u}(k) e^{i(kx - \omega(k)t)} dk + c.c.$$

Carrier/enveloppe :

$$u(x, t) = \frac{1}{2} A(x, t) e^{i(k_0 x - \omega_0 t)} + c.c.$$

Enveloppe :

$$A(x, t) = \int_0^{\infty} \hat{u}(k) e^{i(k - k_0)x - i(\omega - \omega_0)t} dk.$$

Spectral analysis at time=0

Fourier transform:

$$u(x, 0) = \frac{1}{2} \int_0^{+\infty} \hat{u}(k) e^{i(kx)} dk + \text{c.c.}$$

$\hat{u}(k)$ is given by Fourier transform at time $t=0$

Envelope :

$$A(x, 0) = \int_0^{\infty} \hat{u}(k) e^{i(k-k_0)x} dk + \text{c.c.}$$


Spectral analysis

Gaussian spectrum: $\hat{u}(k) = u_0 e^{-\sigma^2(k-k_0)^2}$


Initial envelope : $A(x, 0) = \frac{u_0 \sqrt{\pi}}{2\sigma} e^{-\frac{x^2}{4\sigma^2}}$

Gaussian spectrum


spectrum



wave



Gaussian wavepackets



$u_0(x)$	$\hat{u}_0(k_1)$
$\exp\left(-\frac{x^2}{2\sigma^2}\right)$	$\frac{\sigma}{\sqrt{2\pi}} \exp\left(-\frac{\sigma^2 k^2}{2}\right)$
$\delta(x)$	$\frac{1}{2\pi}$

Spectral analysis

Initial envelope :

$$A(x, 0) = \frac{u_0 \sqrt{\pi}}{2\sigma} e^{-\frac{x^2}{4\sigma^2}}$$

Gaussian spectrum:

$$\hat{u}(k) = u_0 e^{-\sigma^2(k-k_0)^2}$$

Spectral analysis

Initial envelope :

$$A(x, 0) = \frac{u_0 \sqrt{\pi}}{2\sigma} e^{-\frac{x^2}{4\sigma^2}}$$

Gaussian spectrum:

$$\hat{u}(k) = u_0 e^{-\sigma^2(k-k_0)^2}$$

Evolution of envelope :

$$A(x, t) = \int_0^\infty \hat{u}(k) e^{i(k-k_0)x - i(\omega - \omega_0)t} dk.$$

Spectral analysis

Initial envelope :

$$A(x, 0) = \frac{u_0 \sqrt{\pi}}{2\sigma} e^{-\frac{x^2}{4\sigma^2}}$$

Gaussian spectrum:

$$\hat{u}(k) = u_0 e^{-\sigma^2(k-k_0)^2}$$

Evolution of envelope :

$$A(x, t) = \int_0^\infty \hat{u}(k) e^{i(k-k_0)x - i(\omega - \omega_0)t} dk.$$

Definition group velocity


$$\omega - \omega_0 = c_g(k - k_0), \quad c_g = \frac{\partial \omega}{\partial k}(k_0)$$

Spectral analysis

Definition of group velocity $\omega - \omega_0 = c_g(k - k_0), \quad c_g = \frac{\partial \omega}{\partial k}(k_0)$

$$A(x, t) = \frac{u_0 \sqrt{\pi}}{2\sigma} e^{-\frac{(x - c_g t)^2}{4\sigma^2}}$$

Group velocity



Wavepacket


Spectral analysis

Higher order
development

$$\omega - \omega_0 = c_g(k - k_0) + \frac{\omega_0''}{2}(k - k_0)^2$$
$$c_g = \frac{\partial \omega}{\partial k}(k_0), \quad \omega_0'' = \frac{\partial^2 \omega}{\partial k^2}(k_0)$$


$$A(x, t) = \frac{u_0}{2} \sqrt{\frac{\pi}{\sigma^2 + \frac{1}{2}i\omega_0''t}} \exp\left(-\frac{(x - c_g t)^2}{4(\sigma^2 + \frac{1}{2}i\omega_0''t)}\right)$$

Wave packet dispersion




Onde correspondant à l'enveloppe $\hat{u}(k)$ pour $\sigma^{-1}k_0 = 0,1$ et $\omega_0'' = 4c_g/k_0$: (a), instant initial $t = 0$; (b), $c_g t = 100/k_0$.


Kelvin's wake




Gravity waves created by a ship




Gravity waves created by a ship




Gravity waves created by a ship



Gravity waves created by a ship




Gravity waves created by a ship



Gravity waves created by a ship

$$\sin(\alpha)/OG = \cos(\theta)/AG$$

$$\sin(\theta - \alpha)AG = GC = OG$$




$$\Rightarrow \sin(\alpha) = \cos(\theta) \sin(\theta - \alpha)$$


$$\Rightarrow \sin(\alpha) = \cos(\theta) (\sin(\theta) \cos(\alpha) + \cos(\theta) \sin(\alpha))$$

$$\Rightarrow \tan(\alpha) = \cos(\theta) \sin(\theta) / (1 + \cos^2(\theta))$$



Gravity waves created by a ship




Gravity waves created by a ship



Gravity waves created by a ship




But observations show



Moisy and Rabaud 2013

But observations show




Moisy and Rabaud 2013

Generalization: Spatio-temporal instability theory


First find the zero group velocity wave: $d\omega/dk=0 \Rightarrow (k_0, \omega_0)$
and consider the sign of $\text{Im}(\omega_0)$

👉 Convective instability



$$\text{Im}(\omega_0) < 0$$


✖ Absolute instability



$$\text{Im}(\omega_0) > 0$$

LINEAR IMPULSE RESPONSE: ABSOLUTE/CONVECTIVE INSTABILITY

Green's function or impulse response



Briggs (1964) Bers (1983)
Huerre and Monkewitz (1985)

LINEAR IMPULSE RESPONSE: ABSOLUTE/CONVECTIVE INSTABILITY

Linearly stable flow


$$\lim_{t \rightarrow \infty} G(x, t) = 0 \quad \text{along all rays } x/t = \text{const.}$$

Linearly unstable flow

$$\lim_{t \rightarrow \infty} G(x, t) = \infty \quad \text{along at least one ray } x/t = \text{const.}$$

LINEAR IMPULSE RESPONSE: ABSOLUTE/CONVECTIVE INSTABILITY

Green's function or impulse response



Briggs (1964) Bers (1983)
Huerre and Monkewitz (1985)

LINEAR IMPULSE RESPONSE: ABSOLUTE/CONVECTIVE INSTABILITY

Convectively unstable flow


$$\lim_{t \rightarrow \infty} G(x, t) = 0 \quad \text{along the ray } x/t = 0$$

Absolutely unstable flow


$$\lim_{t \rightarrow \infty} G(x, t) = \infty \quad \text{along the ray } x/t = 0$$

ANALYSIS IN COMPLEX FOURIER SPACE: AU/CU CRITERION

Temporal growth rate « at velocity v »



Convective instability



Absolute instability

ANALYSIS IN COMPLEX FOURIER SPACE: AU/CU CRITERION

Important notions

Absolute wavenumber k_0 and frequency $\omega_0 = \omega(k_0)$
observed along ray $v = 0$, i.e. for a stationary observer,
defined by


$$\frac{\partial \omega}{\partial k}(k_0) = 0$$

Absolute growth rate is

$$\sigma(0) = \omega_{0,i}$$

Isovaleurs de ω_i

Absolute frequency ω_0 : Saddle point condition



ANALYSIS IN COMPLEX FOURIER SPACE: AU/CU CRITERION

Instability criteria

$\omega_{i,max} < 0$ linearly stable

$\omega_{i,max} > 0$ linearly unstable

$\omega_{0,i} < 0$ convectively unstable

$\omega_{0,i} > 0$ absolutely unstable

Hyperbolic tangent mixing layer


$$U(y) = \bar{U} + \frac{\Delta U}{2} \tanh \left(\frac{2y}{\delta_\omega} \right)$$

$$\delta_\omega(x) \equiv \frac{(U_1 - U_2)}{(dU/dy)_{\max}}$$

Velocity ratio


$$R \equiv \frac{U_1 - U_2}{U_1 + U_2} = \frac{\Delta U}{2\bar{U}}$$

$$U(y; R) = 1 + R \tanh y$$




APPLICATION TO MIXING LAYERS

Locus of complex absolute frequency




H.&Monkewitz (1985)

Influence of countercurrent shear on turbulence level




Influence of countercurrent shear on turbulence level

Base flow




a) Single stream shear layer




b) Countercurrent shear layer

Influence of countercurrent shear on turbulence level

Turbulence intensity





a) Single-stream shear layer



b) Countercurrent shear layer

THE MIXING LAYER: SHIFT TO OSCILLATOR !




Strykowski & Niccum (1991)

Direct Numerical Simulations with top-hat profile at inlet


Viscous diffusion $\longrightarrow$ Non-parallel flow

■ $\Lambda_{loc} = (U_{max} - U_{min}) / (U_{max} + U_{min})$


■ $\delta = (U_{max} - U_{min}) / (|dU/dy|_{max})$



Vorticity field: $Re = 100$, $h = 1$




$$\Lambda = -0.739$$




$$\Lambda = -0.667$$

An increase in Λ (more coflow) advects the perturbation

Vorticity field: $Re = 100$, $\Lambda = -0.739$




$h=1$




$h=3$

Destabilizing influence of confinement!

Spatio-temporal diagram, $h=1$ and $\Lambda = -0.667$



Spatio-temporal diagram, $h=1$ and $\Lambda = -0.739$



THE BLUFF BODY WAKE: A TYPICAL FLOW OSCILLATOR



$Re = 140$
Periodic
flow


Taneda (1982)

ABSOLUTE/CONVECTIVE INSTABILITY IN PARALLEL WAKES

Family of wake profiles

$$U(y) = U_{\infty} + (U_c - U_{\infty}) U_1 \left(\frac{y}{\delta}; N \right)$$


$$U_1(\xi; N) = [1 + \sinh^{2N} \{ \xi \sinh^{-1}(1) \}]^{-1}$$



Monkewitz (1988)

ABSOLUTE/CONVECTIVE INSTABILITY IN PARALLEL WAKES

Family of wake profiles




Effect of velocity ratio R

Effect of N

ABSOLUTE/CONVECTIVE INSTABILITY IN PARALLEL 2D PARALLEL FLOW CONCEPTS

$\text{sech}^2 y$ wake

Temporal approach




Betchov & Criminale (1966)


2D PARALLEL FLOW CONCEPTS

$\text{sech}^2 y$ wake

Sinuuous and varicose modes




sinuous



varicose

ABSOLUTE/CONVECTIVE INSTABILITY IN PARALLEL WAKES

Effect of steepness, velocity ratio and Reynolds number



Monkewitz (1988)

LOCAL INSTABILITY BEHAVIOR OF CYLINDER WAKE

$$5 < Re < 25$$

Convective instability

$$25 < Re < 48.5$$

Absolute instability