


## **2.3 Formation of Multiple Bonds**

# Reading Recommendations

- Clayden, Greeves, Warren, *Organic Chemistry*, Oxford University Press, 2<sup>nd</sup> ed., **2012**, pp 99–106.
- Jamart, Bodiguel, Brosse, *Les cours de Paul Arnaud - Cours de chimie organique*, Dunod, 19<sup>th</sup> ed., **2015**, pp 79–86.


# Formation of Carbon-Carbon Multiple Bonds



- double or triple bonds are one  $\sigma$  bond plus one or two  $\pi$ -bonds, respectively
- $\pi$  bonds between residual p orbitals, node plane along bond, no rotational symmetry
- rotation around  $\pi$ -bonds requires breaking them, energetically too costly at r. t.

# Molecular Orbital View of the Carbon-Carbon Double Bond


- simplified and schematic molecular orbital energy diagram of the ethene molecule



- only orbitals of **matching symmetry & orientation** interact,  **$sp^2$  with  $sp^2$** , and  **$p_z$  with  $p_z$**
- distinct  **$\sigma$  bond (from two  $sp^2$ )** and  **$\pi$  bond (from two  $p_z$ )** with different energy, symmetry
- chemistry ruled by highest occupied, lowest unoccupied molecular orbitals (HOMO, LUMO)
- typically  **$\pi$  HOMO and  $\pi^*$  LUMO**, located between  $\sigma$  and  $\sigma^*$  because  $p_z$  overlap much smaller


# Rotation Around a Double Bond

- $\sigma$  bond has rotational symmetry relative to carbon-carbon bond axis, but  $\pi$  bond does not



- rotating  $\pi$  orbitals by  $90^\circ$  requires breaking the  $\pi$  bond ( $\approx 260$  kJ/mol), unfavorable at r.t.


- simplified and schematic molecular orbital energy diagram of the ethyne molecule



- only orbitals of **matching symmetry & orientation** interact ( $sp$  with  $sp$ ,  $p_z$  with  $p_z$ ,  $p_y$  with  $p_y$ )
- distinct  **$\sigma$ -bond (from two  $sp^2$ )** and **two independent  $\pi$ -bonds (each from two  $p_z$  and two  $p_y$ )**
- the two  $\pi$ -bonds are identical in energy, symmetry, just different in orientation
- chemistry dominated by  **$\pi$  HOMO and  $\pi^*$  LUMO**, reactivity of ethyne similar to ethene

# Molecular Orbital View of the Carbon-Oxygen Double Bond in the Carbonyl Group

- simplified and schematic molecular orbital energy diagram of the carbon-oxygen double bond



- $\sigma$  bond from  $sp^2(\text{C})$  with  $sp^2(\text{O})$ ,  $\pi$ -bond from  $p_z(\text{C})$  with  $p_z(\text{O})$  with different bond energies
- oxygen orbitals lower in energy due to electronegativity difference of carbon and oxygen
- bonding  $\sigma$  orbital and, in particular, bonding  $\pi$  orbital “look like” oxygen orbitals
- both the  $\sigma$  bond and, in particular, the more polarizable  $\pi$  bond polarized towards oxygen

# Learning Outcomes

- **multiple bonds can be described as distinct  $\sigma$  bond plus one or two  $\pi$  bonds**
  - $\sigma$  bond has rotational symmetry with respect to carbon-carbon bond axis
  - $\pi$  bond does not have rotational symmetry; rotation requires breaking it
- **chemistry & physics dominated by frontier orbitals:  $\pi$  HOMO and  $\pi^*$  LUMO**