



ELECTRONS IN CRYSTALS

Chris J. Pickard




ELECTRONS IN CRYSTALS



Cartoon of electrons (blue) in motion

- The electrons in a crystal experience a potential with the periodicity of the Bravais lattice: $U(\mathbf{r} + \mathbf{R}) = U(\mathbf{r})$
- The scale of the periodicity is of the order of the de Broglie wavelength of an electron — $1\text{\AA}$ — so we must use Quantum Mechanics
- Of course, the periodicity is an idealisation: impurities, defects, thermal vibrations, finite size effects

THE PERIODIC POTENTIAL



A 1D periodic crystalline potential

- In principle, we are faced with a many electron problem
- But we can make a lot of progress using the *independent electron* approximation
- We investigate the properties of the Schrödinger equation for a single electron:
$$H\Psi = \left(-\frac{\hbar^2}{2m}\nabla^2 + U(\mathbf{r})\right)\Psi = E\Psi$$
with $U(\mathbf{r} + \mathbf{R}) = U(\mathbf{r})$


BLOCH'S THEOREM

$$\Psi_{n\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}} u_{n\mathbf{k}}(\mathbf{r})$$

$$u_{n\mathbf{k}}(\mathbf{r} + \mathbf{R}) = u_{n\mathbf{k}}(\mathbf{r})$$

$$\Psi_{n\mathbf{k}}(\mathbf{r} + \mathbf{R}) = e^{i\mathbf{k}\cdot\mathbf{R}} \Psi_{n\mathbf{k}}(\mathbf{r})$$

- Independent electrons which obey the one electron Schrödinger equation for a periodic potential are called *Bloch electrons* and obey Bloch's theorem
- Bloch's theorem can be written in two equivalent forms



PROOF OF BLOCH'S THEOREM

Consider the translation operator:

$$T_{\mathbf{R}}f(\mathbf{r}) = f(\mathbf{r} + \mathbf{R})$$

It forms a commuting set for all $\mathbf{R}$ and H :

$$T_{\mathbf{R}}H\Psi(\mathbf{r}) = H(\mathbf{r} + \mathbf{R})\Psi(\mathbf{r} + \mathbf{R}) = H(\mathbf{r})\Psi(\mathbf{r} + \mathbf{R}) = HT_{\mathbf{R}}\Psi(\mathbf{r})$$

$$T_{\mathbf{R}}H = HT_{\mathbf{R}}$$

$$T_{\mathbf{R}}T_{\mathbf{R}'} = T_{\mathbf{R}'}T_{\mathbf{R}} = T_{\mathbf{R}+\mathbf{R}'}$$

The eigenstates of H are simultaneous eigenstates of all $T_{\mathbf{R}}$:

$$H\Psi(\mathbf{r}) = E\Psi(\mathbf{r})$$

$$T_{\mathbf{R}}\Psi(\mathbf{r}) = c(\mathbf{R})\Psi(\mathbf{r})$$

The properties of $T_{\mathbf{R}}$ imply a relationship between the eigenvalues:

$$T_{\mathbf{R}}T_{\mathbf{R}'}\Psi(\mathbf{r}) = c(\mathbf{R})T_{\mathbf{R}'}\Psi(\mathbf{r}) = c(\mathbf{R})c(\mathbf{R}')\Psi(\mathbf{r})$$

$$T_{\mathbf{R}}T_{\mathbf{R}'}\Psi(\mathbf{r}) = T_{\mathbf{R}+\mathbf{R}'}\Psi(\mathbf{r}) = c(\mathbf{R} + \mathbf{R}')\Psi(\mathbf{r})$$

and so:

$$c(\mathbf{R})c(\mathbf{R}') = c(\mathbf{R} + \mathbf{R}')$$

If $\mathbf{a}_i$ are the primitive lattice vectors, we can always write:

$$c(\mathbf{a}_i) = e^{2\pi i x_i}$$

For an arbitrary Bravais lattice vector:

$$\mathbf{R} = n_1 \mathbf{a}_1 + n_2 \mathbf{a}_2 + n_3 \mathbf{a}_3$$

and so, considering repeated applications of $T_{\mathbf{a}_i}$:


$$c(\mathbf{R}) = c(\mathbf{a}_1)^{n_1} c(\mathbf{a}_2)^{n_2} c(\mathbf{a}_3)^{n_3} = e^{i\mathbf{k} \cdot \mathbf{R}}$$

where $\mathbf{b}_i \cdot \mathbf{a}_j = 2\pi\delta_{ij}$ and $\mathbf{k} = x_1 \mathbf{b}_1 + x_2 \mathbf{b}_2 + x_3 \mathbf{b}_3$

We arrive at the second form of Bloch's Theorem:

$$T_{\mathbf{R}}\Psi(\mathbf{r}) = \Psi(\mathbf{r} + \mathbf{R}) = c(\mathbf{R})\Psi(\mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{R}}\Psi(\mathbf{r})$$


THE FIRST BRILLOUIN ZONE



The first FCC Brillouin zone

- The wave vector $\mathbf{k}$ can always be confined to the first Brillouin zone (or any primitive cell of the reciprocal lattice)
- Any $\mathbf{k}'$ not in the first Brillouin zone can be written as: $\mathbf{k}' = \mathbf{k} + \mathbf{K}$, where $\mathbf{k}$ is in the first Brillouin zone and $e^{i\mathbf{K} \cdot \mathbf{R}} = 1$
- The labels K,L,W,X and Γ are high symmetry points in the Brillouin zone

BAND STRUCTURE



- For a given $\mathbf{k}$ there many solutions to the Schrödinger equation:
$$H_{\mathbf{k}}u_{\mathbf{k}} = E_{\mathbf{k}}u_{\mathbf{k}}(\mathbf{r}), u_{\mathbf{k}}(\mathbf{r}) = u_{\mathbf{k}}(\mathbf{r} + \mathbf{R})$$
- The boundary condition ensure that there are many (labelled n) discretely spaced eigenvalues
- The Hamiltonian depends on $\mathbf{k}$ as a parameter, and so the eigenvalues vary continuously with wave vector for a given n . Hence, they are *bands*


CRYSTAL MOMENTUM

- For Bloch electrons $\mathbf{k}$ is not proportional to electronic momentum $\mathbf{p}$, and known as *crystal momentum*

$$\begin{aligned}\frac{\hbar}{i}\nabla\Psi_{n\mathbf{k}} &= \frac{\hbar}{i}\nabla(e^{i\mathbf{k}\cdot\mathbf{r}}u_{n\mathbf{k}}(\mathbf{r})) \\ &= \hbar\mathbf{k}\Psi_{n\mathbf{k}} + e^{i\mathbf{k}\cdot\mathbf{r}}\frac{\hbar}{i}\nabla u_{n\mathbf{k}}\end{aligned}$$

- The $\Psi_{n\mathbf{k}}$ are not momentum eigenstates
- However, $\hbar\mathbf{k}$ is a natural extension of
- The dynamical significance of $\hbar\mathbf{k}$ is revealed by considering electrons response to applied electromagnetic fields
- A quantum number characteristic of the translational symmetry of the periodic potential, as $\mathbf{p}$ is characteristic of the full translational symmetry of free space

VELOCITY AND EFFECTIVE MASS



- The velocity of an electron at $\mathbf{k}$ in band n is given by the gradient of the band and the inverse effective mass is given by the curvature
- The velocity operator is:

$$\mathbf{v} = d\mathbf{r}/dt = (1/i\hbar)[\mathbf{r}, H] = \mathbf{p}/m$$

$$= \hbar \nabla / im$$
- Electrons in a perfect crystal move at a constant mean velocity

DENSITY OF STATES

- Many electronic properties are weighted sums over the electronic levels of the form:

$$Q = 2 \sum_{n\mathbf{k}} Q_n(\mathbf{k})$$

which is an integral in a large crystal:

$$q = 2 \sum_n \int \frac{d\mathbf{k}}{(2\pi)^3} Q_n(\mathbf{k})$$

- Often $Q_n(\mathbf{k})$ depends only on n and $\mathbf{k}$ through $E_n(\mathbf{k})$, and the *density of states* $g(E) = \sum_n g_n(E)$ is a useful construct:

$$q = \int dE g(E) Q(E)$$

- The density of states of a band is:

$$g_n(E) = \int \frac{d\mathbf{k}}{4\pi^3} \delta(E - E_n(\mathbf{k}))$$

- It can be written as a surface integral:



$$g_n(E) = \int_{S_n(E)} \frac{dS}{4\pi^3} \frac{1}{|\nabla E_n(\mathbf{k})|}$$

with $S_n(E)$ a surface of constant energy



VAN HOVE SINGULARITIES

- Because $E_n(\mathbf{k})$ is periodic in reciprocal space, and for each n bounded from above and below, and differentiable everywhere there must be $\mathbf{k}$ for which $|\nabla E| = 0$
- In 1D this results in a divergence of the density of states itself
- In 3D the divergence is integrable, and results in discontinuities in dg_n/dE
- Thus, the integrand in the expression for $g_n(E)$ diverges
- These are the *van Hove singularities*

METALS AND INSULATORS




- Fill the electronic states, lowest energy first across the whole first Brillouin zone (so that each level is counted only once), until all the electrons are accommodated




- If there is a *gap* between the highest occupied state and the lowest unoccupied the crystal is an *insulator* (and a called a *semiconductor* if the gap is close to $k_B T$)

THE FERMI SURFACE




- If there is no gap then the crystal is a metal
- There will be a surface in k-space separating occupied from unoccupied levels: this is known as the *Fermi surface* and may consist of several *branches*. It determines the transport and optical properties of the metal

THE FERMI SURFACE




- If there is no gap then the crystal is a metal
- There will be a surface in k-space separating occupied from unoccupied levels: this is known as the *Fermi surface* and may consist of several *branches*. It determines the transport and optical properties of the metal

ELECTRONS IN A REALISTIC CRYSTAL POTENTIAL



- In principle, it remains only to solve the Schrödinger's equation for the Bloch wavefunctions
- This might be viewed as a job of pure numerics – aside from the choice of $U(\mathbf{r})$
- But we can do better than brute force – and with greater understanding


THE NEARLY FREE ELECTRON APPROXIMATION



Defining near degeneracy

- Possibly suprisingly, the electronic structure of some metals is considered to arise from a weak periodic perturbation of the free electron gas
- This is due to the combined effects of the Pauli exclusion principle and *screening*
- The perturbation has different results, depending whether the free electron states are nearly degenerate or not


THE NEARLY FREE ELECTRON APPROXIMATION



Splitting at a single Bragg plane


- If there is no degeneracy, the perturbation is second order in U_K
- If there is near degeneracy, the perturbation can be linear in the potential
- Symmetries and structure factor effects can eliminate the splitting – returns with *spin-orbit coupling*
- The bandstructure can be plotted in the *reduced* or *extended* or *repeated* zone schemes

FREE ELECTRON AND A REAL BANDSTRUCTURE



The free electron and DFT bandstructure of FCC Aluminium


THE TIGHT BINDING APPROXIMATION



*Sodium electronic wavefunctions –
separated by $3.7\text{\AA}$*

- Now, let's think of the electronic structure as being a modification of that of the isolated constituent atoms
- This is a good picture if the overlap of the atomic wavefunctions is small
- Clearly, this is not the case for metallic sodium – it is a nearly free electron metal

THE TIGHT BINDING APPROXIMATION



*When $r\phi(\mathbf{r})$ is large, $\Delta U(\mathbf{r})$ is small
and vice versa*

- To calculate the corrections, consider:
$$H = H_{\text{at}} + \Delta U(\mathbf{r})$$
- The general form of the wavefunction is: $\psi(r) = \sum_{\mathbf{R}} e^{i\mathbf{k} \cdot \mathbf{R}} \phi(\mathbf{r} - \mathbf{R})$ where $\phi(\mathbf{r}) = \sum_n b_n \psi_n(\mathbf{r})$
- There is a strong *hybridisation* and splitting of levels close to each other in energy – recall the nearly free electron model

THE TIGHT BINDING s-BAND

Consider a single s-band: $|\psi_{\mathbf{k}}\rangle = \sum_{\mathbf{R}} e^{i\mathbf{k}\cdot\mathbf{R}}|\psi_s\rangle$


Multiply: $H|\psi_{\mathbf{k}}\rangle = (H_{\text{at}} + \Delta U)|\psi_{\mathbf{k}}\rangle = \mathcal{E}_{\mathbf{k}}|\psi_{\mathbf{k}}\rangle$ through by $\langle\psi_s|$ and integrate using $\langle\psi_s|H_{\text{at}}|\psi_s\rangle = E_s$:

$$(\mathcal{E}_{\mathbf{k}} - E_s)\langle\psi_s|\psi_{\mathbf{k}}\rangle = \langle\psi_s|\Delta U|\psi_{\mathbf{k}}\rangle \Rightarrow \mathcal{E}_{\mathbf{k}} = E_s + \frac{\langle\psi_s|\Delta U|\psi_{\mathbf{k}}\rangle}{\langle\psi_s|\psi_{\mathbf{k}}\rangle}$$

Ignoring the deviations from unity of the denominator, summing over the nearest neighbours only, and using the inversion symmetry of the potential:

$$\mathcal{E}_{\mathbf{k}} = E_s + \langle\psi_s|\Delta U|\psi_s\rangle + \sum_{nn} \cos(\mathbf{k} \cdot \mathbf{R}) \int \psi_s^*(\mathbf{r}) \Delta U(\mathbf{R}) \psi_s(\mathbf{r} - \mathbf{R}) d\mathbf{r}$$

THE TIGHT BINDING S-BAND



Sodium in a Simple Cubic cell of side $3.7\text{\AA}$:

$$\mathcal{E}_{\mathbf{k}} = \alpha - \beta(\cos ak_x + \cos ak_y + \cos ak_z)$$