

Lecture 14

- Review
- Lorentz transformations of EM fields
- Kramers-Kronig relations, dispersion

- We return to the study of EM fields in media, but at a more advanced level

- absorption

- non-locality in time

- convolution and frequency

- analytic properties

- kramers-kronig relations

- example: resonant absorption model. →

- very general!

- Let us remember the following effect:
polarisation of dielectric media
by external electric field

$$\vec{P}(\vec{x}) = \sum_n \vec{d}_n \cdot f(\vec{x}_n - \vec{x}) \quad \rightarrow \text{smearing function}$$

↓ electric polarization of the medium.

$$\langle \eta \rangle(\vec{x}) = \rho(\vec{x}) - \vec{\nabla}_x \cdot \vec{P}(\vec{x})$$

From this we derived averaged Maxwell equations:

$$\vec{\nabla} \cdot \vec{E} = \frac{\rho}{\epsilon_0} - \frac{\vec{\nabla} \cdot \vec{P}}{\epsilon_0}$$

We also defined the Displacement vector

$$\vec{D} = \epsilon_0 \vec{E} + \vec{P},$$

And we did a similar treatment for the magnetic phenomenon

$$\vec{H} = \frac{1}{\mu_0} \vec{B} - \vec{M} \quad \rightarrow \text{magnetisation}$$

The Maxwell equations eventually read:

$$\vec{\nabla} \cdot \vec{D} = \rho$$

$$\vec{\nabla} \times \vec{E} + \frac{\partial \vec{B}}{\partial t} = 0$$

$$\vec{\nabla} \times \vec{H} - \frac{\partial \vec{D}}{\partial t} = \vec{J}$$

$$\vec{\nabla} \cdot \vec{B} = 0$$

To make use of them we need the relation between $\vec{E}$ and $\vec{D}$ and $\vec{H}$ and $\vec{B}$ which we chose as

$$\vec{D}(\vec{x}, t) = \epsilon \vec{E}(\vec{x}, t)$$

$$\vec{H}(\vec{x}, t) = \frac{1}{\mu} \vec{B}(\vec{x}, t)$$

This result is correct for static fields, however, we also used it for our discussion of EM waves. In the case of time-dependant fields there are two new effects:

→ absorption (ϵ and μ have imaginary parts)

→ dispersion (relation between $\vec{E}$ and $\vec{B}$ ($\vec{H}$ and $\vec{B}$) is non-local in time)

The two effects turn out to be closely related.

• Let us first understand why imaginary parts of ϵ and μ lead to absorption of energy of EM fields by the medium.

Let us consider a monochromatic field:

$$\vec{E} = \vec{E} e^{i\omega t} +$$

Let us consider a monochromatic plane wave in a material:

$$\vec{E} = \frac{1}{E} e^{i\omega t + i\vec{k} \cdot \vec{x}}$$

suppose $\vec{k} = (k_x, 0, 0)$

then $k_x = \sqrt{\epsilon \mu} \omega \frac{1}{c}$

If ϵ or μ have imaginary part then k_x has imaginary part $\Rightarrow$ wave exponentially decays

From now on we will focus on ϵ , assuming μ real and constant. Then a general relation between D and E is

$$D(t) = \epsilon_0 E(t) + \epsilon_0 \int_0^{\infty} \chi(\tau) E(t-\tau) d\tau$$

Here we singled out the first term

for convenience, but most importantly, integrated is over positive $\tau \Rightarrow$ causality!

Let us now Fourier transform this expression:

$$D(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} D(\omega) e^{-i\omega t} d\omega$$

$$D(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} D(t) e^{i\omega t} dt$$

$$\begin{aligned} D(\omega) &= \epsilon_0 \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\omega t} E(t) dt + \\ &+ \epsilon_0 \int_0^{\infty} d\tau \int_{-\infty}^{\infty} \frac{dt}{\sqrt{2\pi}} e^{i\omega t} f(\tau) E(t-\tau) = \\ &= \epsilon_0 E(\omega) + \epsilon_0 E(\omega) \int_0^{\infty} d\tau f(\tau) e^{i\omega\tau} \equiv \\ &\equiv \epsilon(\omega) E(\omega) \end{aligned}$$

$$\frac{\epsilon(\omega)}{\epsilon_0} = 1 + \int_0^{\infty} d\tau f(\tau) e^{i\omega\tau} \quad (*)$$

$\epsilon(\omega)$ has real and imaginary parts:

$$\epsilon(\omega) = \epsilon'(\omega) + i\epsilon''(\omega)$$

Because $f(\tau)$ is real

$$\epsilon(-\omega) = \epsilon^*(\omega):$$

$$\epsilon'(-\omega) = \epsilon'(\omega) \text{ and } \epsilon''(-\omega) = -\epsilon''(\omega)$$

In dielectrics when $\omega \rightarrow 0$ we get the static situation (fields do not depend on time) ϵ becomes equal to real static susceptibility.

Causality and analytic properties of $\epsilon(\omega)$

- We will now consider ϵ as a function of complex variable ω , like we did for the Green's function of $\square$ operator.

- Because integral in (*) starts at $\tau=0$ $\epsilon(\omega)$ is an analytic function in the upper half plane: integral converges if $\text{Im } \omega > 0$ $e^{i\omega\tau} \sim e^{-(\text{Im } \omega) \cdot \tau}$.

We assume that the "memory" is finite so integral also converges for real ω (it is not true for conductors for $\omega=0$)

For very large real ω we can show that
$$\frac{\epsilon(\omega)}{\epsilon_0} = 1 - \frac{\text{const}}{\omega^2} + \dots$$

This is because at very high frequencies charged particles are basically free.


Then

$$m \ddot{r} = e E^1 e^{i\omega t} \Rightarrow$$

$$\Rightarrow r \sim \frac{1}{\omega^2}$$

$$\vec{P} = \sum e \vec{r} \approx \frac{1}{\omega^2} \Rightarrow \vec{D} - \epsilon_0 \vec{E} \sim \frac{1}{\omega^2}$$

$\epsilon(\omega)$



We will now use Cauchy's theorem
to relate $\epsilon'(\omega)$ and $\epsilon''(\omega)$ for
real ω (physical)

$$F(z) = \frac{1}{2\pi i} \oint_{\gamma} \frac{F(\omega')}{\omega' - z} d\omega'$$


take $F(z) = \frac{\epsilon(z)}{\epsilon_0} - 1$



$$\frac{\epsilon(z)}{\epsilon_0} - 1 = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\epsilon(x)/\epsilon_0 - 1}{x - z} dx$$

now we take

$$z = \omega + i\delta$$

↑ real ↑ infinitesimal

$$\frac{1}{x - \omega - i\delta} = P \left(\frac{1}{x - \omega} \right) + \pi i \delta(\omega' - \omega)$$

$$\frac{1}{2} \left(\frac{\epsilon(\omega)}{\epsilon_0} - 1 \right) = \frac{1}{2\pi i} P \int_{-\infty}^{\infty} \frac{\epsilon(x)/\epsilon_0 - 1}{x - \omega} d\omega'$$

Let's take real and imaginary parts:

$$\frac{\epsilon'(\omega)}{\epsilon_0} - 1 = \frac{1}{\pi} P \int_{-\infty}^{\infty} \frac{\epsilon''(\omega')}{x - \omega} dx$$

$$\frac{\epsilon''(\omega)}{\epsilon_0} = -\frac{1}{\pi} P \int \frac{\epsilon'(x) - 1}{x - \omega} dx$$

These are the Kramers-Kronig relations

Using symmetry properties we get

$$\frac{\epsilon'(\omega)}{\epsilon_0} = 1 + \frac{2}{\pi} P \int_0^{\infty} \frac{x \epsilon''(x)}{x^2 - \omega^2} dx$$

$$\frac{\epsilon''(\omega)}{\epsilon_0} = -\frac{2\omega}{\pi} P \int_0^{\infty} \frac{\epsilon'(\omega)/\epsilon_0 - 1}{x^2 - \omega^2} dx$$

Importance of these relations is in the fact that measuring absorption $\epsilon''(\omega)$ allows to reconstruct the entire dispersion of $\epsilon(\omega)$.

Let us consider an example

$$\frac{\epsilon}{\epsilon_0} = 1 + \frac{\gamma}{\omega_0^2 - \omega^2 - i\gamma\omega}$$

(resonant absorption on a line)

$$\frac{\epsilon}{\epsilon_0} = 1 + \frac{\text{const}}{\omega^2} \quad \text{at } \omega \rightarrow \infty \quad \checkmark$$

$$\frac{\epsilon'}{\epsilon_0} = 1 + \gamma \frac{\omega_0^2 - \omega^2}{(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2} \approx$$

$$\approx 1 + \gamma \frac{1}{\omega_0^2 - \omega^2}, \quad \text{when } \gamma \text{ is small,}$$

and $\omega \neq \omega_0$

$$\frac{\epsilon''}{\epsilon_0} = \frac{\gamma \omega}{(\omega_0^2 - \omega^2)^2 + \gamma^2 \omega^2}$$

Let's check KK: (for $\omega \neq \omega_0$ and γ -small)

$$\int_0^{\infty} \frac{dx}{x^2 - \omega^2} \frac{\gamma x}{(\omega_0^2 - x^2)^2 + \gamma^2 x^2} \approx$$

$$\approx \int_{\omega_0 - \gamma}^{\omega_0 + \gamma} \frac{1}{\omega_0^2 - \omega^2} \frac{\gamma x}{(\omega_0^2 - x^2)^2 + \gamma^2 x^2}$$

$$\int dz \frac{\gamma \gamma \omega_0}{(z \omega_0)^2 z^2 + \gamma^2 \omega_0^2} \approx \gamma$$

- Let's come back to plane waves. $\vec{E} = E \cdot e^{i\vec{k}\vec{r} - i\omega t}$

Since we derived all equations in frequency space, the relations between k and ω still hold:

$$i\omega \mu(\omega) \vec{H} = c \vec{\nabla} \times \vec{E}$$

$$i\omega \epsilon(\omega) \vec{E} = -c \vec{\nabla} \times \vec{H} \Rightarrow$$

$$k^2 = \epsilon(\omega) \mu \frac{\omega^2}{c^2}$$

Let us first assume that ϵ is real. Then group velocity of

light is given by $u = \frac{d\omega}{dk} = \frac{c}{d(n\omega)/d\omega}$

where $n(\omega) = \sqrt{\epsilon(\omega)\mu}$.

If ϵ is imaginary k develops imaginary part. Assume $\vec{k} \sim (k_x, 0, 0)$

$$k_x = \sqrt{\epsilon\mu} \frac{\omega}{c} \equiv [n(\omega) + i\alpha(\omega)] \frac{\omega}{c}$$

↑
extinction
coefficient.

KK-like relations can be also derived for n and α