




Particle Physics I

Lecture 5: Particle scattering, cross section

Prof. Radoslav Marchevski
October 2nd 2024

Today's learning targets


- Relation between decay width and cross section using Fermi's Golden rule
- Calculate cross section for $2 \rightarrow 2$ scattering in the center of mass and laboratory frames



Cross section

Cross section

- Consider a single particle of type a with velocity v_a , traversing a region of area A containing n_b particles of type b per unit volume



- In time δt a particle of type a traverses a region containing $\delta N = n_b(v_a + v_b)A\delta t$ particles of type b
- Interaction probability obtained from the effective cross-sectional area occupied by $n_b(v_a + v_b)A\delta t$ particles of type b :

$$\delta P = \frac{\delta N \sigma}{A} = \frac{n_b(v_a + v_b)A\delta t \sigma}{A} = n_b v \delta t \sigma$$

$$v = (v_a + v_b)$$

- $\Rightarrow$ rate per particle a is $r_a = \frac{dP}{dt} = n_b v \sigma$

Cross section

- Consider a volume V , where the total reaction rate, R , is

$$R = (n_b v \sigma) \cdot (n_a V) = (n_b V)(n_a v) \sigma = N_b \phi_a \sigma$$

- **i.e. reaction rate = flux $\times$ number of target particles $\times$ cross section**

Cross section definition


$$\sigma = \frac{\text{number of interactions per unit time per target}}{\text{incident flux}}$$

Incident flux = number of incident particles/unit area/unit time

- The “cross section”, σ , can be thought of as the effective cross-sectional area representing the size of the target object that the incoming particles must hit for the interaction to occur
- It is a measure of the probability of the interaction
- In general, this has nothing to do with the physical size of the target although there are exceptions, e.g. neutron absorption

Differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{\text{number of interactions per unit time per target into a solid angle } d\Omega}{\text{incident flux}}$$



$$\sigma = \int \frac{d\sigma}{d\Omega} d\Omega$$

Cross section calculation

- Consider scattering process $1 + 2 \rightarrow 3 + 4$
- Start from Fermi's Golden Rule:

$$\Gamma_{fi} = (2\pi)^4 \int |T_{fi}|^2 \delta(E_1 + E_2 - E_3 - E_4) \delta^3(\vec{p}_1 + \vec{p}_2 - \vec{p}_3 - \vec{p}_4) \frac{d^3\vec{p}_3}{(2\pi)^3} \frac{d^3\vec{p}_4}{(2\pi)^3} \quad (1)$$

- Here T_{fi} is the transition matrix for a normalization of 1 particle per unit volume
- Rate/Volume = (flux of 1) $\times$ (number density of 2) $\times \sigma = n_1(v_1 + v_2)n_2\sigma$
- For 1 target particle per unit volume, the rate is: $(v_1 + v_2)\sigma \Rightarrow \sigma = \frac{\Gamma_{fi}}{v_1 + v_2}$

$$\sigma = \frac{(2\pi)^4}{(v_1 + v_2)} \int |T_{fi}|^2 \delta(E_1 + E_2 - E_3 - E_4) \delta^3(\vec{p}_1 + \vec{p}_2 - \vec{p}_3 - \vec{p}_4) \frac{d^3\vec{p}_3}{(2\pi)^3} \frac{d^3\vec{p}_4}{(2\pi)^3} \quad (2)$$

These are not Lorentz-invariant terms

Cross section calculation

- To obtain Lorentz-invariant form we start by using wave functions normalised to $2E$ particles per unit volume: $\Psi' = \sqrt{2E} \Psi$

- Again define Lorentz-invariant matrix element $|M_{fi}| = \sqrt{2E_1 2E_2 2E_3 2E_4} |T_{fi}|$

$$\sigma = \frac{(2\pi)^{-2}}{2E_1 2E_2 (v_1 + v_2)} \int |M_{fi}|^2 \delta(E_1 + E_2 - E_3 - E_4) \delta^3(\vec{p}_1 + \vec{p}_2 - \vec{p}_3 - \vec{p}_4) \frac{d^3\vec{p}_3}{2E_3} \frac{d^3\vec{p}_4}{2E_4} \quad (3)$$

- The integral is now written in Lorentz-invariant form
- The quantity $F = 2E_1 2E_2 (v_1 + v_2)$ can be written in terms of a scalar product of 4-vectors and is also LI

$$F = 4 \sqrt{(p_1^\mu p_{2,\mu}) - m_1^2 m_2^2} \quad (4)$$

- $\Rightarrow$ the cross section is a Lorentz invariant quantity

Two special cases of Lorentz-invariant flux


1. Center-of-Mass frame (CoM):

$$\begin{aligned} F &= 4E_1 E_2 (v_1 + v_2) \\ &= 4E_1 E_2 \left(\frac{p^*}{E_1} + \frac{p^*}{E_2} \right) \\ &= 4p^* (E_2 + E_1) \\ &= 4p^* \sqrt{s} \end{aligned}$$

2. Target particle (particle 2) at rest:

$$\begin{aligned} F &= 4E_1 E_2 (v_1 + v_2) \\ &= 4E_1 m_2 v_1 \\ &= \frac{4E_1 m_2 |\vec{p}_1|}{E_1} \\ &= 4m_2 |\vec{p}_1| \end{aligned}$$

2 → 2 body scattering in CoM frame



- We will apply the Lorentz-invariant formula for the interaction cross section to the most common cases
- 2 → 2 body scattering in the CoM frame:

$$\sigma = \frac{(2\pi)^{-2}}{2E_1 2E_2 (v_1 + v_2)} \int |M_{fi}|^2 \delta(E_1 + E_2 - E_3 - E_4) \delta^3(\vec{p}_1 + \vec{p}_2 - \vec{p}_3 - \vec{p}_4) \frac{d^3\vec{p}_3}{2E_3} \frac{d^3\vec{p}_4}{2E_4} \quad (5)$$

- We can use $\vec{p}_1 + \vec{p}_2 = 0$ and $E_1 + E_2 = \sqrt{s}$

$$\sigma = \frac{(2\pi)^{-2}}{4|\vec{p}_i| \sqrt{s}} \int |M_{fi}|^2 \delta(\sqrt{s} - E_3 - E_4) \delta^3(\vec{p}_3 + \vec{p}_4) \frac{d^3\vec{p}_3}{2E_3} \frac{d^3\vec{p}_4}{2E_4} \quad (6)$$

2 → 2 body scattering in CoM frame

- The integral is exactly the same as in the particle decay calculation but with m_i replaced by $\sqrt{s}$
- 2 → 2 body scattering in the CoM frame:


$$\sigma = \frac{(2\pi)^{-2}}{4|\vec{p}_i|\sqrt{s}} \frac{|\vec{p}_f|}{4\sqrt{s}} \int |M_{fi}|^2 d\Omega^*$$

$$\sigma = \frac{1}{64\pi^2 s} \frac{|\vec{p}_f^*|}{|\vec{p}_i^*|} \int |M_{fi}|^2 d\Omega^* \quad (7)$$

$2 \rightarrow 2$ body scattering in CoM frame

- Elastic scattering: $|\vec{p}_i^*| = |\vec{p}_f^*|$

$$\sigma_{\text{elastic}} = \frac{1}{64\pi^2 s} \int |M_{fi}|^2 d\Omega^* \quad (8)$$



- For calculating the total Lorentz-invariant cross section, the result from the previous page is sufficient
 - not so useful for computing the differential cross section in a rest frame other than CoM
 - $d\Omega^* = d(\cos\theta^*)d\phi^*$ refers to the angles in CoM frame

$$d\sigma = \frac{1}{64\pi^2 s} \frac{|\vec{p}_f^*|}{|\vec{p}_i^*|} |M_{fi}|^2 d\Omega^* \quad (9)$$

- We would need to find a Lorentz-invariant expression for $d\sigma$

2 → 2 body scattering in CoM frame

- Express $d\Omega^*$ in terms of the Mandelstam t :

- $t = q^2 = (p_1 - p_3)^2 = m_1^2 + m_2^2 - 2p_1 \cdot p_3$

- In CoM frame:

- $p_1^{*\mu} = (E_1^*, 0, 0, |\vec{p}_1^*|), p_3^{*\mu} = (E_3^*, |\vec{p}_3^*|\sin\theta^*, 0, |\vec{p}_3^*|\cos\theta^*)$

- $p_1^{*\mu} \cdot p_3^{*\mu} = E_1^* E_3^* - |\vec{p}_1^*| |\vec{p}_3^*| \cos\theta^*$



- $t = m_1^2 + m_2^2 - 2E_1^* E_3^* + 2|\vec{p}_1^*| |\vec{p}_3^*| \cos\theta^*$

- $\Rightarrow dt = 2|\vec{p}_1^*| |\vec{p}_3^*| d(\cos\theta^*)$

- $d\Omega^* = d(\cos\theta^*) d\phi^* = \frac{dt d\phi^*}{2|\vec{p}_1^*| |\vec{p}_3^*|}$

$$d\sigma = \frac{1}{128\pi^2 s |\vec{p}_1^*|^2} |M_{fi}|^2 dt d\phi^*$$

(10)



2 → 2 body scattering in CoM frame

- Finally, integrating over $d\phi^*$ (assuming no ϕ^* dependence on $|M_{fi}|^2$)



$$\frac{d\sigma}{dt} = \frac{1}{64\pi s |\vec{p}_1^*|^2} |M_{fi}|^2 \quad (11)$$

- All quantities are Lorentz-invariant and therefore it applies to any rest frame
 - $|\vec{p}_1^*|$ is constant fixed by energy-momentum conservation
 - $|\vec{p}_1^*|^2 = \frac{1}{4s} [s - (m_1 + m_2)^2][s - (m_1 - m_2)^2]$
- Example of how to use $d\sigma/dt$: consider elastic scattering in the lab. frame where we can neglect the mass of the incoming particle (e.g. electron or neutrino scattering): $|\vec{p}_1^*|^2 = (s - m_2^2)^2 / (4s)$

$$\frac{d\sigma}{dt} = \frac{1}{16\pi (s - m_2^2)^2} |M_{fi}|^2 \quad (12)$$

2 → 2 body scattering in laboratory frame

- The other commonly occurring case is scattering from a fixed target in a laboratory frame (e.g. electron-proton scattering)
- Take the case of elastic scattering at high energy where the mass of the incoming particles can be neglected: $m_1 = m_3 = 0, m_2 = m_4 = M$



- Express the cross section in terms of the scattering angle of the e^- : $d\Omega = 2\pi d(\cos\theta)$

$$\frac{d\sigma}{d\Omega} = \frac{d\sigma}{dt} \frac{dt}{d\Omega} = \frac{1}{2\pi} \frac{dt}{d(\cos\theta)} \frac{d\sigma}{dt} \quad (13)$$

$2 \rightarrow 2$ body scattering in laboratory frame

- Four-momenta of the particles
 - $p_1 = (E_1, 0, 0, E_1)$
 - $p_2 = (M, 0, 0, 0)$
 - $p_3 = (E_3, E_3 \sin \theta, 0, E_3 \cos \theta)$
 - $p_4 = (E_4, \vec{p}_4)$
 - $\Rightarrow t = (p_1 - p_3)^2 = -2p_1 p_3 = -2E_1 E_3 (1 - \cos \theta)$
- From $(E, \vec{p})$ conservation $p_1 + p_2 = p_3 + p_4$ we can express t in terms of p_2 and p_4
 - $t = (p_2 - p_4)^2 = 2M^2 - 2ME_4 = -2M(M - E_4) = -2M(E_1 - E_3)$
- E_1 is constant (the energy of the incoming particle):

$$\frac{dt}{d(\cos \theta)} = 2M \frac{dE_3}{d(\cos \theta)} \quad (14)$$

2 → 2 body scattering in laboratory frame

- Equating the two expressions for t we get

- $E_3 = \frac{E_1 M}{M + E_1 - E_1 \cos \theta}$

- $\frac{dE_3}{d(\cos \theta)} = \frac{E_1^2 M}{(M + E_1 - E_1 \cos \theta)^2} = E_1^2 M \left(\frac{E_3}{E_1 M} \right)^2 = \frac{E_3^2}{M}$

$$\frac{d\sigma}{d\Omega} = \frac{1}{2\pi} \frac{dt}{d(\cos \theta)} \frac{d\sigma}{dt} = \frac{1}{2\pi} 2M \frac{E_3^2}{M} \frac{d\sigma}{dt} = \frac{E_3^2}{\pi} \frac{d\sigma}{dt} = \frac{E_3^2}{16\pi^2 (s - M^2)^2} |M_{fi}|^2 \quad (15)$$

- Using $s = (p_1 + p_2)^2 = M^2 + 2p_1 \cdot p_2 = M^2 + 2ME_1$, as $p_1^2 = 0$, gives $(s - M^2) = 2ME_1$

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \left(\frac{E_3}{ME_1} \right)^2 |M_{fi}|^2 \quad (\text{in the limit of } m_1 \rightarrow 0) \quad (16)$$

2 → 2 body scattering in laboratory frame

- Express E_3 as a function of θ : $E_3 = \frac{ME_1}{M+E_1(1-\cos\theta)}$

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \left(\frac{1}{M + E_1(1 - \cos\theta)} \right)^2 |M_{fi}|^2 \quad (17)$$

- General form of 2 → 2 body scattering in the lab frame in case the mass m_1 can't be neglected

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \frac{1}{|\vec{p}_1| m_1} \frac{|\vec{p}_3|^2}{|\vec{p}_3|(E_1 + m_2) - E_3 |\vec{p}_1| \cos\theta} |M_{fi}|^2 \quad (18)$$

- There is only one independent variable: the angle θ , from conservation of energy:

$$(E_1 + m_2) = \sqrt{|\vec{p}_3|^2 + m_3^2} + \sqrt{|\vec{p}_1|^2 + |\vec{p}_3|^2 - 2|\vec{p}_1||\vec{p}_3|\cos\theta + m_4^2}$$

Summary

- We used a Lorentz-invariant formulation of Fermi's golden rule to derive decay rates and cross sections
 - Expressed in the Lorentz-invariant Matrix Element (wave-functions normalised to $2E/\text{unit volume}$)
- **Particle decay width:**

$$\Gamma = \frac{|\vec{p}^*|}{32\pi^2 m_i^2} \int |M_{fi}|^2 d\Omega \quad (19)$$

$$p^* = \frac{1}{2m_i} \sqrt{[m_i^2 - (m_1 + m_2)^2][m_i^2 - (m_1 - m_2)^2]} \text{(function of the mass of the particles)}$$

- **Scattering cross section in CoM frame:**

$$\sigma = \frac{1}{64\pi^2 s} \frac{|\vec{p}_f^*|}{|\vec{p}_i^*|} \int |M_{fi}|^2 d\Omega^* \quad (20)$$

Summary

- Invariant differential cross section valid in all frames:

$$\frac{d\sigma}{dt} = \frac{1}{64\pi s |\vec{p}_i^*|^2} \int |M_{fi}|^2 \quad (21)$$

$$|\vec{p}_i^*|^2 = \frac{1}{4s} [s - (m_1 + m_2)^2][s - (m_1 - m_2)^2]$$

- Differential cross section in the lab. frame ($m_1 = 0$):

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \left(\frac{E_3}{ME_1} \right)^2 |M_{fi}|^2 \Leftrightarrow \frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \left(\frac{1}{M + E_1(1 - \cos\theta)} \right)^2 |M_{fi}|^2 \quad (22)$$

- Differential cross section in the lab. frame ($m_1 \neq 0$):

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \frac{1}{|\vec{p}_1| m_1} \frac{|\vec{p}_3|^2}{|\vec{p}_3|(E_1 + m_2) - E_3 |\vec{p}_1| \cos\theta} |M_{fi}|^2 \quad (23)$$

$$(E_1 + m_2) = \sqrt{|\vec{p}_3|^2 + m_3^2} + \sqrt{|\vec{p}_1|^2 + |\vec{p}_3|^2 - 2|\vec{p}_1||\vec{p}_3|\cos\theta + m_4^2}$$

Examples: relativistic Rutherford cross section

- Need to get a differential cross section in the lab frame with a massless incident particle $M \gg E_1$ (Eq. 22)

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2 M^2} |M_{fi}|^2 \quad (24)$$

- using

$$|M_{fi}|^2 = (2E_1 \cdot 2M \cdot 2E_3 \cdot 2M) |T_{fi}|^2 \quad (25)$$

$$\Rightarrow \frac{d\sigma}{d\Omega} = \frac{16M^2 E_1 E_3}{64\pi^2 M^2} |T_{fi}|^2 = \frac{E^2}{(2\pi)^2} |T_{fi}|^2 \quad (26)$$

- Hamiltonian for a Coulomb potential: $\hat{H} = e\phi(x)$
- Initial and final state wave function are plane waves: $|\Psi_i\rangle = e^{-ip_1 \cdot x}, |\Psi_f\rangle = e^{-ip_3 \cdot x}$
- Matrix element T_{fi} :

$$T_{fi} = \langle \Psi_f | \hat{H} | \Psi_i \rangle = \int e^{ip_3 \cdot x} e\phi(x) e^{-ip_1 \cdot x} d^3x \quad (27)$$

Examples: relativistic Rutherford cross section

- Define momentum transfer as $q = p_3 - p_1$

$$T_{fi} = \langle \Psi_f | \hat{H} | \Psi_i \rangle = e \int \phi(x) e^{iq \cdot x} d^3x \quad (28)$$

- use

$$e^{iq \cdot x} = -\frac{1}{|q|^2} \cdot \nabla^2 e^{-iq \cdot x} \text{ and } \int (u \nabla^2 v - v \nabla^2 u) d^3x = 0 \text{ (Green's theorem)} \quad (29)$$

$$T_{fi} = -\frac{e}{|q|^2} \int (\nabla^2 \phi(x)) e^{-iq \cdot x} d^3x \quad (30)$$

- From Poisson equation: $\nabla^2 \phi(x) = -\rho(x)$, where $\rho(x) = Ze f(x)$ is a static potential with a normalisation condition $\int f(x) d^3x = 1$

$$T_{fi} = \frac{4\pi\alpha Z}{|q|^2} \int f(x) e^{-iq \cdot x} d^3x \quad (31)$$

Examples: relativistic Rutherford cross section

- Definition: $F(q) = \int f(x) e^{-iq \cdot x} d^3x$ is the Fourier transformation of the charge function $f(x)$, called also a **Form Factor (FF)** of the charge distribution
- The FF contains all the information about the spatial distribution of the charge of the studied object
- In our case we replace it with a delta function $F(q) = 1$
- This gives for the matrix element simply:


$$T_{fi} = \frac{4\pi\alpha Z}{|q|^2} \quad (32)$$

- So the Rutherford cross section becomes:

$$\frac{d\sigma}{d\Omega} = \frac{E^2}{(2\pi)^2} \left| \frac{4\pi\alpha Z}{|q|^2} \right|^2 = \frac{4Z^2\alpha^2 E^2}{|q|^4} \quad (33)$$

Examples: relativistic Rutherford cross section

- **Quantum Field theory:** the electron interacts with a nucleus (charge = Ze) via the exchange of a photon
 - photon momentum: $q = p - p'$
 - de Broglie wavelength: $\lambda = 1/|q|$
- If λ is large, the internal structure of the nucleus can not be resolved and can be considered as a point-like object (that is what we assumed so far in our calculations)



Particle accelerators: motivations

- Accelerators serve as “microscopes”
 - $|q| = 1 \text{ GeV} \Rightarrow \lambda \approx 1.2 \times 10^{-15} \text{ m}$ – size of a proton
 - $|q| = 10^3 \text{ GeV} \Rightarrow \lambda \approx 1.2 \times 10^{-18} \text{ m}$ – size of a proton substructure (e.g. quarks)
 - $\Rightarrow$ accelerators allow us to look for the substructure of particles
- Types of operation modes:
 - **fixed target** of mass m , beam with energy E : $\sqrt{s} \approx \sqrt{2mE}$
 - **collider** with two beams of energy E : $\sqrt{s} = 2E$
- Example: collider with two 22 GeV (1 TeV) beams gives the same center-of-mass energy as a fixed target with a beam of 1 TeV (10^3 TeV)
- When looking for massive particles produced in the interactions, aim for the highest energy possible and the collider mode is the more appropriate one


Particle accelerators: luminosity

- Luminosity ($\mathcal{L}$) is the exposure of the target (beam) to scattering (collision) per unit time and unit area
- **Fixed target experiments**
 - $\mathcal{L} = \text{flux} \times \text{number of scattering centers} = \Phi_a \times N_b = n_a \times v_a \times N_b$
- **Colliding beams**
 - $\mathcal{L} = \frac{n_a \times n_b}{A} b f$, where $A = 4\pi\sigma_x\sigma_y$
 - here n_i is the number of particles per bunch, b is the number of bunches, f is the frequency of the orbit
 - typical beam size of the LHC $\sigma_x \approx \sigma_y \approx 15\mu\text{m}$
- The number of produced events for a process with cross section σ is:


$$N = \mathcal{L}_{\text{int}} \times \sigma, \quad \mathcal{L}_{\text{int}} = \int \mathcal{L} dt$$

LHC: instantaneous luminosity $\mathcal{L}$

Peak instantaneous luminosity :




LHC: integrated luminosity $\mathcal{L}_{\text{int}}$



LHC: example production cross section

- $N_{pp} = \mathcal{L}_{\text{int}} \sigma_p^{\text{inel}} = 20 \cdot 100 \cdot 10^{12}$ (with $\sigma_p^{\text{inel}} = 100 \text{ mb}$) cross section at 13.6 TeV
 - W boson (leptonic decays): 63000 pb
 - Z boson: $Z \rightarrow \mu\mu$: 2103 pb
 - $t\bar{t}$: 920 pb
 - gluon fusion Higgs production: 53 pb
 - $t\bar{t}H$: 600 fb
 - HH : 34 fb
- How many given interactions did LHC produce in 2022?
 - assume $\mathcal{L} = 20 \text{ fb}^{-1}$?




Summary of Lecture 5

Main learning outcomes

- How to deal with kinematics of cross sections
- The fundamental particle physics is in the matrix element
- The above equations are the basis for all calculations that follow

Additional slides: Dirac δ –function

- In the relativistic formulation of decay rates and cross sections we will make use of the Dirac δ function: “infinitely narrow spike of unit area”



$$\int_{-\infty}^{\infty} \delta(x - a) dx = 1$$

$$\int_{-\infty}^{\infty} f(x) \delta(x - a) dx = f(a)$$

- Any function with the above properties can represent $\delta(x)$, e.g.:

$$\delta(x) = \lim_{\sigma \rightarrow 0} \frac{1}{\sqrt{2\pi}\sigma} e^{-\left(\frac{x^2}{2\sigma^2}\right)}$$

Infinitesimally narrow Gaussian


Additional slides: Dirac δ –function of a function

- An expression for the δ –function of a function $\delta(f(x))$:
 - start from the definition of a δ –function:

$$\int_{y_1}^{y_2} \delta(y) dy = \begin{cases} 1 & \text{if } y_1 < 0 < y_2 \\ 0 & \text{otherwise} \end{cases}$$

- Now express in terms of $y = f(x)$, where $f(x_0) = 0$ and change variables

$$\int_{x_1}^{x_2} \delta(f(x)) \frac{df}{dx} dx = \begin{cases} 1 & \text{if } x_1 < 0 < x_2 \\ 0 & \text{otherwise} \end{cases}$$



Additional slides: Dirac δ –function of a function

- From the properties of a δ –function (i.e. only non-zero at x_0)

$$\left| \frac{df}{dx} \right| \int_{x_1}^{x_2} \delta(f(x)) dx = \begin{cases} 1 & \text{if } x_1 < 0 < x_2 \\ 0 & \text{otherwise} \end{cases}$$

- Rearranging and expressing RHS as a δ –function

$$\int_{x_1}^{x_2} \delta(f(x)) dx = \frac{1}{\left| \frac{df}{dx} \right|_{x_0}} \int_{x_1}^{x_2} \delta(x - x_0) dx \Rightarrow \delta(f(x)) = \left| \frac{df}{dx} \right|_{x_0}^{-1} \delta(x - x_0)$$