
QUANTUM PHYSICS III

Problem Set 11

26 November 2024

1. Properties of spherical Bessel functions

Recall the definitions of the spherical Bessel and von Neumann functions (for $l \in \mathbb{N}$) :

$$j_l(\rho) = (-1)^l \rho^l \left(\frac{1}{\rho} \frac{d}{d\rho} \right)^l \frac{\sin \rho}{\rho}, \quad (1)$$

$$y_l(\rho) = -(-1)^l \rho^l \left(\frac{1}{\rho} \frac{d}{d\rho} \right)^l \frac{\cos \rho}{\rho}. \quad (2)$$

1. Compute the asymptotic behavior of j_l and y_l for $\rho \rightarrow 0$:

$$j_l(\rho) \sim \frac{\rho^l}{(2l+1)!!}, \quad (3)$$

$$y_l(\rho) \sim \frac{(2l-1)!!}{\rho^{l+1}}, \quad (4)$$

”!!” is the double factorial, defined as $n!! = n \cdot (n-2) \cdot (n-4) \cdot \dots$, with the usual convention $0!! = 1$.

2. Compute the asymptotic behavior of j_l and y_l for $\rho \rightarrow \infty$:

$$j_l(\rho) \sim \frac{1}{\rho} \sin\left(\rho - l\frac{\pi}{2}\right), \quad (5)$$

$$y_l(\rho) \sim \frac{1}{\rho} \cos\left(\rho - l\frac{\pi}{2}\right). \quad (6)$$

2. Scattering phase shift in a Yukawa potential

For a Yukawa potential,

$$V(r) = \frac{V_0}{\mu r} e^{-\mu r} \quad (7)$$

it can be shown that in the first Born approximation, the scattering amplitude is given by

$$f(\mathbf{p}' \leftarrow \mathbf{p}) = -\frac{2mV_0}{\mu} \frac{1}{2p^2(1 - \cos \theta) + \mu^2} \quad (8)$$

θ is the angle between $\mathbf{p}$ and $\mathbf{p}'$, and $p = |\mathbf{p}| = |\mathbf{p}'|$; (we also consider $\hbar = 1$ as usual).

1. Obtain an expression for the scattering phase shift δ_l in terms of Legendre functions of the second kind :

$$Q_l(\zeta) = \frac{1}{2} \int_{-1}^1 \frac{P_l(\zeta')}{\zeta - \zeta'} d\zeta' \quad (9)$$

We will assume that $|\delta_l| \ll 1$.

Hint. Make use of the orthogonality of the Legendre polynomials :

$$\int_{-1}^1 d\zeta P_l(\zeta) P_{l'}(\zeta) = \frac{2\delta_{ll'}}{2l+1} \quad (10)$$

2. Use the expansion formula :

$$Q_l(\zeta) = \sum_{n=0}^{\infty} \frac{(l+2n)!}{(2l+2n+1)!!(2n)!!} \cdot \frac{1}{\zeta^{l+2n+1}} \quad (11)$$

to show the following :

- (a) δ_l is negative, resp. positive, when the potential is repulsive, resp. attractive.
- (b) When the de Broglie wavelength is much longer than the range of the potential, δ_l is proportional to p^{2l+1} . Find the proportionality constant.

3. Scattering off a spherical potential

Consider the potential $V(r)$ given by

$$V(r) = \begin{cases} 0 & \text{for } r > R, \\ \infty & \text{for } r < R. \end{cases} \quad (12)$$

1. Derive an expression for the s -wave ($l = 0$) phase shift. By obtaining a general expression for the scattering phase shift δ_l in the limit $kR \ll 1$, justify contributions beyond s -wave can be neglected.
2. What is the total cross section $\sigma = \int (\frac{d\sigma}{d\Omega}) d\Omega$ in the extreme low-energy limit $k \rightarrow 0$? Compare your answer with the geometric cross section πa^2 . Use the following relations :

$$\frac{d\sigma}{d\Omega} = |f(\theta)|^2, \quad (13)$$

$$f(\theta) = \left(\frac{1}{k}\right) \sum_{l=0}^{\infty} (2l+1) e^{i\delta_l} \sin \delta_l P_l(\cos \theta). \quad (14)$$

4. Scattering off a constant potential

Consider a potential

$$V = \begin{cases} 0 & \text{for } r > R, \\ V_0 = \text{constant} & \text{for } r < R, \end{cases} \quad (15)$$

where V_0 may be positive or negative.

1. Using the method of partial waves, show that for $|V_0| \ll E = \frac{\hbar^2 k^2}{2m}$ and $kR \ll 1$, the scattering phase shift can be written as :

$$\tan \delta_l = \frac{(kR)^{2l+3}}{(2l+3)!!(2l+1)!!} \left[\frac{\kappa^2}{k^2} - 1 \right], \quad (16)$$

where $E - V_0 \equiv \frac{\hbar^2 k^2}{2m}$.

Furthermore, show that the differential cross section is isotropic and that the total cross section is given by

$$\sigma_{\text{tot}} = \left(\frac{16\pi}{9} \right) \left(\frac{m^2 V_0^2 R^6}{\hbar^4} \right). \quad (17)$$

Hint : the following identity may be useful :

$$f'_l(x) = \frac{l}{x} f_l(x) - f_{l+1}(x), \quad (18)$$

for f any spherical Bessel or von Neumann function.

2. Suppose the energy is raised slightly. Show that the angular distribution can then be written as

$$\frac{d\sigma}{d\Omega} = A + B \cos \theta. \quad (19)$$

Obtain an approximate expression for B/A .

5. Scattering off a $\frac{1}{r^2}$ potential

Let ($A > 0$) and consider the potential

$$V(r) = \frac{\hbar^2 A}{2mr^2}. \quad (20)$$

1. Write down the eigenvalue equation of the Hamiltonian associated with this potential, and find a solution in terms of spherical Bessel functions.

Hint : Even though the spherical Bessel functions $j_l(x)$ were defined for an integer l , they can also be extended to any complex l .

2. Obtain the phase shifts exactly. Show that for $A \ll 1$, one approximately has

$$\delta_l = -\frac{\pi}{2} \frac{A}{2l+1}. \quad (21)$$

What is the value of the total cross section ?