


Radiation Biology, Protection and Applications (FS2024)

Week 7: Radiation Sources



Oskari Pakari, PhD
Scientist, LRS

<https://www.epfl.ch/labs/lrs/>

Oskari.pakari@epfl.ch

Lecture Plan Nov/Dec

01.11.2024	Radiation sources	Oskari
08.11.2024	Radiation shielding	Vincent
15.11.2024	Introduction to Safeguards & the IAEA	Uwe, BFE
22.11.2024	Industrial applications of radiation: Gauges, Tracers, Batteries, Food	Oskari
29.11.2024	Emergency dosimetry	Oskari
06.12.2024	Advances in luminescence dosimetry	Lily, PSI
13.12.2024	Safeguards and Export control	Uwe, BFE
After 15.12: Online	Exam example	Oskari

- Basics of radioactivity: See Lecture 1
- Radiation Concepts
- Fast Electron Sources
 - Beta decay
 - Internal conversion
- Heavy Charged Particle Sources
 - Alpha decay
 - Spontaneous fission
- Electromagnetic Radiation (EMR) Sources
 - Gamma rays
 - X-rays, characteristic X-rays
- Neutron Sources
 - Spontaneous fission
 - Neutrons from (α, n) -reactions
 - Photoneutrons
 - Accelerated charged particles

- There are four general types of radiation generated in **nuclear** and **atomic** processes:
 - **Charged particulate radiation:**
 - Fast electrons: β^+ and β^- from nuclear decay, energetic electrons.
 - Heavy charged particles: all energetic ions with $A \geq 1$ (p^+ , α^{2+} , fission products, nuclear reaction products)
 - **Uncharged radiation:**
 - Electromagnetic radiation: photons, X-rays (from electron transitions between atomic shells), γ -rays (from nuclear transitions)
 - Neutrons: slow and fast (generated in nuclear reactions.)
- **Absolute activity** is defined as rate of decay: It **measures** the **source disintegration rate**, **not** the **emission rate** of radiation.

❑ Energy range of **ionizing** radiation:


- 10 eV: minimum energy for ionization of typical materials.
- to 20 MeV: upper bound for practical applications.

❑ Hard radiation:

- High penetrating power.
- Sources are less affected by self-absorption.
- γ -rays, hard X-rays or neutrons.

❑ Soft radiation:

- Highly ionizing
- Low penetrating power
- Sources must be thin to minimize self-absorption.
- Charged particles, soft X-rays.



□ Radioactive decay in which a beta particle (electron or positron) is emitted.

- electron emission: "beta minus" (β^-),

$$n^0 \rightarrow p^+ + e^- + \bar{\nu}_e$$

$${}_Z^AX \rightarrow {}_{Z+1}^AY + \beta^- + \bar{\nu}_e$$


- positron emission: "beta plus" (β^+).

$$\text{Energy} + p^+ \rightarrow n^0 + e^+ + \nu_e$$


$${}_Z^AX \rightarrow {}_{Z-1}^AY + \beta^+ + \nu_e$$

□ Beta plus decay cannot occur in isolation:

- Neutron mass $m_n > m_p$. (proton is stable)
- Requires available excitation energy from inside nucleus
- The Q value goes into:
 - the process of converting a proton into a neutron,
 - the positron and the neutrino, and into
 - the kinetic energy of these particles.




The Feynman diagram for beta decay of a neutron into a proton, electron, and electron-antineutrino via an intermediate heavy W-boson.




The decay scheme of ${}^{36}\text{Cl}$ and the resulting beta particle energy distribution.

Beta Decay of Th-234

Neutron becomes a proton



${}_{90}^{234}\text{Th}$




- (Artificial) beta emitters can be produced by neutron irradiation of stable materials in nuclear reactors or high neutron flux facilities.


- As most beta decays populate an excited state of the daughter nucleus, they are not “pure”, i.e., they are accompanied by γ -rays.

- Examples for beta decays:

- Beta minus:



- Beta plus:



- Electron capture:






Table 1.1 Some “Pure” Beta-Minus Sources

Nuclide	Half-Life	Endpoint Energy (MeV)
${}^3\text{H}$	12.26 y	0.0186
${}^{14}\text{C}$	5730 y	0.156
${}^{32}\text{P}$	14.28 d	1.710
${}^{33}\text{P}$	24.4 d	0.248
${}^{35}\text{S}$	87.9 d	0.167
${}^{36}\text{Cl}$	3.08×10^5 y	0.714
${}^{45}\text{Ca}$	165 d	0.252
${}^{63}\text{Ni}$	92 y	0.067
${}^{90}\text{Sr}/{}^{90}\text{Y}$	27.7 y/64 h	0.546/2.27
${}^{99}\text{Tc}$	2.12×10^5 y	0.292
${}^{147}\text{Pm}$	2.62 y	0.224
${}^{204}\text{Tl}$	3.81 y	0.766

Data from Lederer and Shirley.¹

The curious case of ^{64}Cu

^{63}Zn β^+	^{64}Zn $2\beta^+$	^{65}Zn β^+	^{66}Zn Stable
^{62}Cu β^+	^{63}Cu Stable	^{64}Cu β^+	^{65}Cu Stable
^{61}Ni Stable	^{62}Ni Stable	^{63}Ni β^-	^{64}Ni Stable




EC = Electron capture

□ Source of nearly **monoenergetic** electrons:

$$E_{e^-} = E_{ex} - E_b$$

□ Process:

- Alternative to de-excitation of an excited nuclear state by emission of a γ -ray.
- Nuclear excitation energy E_{ex} is transferred to an orbital electron.
- Discrete energies represent transitions between atomic energy levels (shells).
- A single excited atom can lead to several groups of electrons with different energies.
- Sometimes sources have superimposed the β -spectrum of the parent nucleus.




Some Common Conversion Electron Sources					
Parent Nuclide	Parent Half-Life	Decay Mode	Decay Product	Transition Energy of Decay Product (keV)	Conversion Electron Energy (keV)
^{109}Cd	453 d	EC	$^{109\text{m}}\text{Ag}$	88	62 84
^{113}Sn	115 d	EC	$^{113\text{m}}\text{In}$	393	365 389
^{137}Cs	30.2 y	β^-	$^{137\text{m}}\text{Ba}$	662	624 656
^{139}Ce	137 d	EC	$^{139\text{m}}\text{La}$	166	126 159
^{207}Bi	38 y	EC	$^{207\text{m}}\text{Pb}$	$\left\{ \begin{array}{l} 570 \\ 1064 \end{array} \right.$	482 554 976 1048

Data from Lederer and Shirley.¹

Conversion electrons are the **only practical laboratory-size source** of monoenergetic electron groups **in the high keV to MeV** energy range.

- Decay by emission of an alpha particle (or ${}^4\text{He}^{2+}$ nucleus):



- Alpha decay can be described in the framework of Quantum Mechanics: 'tunneling' through a potential barrier
- Probability of emission increases with the energy of the alpha particle E_α ($\sim e^{-G}$, G =Gamow-Factor).





Figure 1.3 Alpha particle groups produced in the decay of ${}^{238}\text{Pu}$. The pulse height spectrum shows the three groups as measured by a silicon surface barrier detector. Each peak is identified by its energy in MeV and percent abundance (in parentheses). The insert shows the decay scheme, with energy levels in the product nucleus labeled in MeV. (Spectrum from Chanda and Deal.)



- Each α particle shares the energy with the recoil nucleus in a unique way ($Q=Q$ -value of the decay):



$$E_{\alpha} = Q (A - 4)/A$$

- Alpha particles appear in one or more (essentially) monoenergetic energy groups.
- Typical kinetic energy $E_{\alpha} \sim 5$ MeV with a speed of 15,000 km/s.
- Alpha particles are among the most hazardous forms of internal radiation:**
 - Energy loss takes place within a very short distance.
 - Significant damage to surrounding biomolecules.
- External alpha irradiation is not harmful:
 - Completely absorbed by a very thin (μm) dead layer of skin as well as by a few centimeters of air.


Common Alpha-Emitting Radioisotope Sources				
Source	Half-Life	Alpha Particle Kinetic Energy (with Uncertainty) in MeV		Percent Branching
^{148}Gd	93 y	3.182787	± 0.000024	100
^{232}Th	1.4×10^{10} y	4.012	± 0.005	77
		3.953	± 0.008	23
^{238}U	4.5×10^9 y	4.196	± 0.004	77
		4.149	± 0.005	23
^{235}U	7.1×10^8 y	4.598	± 0.002	4.6
		4.401	± 0.002	56
		4.374	± 0.002	6
		4.365	± 0.002	12
^{236}U	2.4×10^7 y	4.219	± 0.002	6
		4.494	± 0.003	74
		4.445	± 0.005	26
		4.6875	± 0.0015	76.3
^{230}Th	7.7×10^4 y	4.6210	± 0.0015	23.4
		4.7739	± 0.0009	72
^{234}U	2.5×10^5 y	4.7220	± 0.0009	28
		5.0590	± 0.0008	11
^{231}Pa	3.2×10^4 y	5.0297	± 0.0008	20
		5.0141	± 0.0008	25.4
		4.9517	± 0.0008	22.8
		5.1554	± 0.0007	73.3
^{239}Pu	2.4×10^4 y	5.1429	± 0.0008	15.1
		5.1046	± 0.0008	11.5
		5.16830	± 0.00015	76
^{240}Pu	6.5×10^3 y	5.12382	± 0.00023	24
		5.2754	± 0.0010	87.4
^{243}Am	7.4×10^3 y	5.2335	± 0.0010	11
		5.30451	± 0.00007	99+
^{210}Po	138 d	5.48574	± 0.00012	85.2
^{241}Am	433 y	5.44298	± 0.00013	12.8
^{238}Pu	88 y	5.49921	± 0.00020	71.1
		5.4565	± 0.0004	28.7
^{244}Cm	18 y	5.80496	± 0.00005	76.4
		5.762835	± 0.000030	23.6
^{243}Cm	30 y	6.067	± 0.003	1.5
		5.992	± 0.002	5.7
		5.7847	± 0.0009	73.2
		5.7415	± 0.0009	11.5
^{242}Cm	163 d	6.11292	± 0.00008	74
		6.06963	± 0.00012	26
^{254m}Es	276 d	6.4288	± 0.0015	93
^{253}Es	20.5 d	6.63273	± 0.00005	90
		6.5916	± 0.0002	6.6


Data from Rytz.³

Alpha Decay of U-238


 $^{238}_{92}\text{U}$ 

Cloud chamber: U sample






**How can we get even
heavier charged
particles?**




Reminder:
Chart of nuclides



Primary Decay Mode


Stable	n	e- capture
Fission	2n	p
α	β^+	2p
β^-	$2\beta^+$	3p
$2\beta^-$	e+	
Long-lived		
Estimated		
Unknown		

Primary Decay Mode



Oxygen Z=8	11O 2p	12O 2p	13O β+	14O β+	15O β+	16O Stable	17O Stable	18O Stable	19O β-
Nitrogen Z=7	10N p	11N p	12N β+	13N β+	14N Stable	15N Stable	16N β-	17N β-	18N β-
Carbon Z=6	8C 2p	9C β+	10C β+	11C β+	12C Stable	13C Stable	14C β-	15C β-	16C β-
Boron Z=5	6B 2p	7B p	8B β+	9B p	10B Stable	11B Stable	12B β-	13B β-	14B β-
Beryllium Z=4	5Be p	6Be 2p	7Be e- capture	8Be α	9Be Stable	10Be β-	11Be β-	12Be β-	13Be n
Lithium Z=3	3Li p	4Li p	5Li p	6Li Stable	7Li Stable	8Li β-	9Li β-	10Li n	11Li β-
Helium Z=2	3He Stable	4He Stable	5He n	6He β-	7He n	8He β-	9He n	10He 2n	
Hydrogen Z=1	1H Stable	2H Stable	3H β-	4H n	5H 2n	6H n	7H 2n		
Neutron Z=0	1n β-								

Primary Decay Mode



idium Z=46	90Pd β^+	91Pd β^+	92Pd β^+	93Pd β^+	94Pd β^+	95Pd β^+	96Pd β^+	97Pd β^+	98Pd β^+	99Pd β^+	100Pd e- capture	101Pd β^+	102Pd 2 β^+	103Pd e- capture	104Pd Stable	105Pd Stable	106Pd Stable
88Rh β^+	89Rh β^+	90Rh β^+	91Rh β^+	92Rh β^+	93Rh β^+	94Rh β^+	95Rh β^+	96Rh β^+	97Rh β^+	98Rh β^+	99Rh β^+	100Rh β^+	101Rh e- capture	102Rh β^+	103Rh Stable	104Rh β^-	105Rh β^-
87Ru β^+	88Ru β^+	89Ru β^+	90Ru β^+	91Ru β^+	92Ru β^+	93Ru β^+	94Ru β^+	95Ru β^+	96Ru 2 β^+	97Ru β^+	98Ru Stable	99Ru Stable	100Ru Stable	101Ru Stable	102Ru Stable	103Ru β^-	104Ru 2 β^-
86Tc β^+	87Tc β^+	88Tc β^+	89Tc β^+	90Tc β^+	91Tc β^+	92Tc β^+	93Tc β^+	94Tc β^+	95Tc β^+	96Tc β^+	97Tc e- capture	98Tc β^-	99Tc β^-	100Tc β^-	101Tc β^-	102Tc β^-	103Tc β^-
85Mo β^+	86Mo β^+	87Mo β^+	88Mo β^+	89Mo β^+	90Mo β^+	91Mo β^+	92Mo 2 β^+	93Mo e- capture	94Mo Stable	95Mo Stable	96Mo Stable	97Mo Stable	98Mo 2 β^-	99Mo β^-	100Mo 2 β^-	101Mo β^-	102Mo β^-
84Nb β^+	85Nb β^+	86Nb β^+	87Nb β^+	88Nb β^+	89Nb β^+	90Nb β^+	91Nb e- capture	92Nb β^+	93Nb Stable	94Nb β^-	95Nb β^-	96Nb β^-	97Nb β^-	98Nb β^-	99Nb β^-	100Nb β^-	101Nb β^-
83Zr β^+	84Zr β^+	85Zr β^+	86Zr β^+	87Zr β^+	88Zr e- capture	89Zr β^+	90Zr Stable	91Zr Stable	92Zr Stable	93Zr β^-	94Zr 2 β^-	95Zr β^-	96Zr 2 β^-	97Zr β^-	98Zr β^-	99Zr β^-	100Zr β^-
82Y β^+	83Y β^+	84Y β^+	85Y β^+	86Y β^+	87Y β^+	88Y β^+	89Y Stable	90Y β^-	91Y β^-	92Y β^-	93Y β^-	94Y β^-	95Y β^-	96Y β^-	97Y β^-	98Y β^-	99Y β^-
81Sr β^+	82Sr e- capture	83Sr β^+	84Sr 2 β^+	85Sr e- capture	86Sr Stable	87Sr Stable	88Sr Stable	89Sr β^-	90Sr β^-	91Sr β^-	92Sr β^-	93Sr β^-	94Sr β^-	95Sr β^-	96Sr β^-	97Sr β^-	98Sr β^-
80Rb β^+	81Rb β^+	82Rb β^+	83Rb e- capture	84Rb β^+	85Rb Stable	86Rb β^-	87Rb β^-	88Rb β^-	89Rb β^-	90Rb β^-	91Rb β^-	92Rb β^-	93Rb β^-	94Rb β^-	95Rb β^-	96Rb β^-	97Rb β^-
79Kr β^+	80Kr Stable	81Kr e- capture	82Kr Stable	83Kr Stable	84Kr Stable	85Kr β^-	86Kr 2 β^-	87Kr β^-	88Kr β^-	89Kr β^-	90Kr β^-	91Kr β^-	92Kr β^-	93Kr β^-	94Kr β^-	95Kr β^-	96Kr β^-
78Br β^+	79Br Stable	80Br β^-	81Br Stable	82Br β^-	83Br β^-	84Br β^-	85Br β^-	86Br β^-	87Br β^-	88Br β^-	89Br β^-	90Br β^-	91Br β^-	92Br β^-	93Br β^-	94Br β^-	95Br β^-
77Se Stable	78Se Stable	79Se β^-	80Se 2 β^-	81Se β^-	82Se 2 β^-	83Se β^-	84Se β^-	85Se β^-	86Se β^-	87Se β^-	88Se β^-	89Se β^-	90Se β^-	91Se β^-	92Se β^-	93Se β^-	94Se β^-


																		Mendelevium Z=101																		Fermium Z=100																		Einsteinium Z=99																		Californium Z=98																		Berkelium Z=97																		Curium Z=96																		Americium Z=95																		Plutonium Z=94																		Neptunium Z=93																		Uranium Z=92																		Protactinium Z=91																		Thorium Z=90																		Actinium Z=89																		Radium Z=88																																																																																																																																																																																																																																									
																		244Md245Md246Md247Md																		241Fm242Fm243Fm244Fm245Fm246Fm																		239Es240Es241Es242Es243Es244Es245Es																		237Cf238Cf239Cf240Cf241Cf242Cf243Cf244Cf																		233Bk234Bk235Bk236Bk237Bk238Bk239Bk240Bk241Bk242Bk243Bk																		231Cm232Cm233Cm234Cm235Cm236Cm237Cm238Cm239Cm240Cm241Cm242Cm																		227Am228Am229Am230Am231Am232Am233Am234Am235Am236Am237Am238Am239Am240Am241Am																		221Pu222Pu223Pu224Pu225Pu226Pu227Pu228Pu229Pu230Pu231Pu232Pu233Pu234Pu235Pu236Pu237Pu238Pu239Pu240Pu																		219Np220Np221Np222Np223Np224Np225Np226Np227Np228Np229Np230Np231Np232Np233Np234Np235Np236Np237Np238Np239Np																		215U216U217U218U219U220U221U222U223U224U225U226U227U228U229U230U231U232U233U234U235U236U237U238U																		211Pa212Pa213Pa214Pa215Pa216Pa217Pa218Pa219Pa220Pa221Pa222Pa223Pa224Pa225Pa226Pa227Pa228Pa229Pa230Pa231Pa232Pa233Pa234Pa235Pa236Pa237Pa																		208Th209Th210Th211Th212Th213Th214Th215Th216Th217Th218Th219Th220Th221Th222Th223Th224Th225Th226Th227Th228Th229Th230Th231Th232Th233Th234Th235Th236Th																		205Ac206Ac207Ac208Ac209Ac210Ac211Ac212Ac213Ac214Ac215Ac216Ac217Ac218Ac219Ac220Ac221Ac222Ac223Ac224Ac225Ac226Ac227Ac228Ac229Ac230Ac231Ac232Ac233Ac234Ac235Ac																		201Ra202Ra203Ra204Ra205Ra206Ra207Ra208Ra209Ra210Ra211Ra212Ra213Ra214Ra215Ra216Ra217Ra218Ra219Ra220Ra221Ra222Ra223Ra224Ra225Ra226Ra227Ra228Ra229Ra230Ra231Ra232Ra233Ra234Ra																		199Fr200Fr201Fr202Fr203Fr204Fr205Fr206Fr207Fr208Fr209Fr210Fr211Fr212Fr213Fr214Fr215Fr216Fr217Fr218Fr219Fr220Fr221Fr222Fr223Fr224Fr225Fr226Fr227Fr228Fr229Fr230Fr231Fr232Fr233Fr																		198Rn199Rn200Rn201Rn202Rn203Rn204Rn205Rn206Rn207Rn208Rn209Rn210Rn211Rn212Rn213Rn214Rn215Rn216Rn217Rn218Rn219Rn220Rn221Rn222Rn223Rn224Rn225Rn226Rn227Rn228Rn229Rn230Rn231Rn																		197At198At199At200At201At202At203At204At205At206At207At208At209At210At211At212At213At214At215At216At217At218At219At220At221At222At223At224At225At226At227At228At229At																		196Po197Po198Po199Po200Po201Po202Po203Po204Po205Po206Po207Po208Po209Po210Po211Po212Po213Po214Po215Po216Po217Po218Po219Po220Po221Po222Po223Po224Po225Po226Po227Po																		195Bi196Bi197Bi198Bi199Bi200Bi201Bi202Bi203Bi204Bi205Bi206Bi207Bi208Bi209Bi210Bi211Bi212Bi213Bi214Bi215Bi216Bi217Bi218Bi219Bi220Bi221Bi222Bi223Bi224Bi																		194Pb195Pb196Pb197Pb198Pb199Pb200Pb201Pb202Pb203Pb204Pb205Pb206Pb207Pb208Pb209Pb210Pb211Pb212Pb213Pb214Pb215Pb216Pb217Pb218Pb219Pb220Pb																		193Tl194Tl195Tl196Tl197Tl198Tl199Tl200Tl201Tl202Tl203Tl204Tl205Tl206Tl207Tl208Tl209Tl210Tl211Tl212Tl213Tl214Tl215Tl216Tl217Tl218Tl																		192Hg193Hg194Hg195Hg196Hg197Hg198Hg199Hg200Hg201Hg202Hg203Hg204Hg205Hg206Hg207Hg208Hg209Hg210Hg211Hg212Hg213Hg214Hg215Hg216Hg																		191Au192Au193Au194Au195Au196Au197Au198Au199Au200Au201Au202Au203Au204Au205Au206Au207Au208Au209Au210Au																		190Pt191Pt192Pt193Pt194Pt195Pt196Pt197Pt198Pt199Pt200Pt201Pt202Pt203Pt204Pt205Pt206Pt207Pt208Pt																		189Ir190Ir191Ir192Ir193Ir194Ir195Ir196Ir197Ir198Ir199Ir200Ir201Ir202Ir203Ir204Ir205Ir																		188Os189Os190Os191Os192Os193Os194Os195Os196Os197Os198Os199Os200Os201Os202Os203Os																	

Primary Decay Mode


Stable	n	e- capture
Fission	2n	p
α	β^+	2p
β^-	$2\beta^+$	3p
$2\beta^-$	e+	
Long-lived		
Estimated		
Unknown		

EPFL Heavy Charged Particles: Spontaneous Fission (1)


- ❑ Form of radioactive decay characteristic of very heavy isotopes.
- ❑ Spontaneous fission is **only** energetically feasible for $A > 230$
- ❑ Same process as induced fission:
 - Releases neutrons, fission products, gamma rays, neutrinos, etc
- ❑ Isotopes, for which spontaneous fission is a non-negligible decay mode can be used as neutron sources:
 - ^{252}Cf (half-life 2.645 years)
 - Applied to:
 - Detector calibration (well known spectrum of n)
 - inspect airline luggage for hidden explosives,
 - to gauge the moisture content of soil, materials stored in silos, etc.




Spontaneous Fission



Element-A	Half-life years	Spontaneous fission	
		Fraction	Rate per kg per second
uranium-233	2×10^5	1.6×10^{-12}	0.5
uranium-235	7.0×10^8	7×10^{-11}	0.06
uranium-238	4.5×10^9	5.4×10^{-7}	6
plutonium-239	2.4×10^4	4.4×10^{-12}	10
plutonium-240	6.6×10^3	5.0×10^{-8}	4.1×10^5
californium-252	2.6	0.03	2.3×10^{15}



The mass distribution of ^{252}Cf spontaneous fission fragments. Also shown is the corresponding distribution from fission of ^{235}U induced by thermal neutrons. (From Nervik.⁴)




The distribution in kinetic energy of the ^{252}Cf spontaneous fission fragments. The peak on the left corresponds to the heavy fragments, and that on the right to the light fragments. (From Whetstone.⁵)

- ❑ Fission fragments are medium-weight positive ions.
- ❑ Fission is generally asymmetric: clustering into light ($A \sim 108$) and heavy ($A \sim 143$) groups.
- ❑ Initial charge approaches Z of the fragment and is reduced by interaction with the material.
- ❑ Energy shared by the two fragments: ~ 185 MeV.
- ❑ Asymmetric distribution of kinetic energy: light fragments receive the largest.
- ❑ Sources must be thin to overcome self-absorption.

5 min break

Reminder: Non-ionizing vs. Ionizing radiation



$$\text{Energy of photon} = \Delta E_{\text{excited-final}}$$

- Emitted in the transition to lower energy levels in an excited nucleus
- Excited nuclei are produced by decay of a parent radionuclide:
 - Beta-decay leads to excited nucleus (parent half-life).
 - Energy is emitted as γ -photons (half-life $\sim$ ps).
 - The energy level structure reflects that of the daughter nucleus.
 - The γ -emission half-life is effectively that of the parent nucleus.




Figure 1.5 Decay schemes for some common gamma reference sources. Only major transitions are shown. The energies and yields per disintegration of X- and gamma rays emitted in each decay are listed below the diagram. (Data from Lederer and Shirley.¹)

☐ Nuclear states have very well defined energies:



- γ -rays emitted have well defined energies with very narrow peaks (nearly monoenergetic), $E_{\gamma} = E_i - E_f$.
- Can be used for detector calibration.

☐ Gamma reference sources are essential in radiation measurement laboratories:


- They consist of samples of β -emitters of a few μCi ($\sim 10^5$ Bq).
- Encased in plastic disks or rods and encapsulated to stop particulate radiation.
- Secondary radiation, annihilation photons or Bremsstrahlung can be significant.
- Radiation hazard is minimal due to low absolute activity.

☐ Energy is limited to about 2.8 MeV. Higher energies from:


- ^{56}Co (3.55 MeV, half-life 77 days)
- ^{16}N (6.13 and 7.11 MeV, half-life 7 s).




- Produced in nuclei undergoing β^+ decay.
- The positrons travel only a few millimetres before being stopped and annihilated by matter-antimatter interaction.
- This radiation is super-imposed on any gamma radiation emitted in the decay of the daughter nucleus.



$$e^+ + e^- \rightarrow 2\gamma (m_e c^2 = 0.511 \text{ MeV})$$





γ_1 : 1.274 MeV γ (100%)
annihilation radiation
Ne char. X-rays




- ❑ High energy γ -rays can be produced from nuclear energy transitions of higher-lying nuclear states.
- ❑ Nuclear reactions provide the needed high energy excited states.
- ❑ Reactions used:

- **Alpha absorption:**




Sources are a combination of an α -emitter and the target material. Large α -yields must be used for practical intensities: e.g. $6 \cdot 10^9$ Bq of ${}^{238}\text{PuO}_2$ and 200mg of ${}^{13}\text{C}$ give a source of 770 photons/s of 6.130 MeV γ -rays.

- **γ -rays from absorption of thermal neutrons** (radiative neutron-capture):
 - Intense flux from nuclear reactors or accelerator facilities.
 - Weaker fluxes from radioisotope sources of neutrons.
 - Gamma energies as high as 9 MeV.



Discovery of the neutron (1932, Chadwick)



1930: No effect in E field
 $\rightarrow \gamma?$

312

NATURE

[FEBRUARY 27, 1932]

Letters to the Editor

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, nor to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by α -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about 0.3 (cm.)^{-1} . Recently Mme. Curie-Joliot and M. Joliot found, when measuring the ionisation produced by this beryllium radiation in a vessel with a thin window, that the ionisation increased when matter containing hydrogen was placed in front of the window. The effect appeared to be due to the ejection of protons with velocities up to a maximum of nearly $3 \times 10^8 \text{ cm.}$

This again receives a simple explanation on the neutron hypothesis.


If it be supposed that the radiation consists of quanta, then the capture of the α -particle by the Be^9 nucleus will form a C^{13} nucleus. The mass defect of C^{13} is known with sufficient accuracy to show that the energy of the quantum emitted in this process cannot be greater than about 14×10^6 volts. It is difficult to make such a quantum responsible for the effects observed.

It is to be expected that many of the effects of neutron in passing through matter should resemble those of a quantum of high energy, and it is not easy to reach the final decision between the two hypotheses. Up to the present, all the evidence is in favour of the neutron, while the quantum hypothesis can only be upheld if the conservation of energy and momentum be relinquished at some point.


J. CHADWICK.

Cavendish Laboratory,
 Cambridge, Feb. 17.


1935



- Bremsstrahlung: electromagnetic radiation from acceleration of charged particles, such as e^-
- Bremsstrahlung has a continuous spectrum:
 - The fraction of e^- energy converted into Bremsstrahlung increases with the electron energy and target Z .
 - The average photon energy is a small fraction of the incident electron energy.



Spectrum produced by X-ray tube




□ Bremsstrahlung is used to produce X-rays from conventional X-ray tubes.

- Continuous spectrum altered by:
 - ❖ Filtration with absorber materials.
 - ❖ Peaked spectrum can be created by removing lower energy photons.
 - ❖ Can be used for calibration of detectors whose response changes only gradually with energy.

□ Bremsstrahlung also produced by:

- β -emitters interacting with shielding.
- Changes in nucleus electric field during β -decay.



Examples of measured pulse height spectra [using a NaI(Tl) scintillator] after filtration of an X-ray tube output using the indicated absorbers and tube voltages. (From Storm et al.¹⁵)

- ❑ X-Rays come from the re-arrangement of orbital electrons from excited atomic energy levels to ground states.
- ❑ Characteristic X-ray series depending on the shell with the vacancies.
- ❑ K-series is the most energetic, energy grows with Z:
 - Na (Z=11) 1 keV,
 - Ga (Z=31) 10 keV,
 - Ra (Z=88) 100 keV.
- ❑ The energy of the characteristic X-rays is unique. They can be used for element analysis.


Energies in the K-series:

$$E_{K_{\alpha}} = E_L - E_K$$

$$E_{K_{\beta}} = E_M - E_K$$


M

$$E_{K_{\max}} = E_{\text{free}} - E_K = K_{\text{Binding Energy}}$$



K-series of X-rays

❑ Excitation by **Electron Capture**:



- Nuclear electron capture from a K-shell electron creates a vacancy.
- The daughter atom is still neutral (Z-1), but a hole exists in one of the inner shells.
- The hole is filled by upper-shell electrons and gives a characteristic X-ray.

❑ Excitation by **Internal Conversion**:

- K-electrons are the most readily converted: the K-series is the most prominent.
- Gamma-ray de-excitation competes with this process, thus the K X-rays are produced together with γ -photons.
- If the energy of the converted electrons is high, Bremsstrahlung is also possible.

Some Radioisotope Sources of Low-Energy X-Rays


Nuclide	Half-Life	Weighted K_{α} X-Ray Energy	Fluorescent Yield	Other Radiations
^{37}Ar	35.1 d	2.957 keV	0.086	Some IB ^a
^{41}Ca	8×10^4 y	3.690	0.129	Pure
^{44}Ti	48 y	4.508	0.174	γ Rays at 68 and 78 keV
^{49}V	330 d	4.949	0.200	IB
^{55}Fe	2.60 y	5.895	0.282	Weak IB

^aIB represents inner bremsstrahlung.Data from Amlauer and Tuohy.¹⁷

- ❑ The yield of high-energy γ -rays from nuclear transitions is large compared to the characteristic X-rays.
- ❑ A pure X-ray source needs a radioisotope that decays by electron capture leading directly to a ground nuclear state of the daughter.
- ❑ ^{55}Fe is the most used because of its half-life and specific activity, and its nearly pure source of K-series of Mn at 5.9 keV, with very little Bremsstrahlung.
- ❑ Sources must be thin to prevent self-absorption of X-rays and have a high specific activity.

□ Excitation by **external radiation**:

- external sources of radiation used are: X-rays, e^- , α , etc.
- The radiation excites the parent atom which emits characteristic X-rays (isotropically).




□ The energy of X-rays depends on the target material:

Low $Z \rightarrow$ soft X-rays; High $Z \rightarrow$ hard X-rays.

Alpha Particle Sources Useful for Excitation of Characteristic X-rays		
	^{210}Po	^{244}Cm
Half-life	138 d	17.6 y
Alpha emissions	5.305 MeV (100%)	5.81, 5.77 MeV
Gamma-rays	803 keV (0.0011%)	43 keV (0.02%) 100 keV (0.0015%) 150 keV (0.0013%) 262 keV (1.4×10^{-4} %) 590 keV (2.5×10^{-4} %) 820 keV (7×10^{-5} %)
X-rays	Pb characteristic <i>L</i> and <i>M</i> (trace)	Pu characteristic <i>L</i> and <i>M</i>

Data from Amlauer and Tuohy.¹⁷



Compact source of characteristic X-rays with α -particle excitation.

No off switch



5 min break

❑ Synchrotron radiation:


- Generated by the B-field acceleration of **ultrarelativistic e^-** .
- Artificially by storage rings in a Synchrotron, or naturally by fast electrons moving through magnetic fields in space.
- The radiation spectrum typically spans from infrared (few eV) to X-rays (10 keV).

❑ Characteristics:


- High brightness and intensity, many orders of magnitude more than with X-rays produced in conventional X-ray tubes.
- High collimation, i.e. small angular divergence of beam.
- Low emittance, i.e. the product of source cross section and solid angle of emission is small.
- Wide tunability in energy/wavelength by monochromatization (sub eV up to the MeV range).
- High level of polarization (linear or elliptical) .
- Pulsed light emission (pulse durations at or below 1 ns).
- See, e.g., <http://www.psi.ch/sls/> (Swiss Light Source)




- ❑ Nuclear reactors (fission)
- ❑ Spallation sources



TRIGA Reactor



SINQ
Spallation
Source at PSI

TRIGA reactor



Spallation source example: Swiss Spallation Neutron Source SINQ

Principle:



Spallation source example: Swiss Spallation Neutron Source SINQ


PAUL SCHERRER INSTITUT
PSI




3D view: the Swiss Spallation Neutron Source SINQ




Where am I currently at PSI?



3 Neutron guide
Equipped with highly specialised mirrors built to reflect neutrons, these angular pipes guide the SINQ neutrons to the experimental stations.




2 When a proton collides with a lead nucleus, the nucleus heats up and in the process releases around ten neutrons. This is known as spallation.




1 In the target block
The SINQ produces neutrons when protons from the high intensity proton accelerator (HIPA, see p. 12 – 13) collide with a target block made of lead. Neutrons are then ejected at high speed in all directions. A cold source of liquid deuterium held in a 6,000 litre tank slows down or «moderates» the neutrons, which then pass into one of the neutron guides for use in specialised experiments. Neutrons that escape in other directions are blocked by protective layers and ultimately by the concrete cladding surrounding the target block. Some experiments operate with even slower «cold» neutrons. Their speed is further moderated in a 20-litre tank containing heavy water at round minus 250 centigrade.


Spallation reaction on Pb nucleus




Proton energy at 590 MeV




Spallation target



Proton beam




$$\sigma_x \sim 3.5 \text{ cm}, \sigma_y \sim 2 \text{ cm}$$




~360 Pb rods with SS / Zy-2 tubes

Target system



Application: Neutron science



PHYSICAL REVIEW LETTERS **124**, 081803 (2020)

Editors' Suggestion

Featured in Physics

Measurement of the Permanent Electric Dipole Moment of the Neutron

We present the result of an experiment to measure the electric dipole moment (EDM) of the neutron at the Paul Scherrer Institute using Ramsey's method of separated oscillating magnetic fields with ultracold neutrons. Our measurement stands in the long history of EDM experiments probing physics violating time-reversal invariance. The salient features of this experiment were the use of a ^{199}Hg comagnetometer and an array of optically pumped cesium vapor magnetometers to cancel and correct for magnetic-field changes. The statistical analysis was performed on blinded datasets by two separate groups, while the estimation of systematic effects profited from an unprecedented knowledge of the magnetic field. The measured value of the neutron EDM is $d_n = (0.0 \pm 1.1_{\text{stat}} \pm 0.2_{\text{sys}}) \times 10^{-26} \text{ e}\cdot\text{cm}$.

Cold neutrons via scattering in solid D₂ at 5K
 $\rightarrow E_{\text{kin}} < 500 \text{ neV}$




Applications: Neutron imaging and radiography

ELECTROCHEMISTRY

Revealing the impact of temperature in battery electrolytes via wavelength-resolved neutron imaging

Eric Ricardo Carreon Ruiz¹, Jongmin Lee^{1,2}, Markus Strobl^{2,3}, Natalie Stalder¹,
Genoveva Burca^{4,5,6}, Lorenz Gubler¹, Pierre Boillat^{1,2*}



11/1/2024



Neutron radiography



5 min break

48


<https://youtu.be/NnQdiDVOlvi>



(Don't brew coffee like that)

- ❑ Excitation levels with energies larger than the neutron binding energy are **not** produced as a result of any convenient radioactive decay process.
- ❑ Small neutron sources can be built based on:
 - Spontaneous Fission (SF).
 - (α ,n) nuclear reactions.
 - Neutron ejection of a nucleus induced by gamma radiation (photoneutrons).
 - Photofission: Neutrons are produced when gamma rays with high enough energies cause heavy nuclei to fission.
 - Fusion reactions of deuterium and/or tritium ions (neutron generator)

But wait...



However:

Helium-5 ($Z=2$, $N=3$)

Discovered 1937 [5]

$t_{1/2} = 602 \pm 22$ ys [5]

yocto = 10^{-24}

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--


Red = n emission

- ❑ SF appears in many of the transuranic heavy nuclides ($Z > 92$).
- ❑ It produces:
 - Several fast neutrons.
 - Heavy fission products.
 - Prompt fission γ -rays (in ns).
 - β and γ activity of the fission products.
- ❑ The neutron energy spectrum is peaked between 0.5 and 1 MeV and can be approximated by:

$$\frac{dN}{dE} = E^{1/2} e^{-E/T}$$

Spontaneous fission rates

U-235	5.60E-03 f/s-kg
U-238	6.93 f/s-kg
Pu-239	7.01 f/s-kg
Pu-240	489,000 f/s-kg



Measured neutron energy spectrum from the spontaneous fission of ^{252}Cf .
(From Batenkov et al.¹⁸)

- ❑ The most common SF source is ^{252}Cf ($Z=98$).
- ❑ Characteristics:
 - **2.6 year half-life**
 - Dominant decay mechanism is α -decay
 - **Extremely radioactive** (1 mg spontaneously emits 170 million neutrons per minute)
- ❑ Produced in nuclear reactors: from irradiation of Cm with α -particles (but how to produce Cm?)
- ❑ ^{252}Cf neutron sources are typically 1/4" to 1/2" in diameter and 1" to 2" in length. The price of a typical ^{252}Cf neutron source : \$50,000.
- ❑ Some uses of the ^{252}Cf sources:
 - Neutron start-up source for some nuclear reactors, calibrating instrumentation.
 - Treatment of certain cervical and brain cancers where other radiation therapies are ineffective.
 - Radiography of aircraft to detect metal fatigue.
 - Airport neutron-activation detectors of explosives.
 - Neutron moisture gauges used to find water and petroleum layers in oil wells.
 - Portable neutron source in gold and silver prospecting for on-the-spot analysis.



❑ Neutrons are produced in (α ,n)-reactions.

- The source of α is a radioisotope.
- Self-contained sources have a mixture of an α -emitter and a target material.
- The maximum neutron yield comes from ^9Be as a target material.

❑ Sources of (α ,n) neutrons:

- All α -emitters are actinides: ^{239}Pu , ^{210}Po , ^{241}Am , ^{244}Cm , ^{226}Ra , etc.
- Most α -particles are absorbed in the target, only 1 in 10^4 reacts with Be.
- They form stable alloy of the form MBe_{13} (M=actinide metal), with no intermediate loss of energy for the α -particle.

Basic reaction for Be neutron sources:



Typical double-walled Be(α ,n) source

Characteristics of Be(α , n) Neutron Sources							
	Source	Half-Life	E_α (MeV)	Neutron Yield per 10^6 Primary Alpha Particles		Percent Yield with $E_n < 1.5$ MeV	
				Calculated	Experimental	Calculated	Experimental
Most widely used	$^{239}\text{Pu}/\text{Be}$	24000 y	5.14	65	57	11	9–33
High n yields, high specific activity	$^{210}\text{Po}/\text{Be}$	138 d	5.30	73	69	13	12
Low γ background, simple α decay process.	$^{238}\text{Pu}/\text{Be}$	87.4 y	5.48	79 ^a	—	—	—
	$^{241}\text{Am}/\text{Be}$	433 y	5.48	82	70	14	15–23
	$^{244}\text{Cm}/\text{Be}$	18 y	5.79	100 ^b	—	18	29
Ideal	$^{242}\text{Cm}/\text{Be}$	162 d	6.10	118	106	22	26
Intense γ background radiation from daughters	$^{226}\text{Ra}/\text{Be}$ + daughters	1602 y	Multiple	502	—	26	33–38
	$^{227}\text{Ac}/\text{Be}$ + daughters	21.6 y	Multiple	702	—	28	38

^aFrom Anderson and Hertz.²² All other data as calculated or cited in Geiger and Van der Zwan.²³

^bDoes not include a 4% contribution from spontaneous fission of ^{244}Cm .

- Energy spectra of all sources with ^9Be are similar.
 - Differences reflect the small variations in the primary α -particle energies.
 - Thick sources have more „spread“, i.e, the originally discrete α -energy spectrum is washed out.





Figure 1.12 Measured energy spectra for neutrons from a $^{239}\text{Pu}/\text{Be}$ source containing 80 g of the isotope. (From Anderson and Neff.²⁵)

Additional sources of (α, n)

Alternative (α, n) Isotopic Neutron Sources			
Target	Reaction	Q-Value	Neutron Yield per 10^6 Alpha Particles
Natural B	$^{10}\text{B}(\alpha, n)$	+1.07 MeV	13 for ^{241}Am alpha particles
	$^{11}\text{B}(\alpha, n)$	+0.158 MeV	
F	$^{19}\text{F}(\alpha, n)$	-1.93 MeV	4.1 for ^{241}Am alpha particles
Isotopically separated ^{13}C	$^{13}\text{C}(\alpha, n)$	+2.2 MeV	11 for ^{238}Pu alpha particles
Natural Li	$^7\text{Li}(\alpha, n)$	-2.79 MeV	
Be (for comparison)	$^9\text{Be}(\alpha, n)$	+5.71 MeV	70 for ^{241}Am alpha particles



Data from Lorch¹⁹ and Geiger and Van der Zwan.²⁷





Neutron energy spectra from alternative (α, n) sources. (^7Li data from Geiger and Van der Zwan,²⁷ remainder from Lorch.¹⁹)

□ Excitation of the nucleus by high energy γ -rays:

- Photoneutron sources combine a powerful γ -emitter with a target isotope.
- Only two target nuclei are practical: ^9Be and ^2H :




- Photoneutron sources produce monoenergetic neutrons (for monoenergetic gammas).
- They need very large γ -ray activities (1 out of 10^5 - 10^6 photons produces 1 n).
- Common emitters are:
 ^{226}Ra , ^{124}Sb , ^{72}Ga , ^{140}La , ^{24}Na .
- Very short half-lives require reactivation in a nuclear reactor between uses.
- Alternative: e^- accelerator + target




Neutron spectra calculated for the photoneutron source dimensions shown in Fig. 1.15. The gamma emitters are either ^{72}Ga or ^{24}Na . The outer shells are either deuterated polyethylene (CD_2) or beryllium (Be).

Linear accelerator (linac) + photoneutron target

Cross sections for neutron production (orange)
Vs. linac Xray spectrum (blue)



Measured neutron energies for
different targets and sources
DU = depleted uranium



Photoneutron Source Characteristics					
Gamma-Ray Emitter	Half-Life ^a	Gamma Energy ^a (MeV)	Target	Neutron Energy ^b (keV)	Neutron Yield (n/s) for 10 ¹⁰ Bq Activity ^c
²⁴ Na	15.0 h	2.7541	Be	967	340,000
		2.7541	D	263	330,000
²⁸ Al	2.24 min	1.7787	Be	101	32,600
³⁸ Cl	37.3 min	2.1676	Be	446	43,100
⁵⁶ Mn	2.58 h	1.8107	Be	129	91,500
		2.1131		398	
		2.9598		1,149	
		2.9598	D	365	162
⁷² Ga	14.1 h	1.8611	Be	174	64,900
		2.2016		476	
		2.5077		748	
		2.5077	D	140	25,100
⁷⁶ As	26.3 h	1.7877	Be	109	3,050
		2.0963		383	
⁸⁸ Y	107 d	1.8361	Be	152	229,000
		2.7340		949	
		2.7340	D	253	160
^{116m} In	54.1 min	2.1121	Be	397	15,600
¹²⁴ Sb	60.2 d	1.6910	Be	23	210,000
¹⁴⁰ La	40.3 h	2.5217	Be	760	10,200
		2.5217	D	147	6,600
¹⁴⁴ Pr	17.3 min	2.1856	Be	462	690

^aDecay data from Ref. 1.

^bCalculated for $\theta = \pi/2$, approximate midpoint of primary spectrum.

^cMonte Carlo calculations for the source dimensions given in Fig. 1.15. Outer target shells are either metallic Be or deuterated polyethylene. Core materials assumed to be NaF, Al, CCl₄, MnO₂, Ga₂O₃, As₂O₃, Y₂O₃, In, Sb, La₂O₃, and Pr₂O₃.



Source: G. F. Knoll, "Radioisotope Neutron Sources," Chap. 2 in *Neutron Sources for Basic Physics and Applications*, Pergamon Press, New York, 1983.

- Two of the most common reactions (with their Q-values) are fusion reactions:




- Characteristics of the sources:

- Low coulomb barrier (low Z target) requires accelerating potentials of 100-300 kV for the charged particles.
- All neutrons mono-energetic
- Typical yields: 1 mA ${}^2\text{H}$ beam will produce 10^9 n/s for ${}^2\text{H}$ and 10^{11} n/s for ${}^3\text{H}$ targets $\rightarrow$ compact, portable source
- Other reactions ${}^9\text{Be}(d,n)$, ${}^7\text{Li}(p,n)$ and ${}^3\text{H}(p,n)$ require large accelerators ($Q < 0$).




Neutron generator application: Nuclear security




Active neutron and gamma-ray imaging of highly enriched uranium for treaty verification

Michael C. Hamel , J. Kyle Polack, Marc L. Ruch, Matthew J. Marcath, Shaun D. Clarke & Sara A. Pozzi



^{252}Cf production in a reactor



²⁵²Cf properties

Californium-252 (Z=98, N=154)

Discovered 1954 [5]


$t_{1/2} = 2.645 \pm 0.008$ y [5]

$J^\pi = 0^+$ [5]

Decay modes

$\alpha = 96.9\%$ [5]

Spontaneous fission = 3.10% [5]



Neutron energy spectrum from ²⁵²Cf sf

Uranium-235 (Z=92, N=143)

Discovered 1935 [5]

Natural abundance 0.7204% [5]

$t_{1/2} = 704 \pm 1$ My [5]

$J^\pi = 7/2^-$ [5]

Decay modes

$\alpha = 100\%$ [5]

Spontaneous fission = $7.00 \times 10^{-9}\%$ [5]

$20\text{Ne} = 8.00 \times 10^{-10}\%$ [5]

$25\text{Ne} \sim 8.00 \times 10^{-10}\%$ [5]

$28\text{Mg} = 8.00 \times 10^{-10}\%$ [5]

Uranium-238 (Z=92, N=146)

Discovered 1896 [5]

Natural abundance 99.2742% [5]

$t_{1/2} = 4.463 \pm 0.003$ Gy [5]

$J^\pi = 0^+$ [5]

Decay modes


$\alpha = 100\%$ [5]

Spontaneous fission = 0.0000544% [5]

$2\beta^- = 2.20 \times 10^{-10}\%$ [5]

	β^+	β^+	β^+	β^+	β^+	β^+	β^+	e- capture	β^+	α	β^-	α
Pu	²³¹ Pu β^+	²³² Pu e- capture	²³³ Pu β^+	²³⁴ Pu e- capture	²³⁵ Pu β^+	²³⁶ Pu α	²³⁷ Pu e- capture	²³⁸ Pu α	²³⁹ Pu α	²⁴⁰ Pu α	²⁴¹ Pu β^-	²⁴² Pu α
Np	²³⁰ Np β^+	²³¹ Np β^+	²³² Np β^+	²³³ Np β^+	²³⁴ Np β^+	²³⁵ Np e- capture	²³⁶ Np e- capture	²³⁷ Np α	²³⁸ Np β^-	²³⁹ Np β^-	²⁴⁰ Np β^-	²⁴¹ Np β^-
U	²²⁹ U β^+	²³⁰ U α	²³¹ U e- capture	²³² U α	²³³ U α	²³⁴ U α	²³⁵ U α	²³⁶ U α	²³⁷ U β^-	²³⁸ U α	²³⁹ U β^-	²⁴⁰ U β^-
Pa	²²⁸ Pa β^+	²²⁹ Pa e- capture	²³⁰ Pa β^+	²³¹ Pa α	²³² Pa β^-	²³³ Pa β^-	²³⁴ Pa β^-	²³⁵ Pa β^-	²³⁶ Pa β^-	²³⁷ Pa β^-	²³⁸ Pa β^-	²³⁹ Pa β^-
Th	²²⁷ Th α	²²⁸ Th α	²²⁹ Th α	²³⁰ Th α	²³¹ Th β^-	²³² Th α	²³³ Th β^-	²³⁴ Th β^-	²³⁵ Th β^-	²³⁶ Th β^-	²³⁷ Th β^-	²³⁸ Th β^-
Ac	²²⁶ Ac β^-	²²⁷ Ac β^-	²²⁸ Ac β^-	²²⁹ Ac β^-	²³⁰ Ac β^-	²³¹ Ac β^-	²³² Ac β^-	²³³ Ac β^-	²³⁴ Ac β^-	²³⁵ Ac β^-	²³⁶ Ac β^-	²³⁷ Ac β^-

Comparison of neutron source spectra for detector calibration



Roman K. Piper*, Andrey V. Mozhayev*, Mark K. Murphy*, Alan K. Thompson†

*Pacific Northwest National Laboratory, P.O. Box 999, Richland, WA 99352

†National Institute of Standards and Technology, 100 Bureau Drive, Gaithersburg, MD 20899

Table 1

Overview of Neutron Calibration Sources and Properties

Neutron source (type)	Radionuclide half-life (years)	Number of sources needed for 15 years ^a	Relative neutron intensity of individual source ^a	Radionuclide nominal specific neutron intensity (s ⁻¹ g ⁻¹)	Nominal mass of radionuclide needed for 10 ⁹ s ⁻¹ source (g)
DD (NG)	n/a	3-5	1.000	n/a	n/a
DT (NG) ^b	12.3	3-5	1.147-1.087	2.28×10^{13}	0.00004
²⁵² Cf (SF)	2.645	2	2.285	2.31×10^{12}	0.0004
²¹⁰ PoBe (α,n)	0.378	13	2.409	1.08×10^{10}	0.1
²⁴⁴ CmBe (α,n)	18.1	1	1.315	2.06×10^8	5
²³⁸ PuBe (α,n)	87.74	1	1.060	4.12×10^7	24
²⁴⁸ Cm (SF)	348000	1	1.000	3.94×10^7	25
²²⁶ RaBe (α,n)	1620	1	1.003	1.27×10^7	79
²⁴⁴ Cm (SF)	18.1	1	1.315	1.11×10^7	90
²⁴¹ AmBe (α,n)	433.6	1	1.012	8.23×10^6	122
²³⁹ PuBe (α,n)	24100	1	1.000	1.49×10^5	6700

- ❑ Glenn F. Knoll, “*Radiation Detection and Measurement*”, John Wiley & Sons (4th edition, 2010, and 3rd edition, 2000)
- ❑ James E. Martin, “*Physics for Radiation Protection*”, Wiley-VCH (2nd edition, 2006)
- ❑ James E. Turner, “*Atoms, Radiation, and Radiation Protection*”, Wiley-VCH (3rd edition, 2007)