

BIO-212 – Kinetics and Catalysis – Exercise session 13
11.12.2025

1) True/False

Which of the following statements about enzyme-substrate binding is correct?

- A) Enzyme-substrate binding is always an irreversible process.
- B) Enzymes bind substrates through a covalent bond in most reactions.
- C) The enzyme's active site is specific for a one or multiple substrates.
- D) Substrate binding occurs at random sites on the enzyme.

What is the primary function of an enzyme in a chemical reaction?

- A) To stabilize the transition state, leading to a higher activation energy of the reaction.
- B) To decrease the activation energy, thereby increasing the reaction rate.
- C) To change the equilibrium constant of the reaction to favour substrate formation.
- D) To bind its substrate with high affinity, forming a stable enzyme-substrate complex.

In Michaelis-Menten kinetics, what happens to the reaction rate when the substrate concentration is much higher than the Michaelis constant ($[S] \gg K_m$)?


- A) The reaction rate increases linearly with increasing substrate concentration.
- B) The reaction rate is directly proportional to the substrate concentration.
- C) The reaction rate is at its maximum value
- D) The reaction rate is slower than at higher substrate concentrations due to substrate inhibition.

Which of the following statements about enzyme inhibitors is true?

- A) Competitive inhibitors decrease the V_{max} of the enzyme.
- B) Non-competitive inhibitors increase the K_m , but do not affect the V_{max} .
- C) Competitive inhibitors can be partially overcome by increasing substrate concentration.
- D) Non-competitive inhibitors bind to the enzyme-substrate complex, leading to an increase of the K_m and a decrease of the V_{max} .

2) Effects of enzyme catalysis


Which of the following effects would be brought about by an enzyme catalysing the simple reaction:



- a) Decreased K'_{eq}
- b) Increased k_1
- c) Increased $\Delta G^\ddagger (E_A)$
- d) Decreased $\Delta G^\ddagger (E_A)$
- e) Increased K'_{eq}
- f) Decreased ΔG°
- g) Increased k_2


3) First-order kinetics

You are studying the kinetics of different compounds containing radioactive elements, with particular interest in iodine-123 (^{123}I) which is important for medical imaging of thyroid function and follows a **first-order decay kinetics** into ^{123}Te . A 15 μg sample of ^{123}I has decreased to 7.5 μg after 13 hours. After how much time will it decay to less than 0.95 μg ?



4) Degradation of H₂O₂

- a. During aerobic metabolism reactive oxygen species (ROS) may be formed, among them hydrogen peroxide (H₂O₂). The enzyme **Catalase** is found in most aerobic organisms where it catalyzes the following reaction:



Can you reason why it is prerogative for organisms to prevent the accumulation of hydrogen peroxide?

- b. The uncatalyzed decomposition of hydrogen peroxide as shown above proceeds spontaneously, as it has a negative Gibbs free energy under standard conditions in aqueous solution.

How does the presence of Catalase impact the Gibbs free energy of the reaction? Can you reason why aerobic organisms rely on the enzymatically catalyzed degradation of hydrogen peroxide over just spontaneous decomposition?

- c. In a laboratory course, you are tasked with measuring the activation energy of the degradation of hydrogen peroxide when catalyzed by catalase. For this purpose, you add purified catalase to a solution containing hydrogen peroxide. You follow the reaction by measuring the rate of oxygen formation immediately after mixing. The following rate constants were determined at two temperatures:

Temperature (K)	Rate constant k (s ⁻¹)
293	7
303	8

- i. **Transform the Arrhenius equation into linear form and complete the following table using the values provided above.**

(Hint: apply a ln() transformation to both sides of the equation.)


1/T	ln(k)

- ii. You want to calculate the activation energy (E_a) using the linearized Arrhenius equation. **You may assume the activation energy (E_a) and the pre-exponential factor (A) to be constants.**

Plot $\ln(k)$ against $1/T$. Why does this result in a straight line?

- iii. Using your plot, derive the activation energy (E_a).

5) Carboxypeptidase





The enzyme carboxypeptidase cleaves carboxy-terminal amino acids from its peptide substrates. Carboxypeptidase is constituted by a single peptide chain of 307 amino acids. The two catalytic residues of carboxypeptidase are Arg-145 and Glu-270.

- If carboxypeptidase was a perfect alpha helix what would the distance between these amino acids be in Å?
- Explain how these two amino acids, distant from each other in the primary structure, can nevertheless catalyse a reaction taking place in the range of few nanometers.
- If only these 2 amino acids are involved in catalysis, why is carboxypeptidase constituted by 307 residues?

6) Michaelis Menten Kinetics

- a) At what substrate concentration would an enzyme with a k_{cat} of 30.0 s^{-1} and a K_m of 0.008 mol/L operate at one-fifth of its maximum rate?
- b) Determine the fraction of the maximum reaction rate (V_o/V_{max}) that would be obtained at the following concentrations of $[S]$: $0.5 K_m$, $2 K_m$, $5 K_m$
- c) Suppose a noncompetitive inhibitor is introduced with a $\alpha'=5$. How does the inhibitor affect the maximum velocity V_{max} and what is the new V_{max} in the presence of the inhibitor? Assume that the enzyme concentration is 50 nmol/L .

7) Thymidylate Synthase Kinetics



The enzyme thymidylate synthase (TS) catalyzes the reductive methylation of 2'-deoxyuridine 5'-monophosphate (dUMP) to 2'-deoxythymidine 5'-monophosphate (dTMP), a key building block for DNA. This reaction is crucial for DNA replication and repair, making TS an important target in cancer treatment. Researchers have found that the K_m for dUMP is $4\mu\text{M}$ and the k_{cat} of the enzyme is 20 min^{-1} .

- In an experiment, the [dUMP] used was 6mM , and the initial velocity (V_0) was 480 nM min^{-1} . What was the concentration of TS used in in this experiment?
- In a second experiment, the enzyme concentration [TS] is $0.5\text{ }\mu\text{M}$ and $V_0 = 5\text{ }\mu\text{M min}^{-1}$. What was the [dUMP] in this experiment?
- You are developing a new chemotherapeutic drug that acts as a competitive inhibitor of TS with an $\alpha = 10$. This inhibitor is added to a mixture of enzyme and substrate, resulting in a V_0 of 240 nM min^{-1} . At an enzyme concentration of 24 nM , which dUMP concentration has been used in the experiment?