




# CH-110 Advanced General Chemistry I


Prof. A. Steinauer  
[angela.steinauer@epfl.ch](mailto:angela.steinauer@epfl.ch)

# Wavefunctions and Energy Levels

Topic 1C




# Overview Chapter 1 (Focus 1: Atoms)




# The Wavefunction and Its Interpretation

Topic 1C.1



Classical  
mechanics:  
Fixed predictable  
path




Quantum  
mechanics:  
Probability  
distribution

Image source: ChatGPT.  
*Classical vs. quantum paths.*

## 1C.1 The wavefunction and its interpretation

### The particle-in-a-box model



The wavefunction of the particle in its lowest-energy state  
(**ground state,  $n = 1$** ):

$$\psi_1(x) = \left(\frac{2}{L}\right)^{\frac{1}{2}} \sin\left(\frac{\pi x}{L}\right)$$

And therefore:

$$\psi_1^2(x) = \left(\left(\frac{2}{L}\right)^{\frac{1}{2}} \sin\left(\frac{\pi x}{L}\right)\right)^2 = \frac{2}{L} \sin^2\left(\frac{\pi x}{L}\right)$$

And  $P(x=0) = P(x=L) = 0$

## 1C.1 The wavefunction and its interpretation

### The particle-in-a-box model

- **Physical analog:** a bead free to slide along a rigid rod lying between two walls a distance  $L$  apart.
- **Classical mechanics:** the bead has the same probability of being found on the rod at any point inside the box, any speed, any kinetic energy




Image source: ChatGPT.  
*Bead on a rigid rod..*

# 1C.1 The wavefunction and its interpretation

## The particle-in-a-box model: $n < 1$

$$\psi_n(x) = \left(\frac{2}{L}\right)^{\frac{1}{2}} \sin\left(\frac{n\pi x}{L}\right), \quad n = 1, 2, \dots$$

- The integer  $n$  labels the wavefunctions and is called a "quantum number".

### A quantum number:

- Is an integer (or sometimes a half-integer, such as  $\frac{1}{2}$ , see Topic 1D)
- Labels a wavefunction
- Specifies a state
- Can sometimes be used to calculate the value of a property of the system, e.g. energy.





Figure 1C.3

## 1C.1 The wavefunction and its interpretation

### Wavefunctions vs. guitar strings

Because the particle acts like a wave with zero amplitude at each end of the box

- Only wavefunctions with **certain wavelengths** can exist in the box
- Think of a **guitar string**: because it is tied down at each end, it can support only shapes like the ones shown.
- The shapes of the wavefunctions for the particle in the box are the same as the displacements of a vibrating string.



# 1C.1 The wavefunction and its interpretation

## Wavefunctions vs. guitar strings





Figure 1C.3

## 1C.1 The wavefunction and its interpretation

### Connecting concepts across Topics 1B and 1C

- **Planck and Einstein:** energy is quantized → light behaves as particles (photons)
- **De Broglie:** matter particles (e.g. electrons) have wave nature
- **Quantum chemistry:** focus on most probable position, not exact location.
- **Schrödinger:** equation provides wavefunction ( $\psi$ ) and energy of a system.
- **Goal:** knowing  $\psi(r)$  and  $E$  gives all electron properties; information extracted through probability.

## 1C.1 The wavefunction and its interpretation

### Summary

The **probability density** for a particle at a location is proportional to the square of the wavefunction at that point; places where the wavefunction passes through zero are called **nodes**, and the particle will not be found there. A wavefunction is found by **solving the Schrödinger equation** for the particle and recognizing the existence of certain **boundary conditions**.




URL

Quantum mechanics and guitars 

# The Quantization of Energy

Topic 1C.2



## 1C.2 The quantization of energy

### Energies of a particle in a box

The wavefunctions associated with different quantum numbers also have different energies associated with them. **How do we calculate these energies?**

## 1C.2 The quantization of energy

### **Energies of a particle in a box**

## 1C.2 The quantization of energy

What does this equation tell you?

$$E_n = \frac{n^2 h^2}{8mL^2}$$

Important formula.  
Need to know how to apply.

- **Mass in denominator:** Heavier particles  $\rightarrow$  lower, closer energy levels. Lighter particles  $\rightarrow$  higher, more widely spaced levels.
- **Box length effect:** Smaller box ( $L \downarrow$ )  $\rightarrow$  higher, more widely spaced levels. Larger box ( $L \uparrow$ )  $\rightarrow$  lower, closer levels.
- $n$  can only be integer values  $\rightarrow$  **energy is quantized!**




Figure 1C.4

## 1C.2 The quantization of energy

### Energy separation between neighboring levels

Important formula.  
Need to know  
how to apply.

$$E_{n+1} - E_n = \frac{(n+1)^2 h^2}{8mL^2} - \frac{n^2 h^2}{8mL^2}$$
$$= \{(n+1)^2 - n^2\} \frac{h^2}{8mL^2} = \frac{(2n+1)h^2}{8mL^2}$$

As  $L$  or  $m$  increases, the separation between neighboring energy levels decreases.

→ For large  $m$  or  $L$  (macroscopic systems), the levels are so close together they appear continuous.




Figure 1C.4

## 1C.2 The quantization of energy


### Quantization of energy and the atomic spectrum of hydrogen

Energy is quantized → this realization is key to understanding the **atomic spectrum of hydrogen** (Topic 1A).

Particle in a box	Hydrogen atom
1D	3D
Physical walls	No physical walls, and electrons are confined by pull of the nucleus
Energy quantized	

## 1C.2 The quantization of energy

What are these lines?!



A spectral line arises from a **transition of an electron between allowed energy levels.**

The difference in energy is carried away as a photon.

$$h\nu = E_{upper} - E_{lower} = \Delta E$$

Important formula.  
Need to know how to apply.

This equation is known as the **Bohr frequency condition.**

## 1C.2 The quantization of energy

### Terminology

- **Ground state:** The lowest energy state of an atom.
- **Excited state:** Any higher energy state above the ground state.
- **Excitation:** Moving to a higher energy state
  1. **Radiative excitation (photon absorption):** photon absorbed, electron promoted.
  2. Nonradiative excitation: energy gained from collisions, chemical reactions, or electric fields.
- **Relaxation:** General term for return from an excited state to a lower state.
  1. **Radiative relaxation (photon emission):** energy released as a photon .
  2. Nonradiative relaxation: energy released in other forms (e.g., vibrations, heat).

## 1C.2 The quantization of energy


### Example 1C.1: Calculating the energies of a particle in a box

We will estimate the energies of the hydrogen atom.

Treat hydrogen atom as a **one-dimensional box of length 150. pm** (the approximate diameter of the atom) with one electron.

Predict energy level separation **between the lowest and next higher** energy levels.

If the electron falls from the **upper level to the lower level**, what would be the wavelength of the radiation emitted as a photon?



## 1C.2 The quantization of energy

### **Example 1C.1: Calculating the energies of a particle in a box**


## 1C.2 The quantization of energy

### **Example 1C.1: Calculating the energies of a particle in a box**

## 1C.2 The quantization of energy

### Example 1C.1: Calculating the energies of a particle in a box

*Evaluate* This wavelength corresponds to 24.7 nm. The experimental value for the actual transition in a hydrogen atom is 122 nm. Although there is a big discrepancy, an atom does not have the hard boundaries that confine a particle in a box, and is three-dimensional. The fact that the predicted wavelength has nearly the same order of magnitude as the actual value suggests that a quantum theory of the atom, based on a more realistic three-dimensional model, should give good agreement.



## 1C.2 The quantization of energy

### Particle in a box: zero-point energy

- Surprising implication of equation:  $E_n = \frac{n^2 h^2}{8mL^2}$
- A particle in a container **cannot have zero energy**.
- The lowest value of  $n$  is 1.
- Lowest energy is  $E_1 = \frac{h^2}{8mL^2}$  (**Zero-point energy**)
- What this means: A particle can never be perfectly still when it is confined between two walls, it must always possess an energy, in this case, at least the kinetic energy  $\frac{h^2}{8mL^2}$ .

## 1C.2 The quantization of energy

### The shapes of the wavefunctions

- The first two wavefunctions ( $n = 1, 2$ ) illustrate how **particle location depends on quantum state**.
- $n = 1$  with  $E = h^2/8mL^2$ : Particle most likely at the center.
- $n = 2$ ,  $E = h^2/2mL^2$ : Particle least likely at the center, most likely between center and walls.
- Shading indicates probability density – the likelihood of finding the particle at each position.




Figure 1C.5

## The skills you have mastered are the ability to

- ❑ Describe the origin and shapes of the wavefunctions of a particle in a box.
- ❑ Calculate the allowed energies of a particle in a box and explain how they depend on the length of the box and the mass of the particle.
- ❑ Explain what is meant by zero-point energy and accounts for its origin

**Summary: You now know that the location of a particle is expressed by a wavefunction, the square of which expresses the probability (as a probability density) that the particle will be found in each region of space. You also know that a wavefunction is found by solving the Schrödinger equation and that one consequence of the wavefunction having to fit into a region of space is that a particle confined to a region can have only certain discrete energies known as energy levels.**