


# Recap from last lecture

- What are anomalies and how are they calculated?
- Paris agreement: keep global temperature rise below  $1.5^{\circ}\text{C}$  above preindustrial levels
- Warming is heterogeneous




	No.	Date	Topics	Deadlines / tentative
Basics	1.	11.09.2025	Introduction to the climate system	Questionnaire (not graded)
	2.	18.09.2025	Climate System, Radiation	
	3.	25.09.2025	Earth's Energy balance, Greenhouse effect, Radiative transfer, aerosols & clouds	launch of first assignment
	4.	02.10.2025	Aerosols & clouds, Radiative Forcing, Feedback mechanisms	Launch of poster project
	5.	09.10.2025	Feedback mechanisms, Climate Sensitivity	
	6.	16.10.2025	Paleoclimate	submission of Poster proposal (graded)
Present and future Climate change	7.	30.10.2025	Climate variability	
	8.	06.11.2025	Paris Agreement, IPCC – report, Climate scenarios (RCPs, SSPs),	
	9.	13.11.2025	Tipping elements, 1.5 vs 2.0°C vs warmer, Emissions, Carbon budget, Emissions Gap	submission of Poster draft (graded)
	10.	20.11.2025	Carbon budget, Emissions Gap, Metrics	
	11.	27.11.2025	Metrics, Extreme Events	submission of assignment (graded)
Actions	12.	04.12.2025	Polar climate change	
	13.	11.12.2025	Mitigation measures	Poster Conference (graded)
	14.	18.12.2025	Climate engineering, questions and answers session	fill in Questionnaire in exercises (not graded)

# Fundamentals of Radiation

Chapter 4 in Wallace and Hobbs

# Spectrum of electromagnetic radiation



## Some definitions

**Visible:** 0.39 – 0.76  $\mu\text{m}$ , colors

**Monochromatic:** one single color, i.e. one single wavelength


**Solar radiation:** < 4  $\mu\text{m}$ , shortwave

**Terrestrial radiation:** > 4  $\mu\text{m}$ , longwave

**Near infrared:** 0.76 – 4.0  $\mu\text{m}$  (solar radiation)

**Thermal Infrared:** > 4  $\mu\text{m}$ , terrestrial radiation

**Microwave** ( $10^{-3}$  –  $10^{-1}$  m): not important for Earth's energy balance, but used in remote sensing (radar), can penetrate through clouds



- Globally averaged, the surface has a net surplus of **radiant** energy while the atmosphere has a net loss. To make up for this imbalance, **sensible** (conduction & convection) and **latent heat** (evaporation) are transferred from the surface to the atmosphere.
- At the surface, temperatures would be 33°C cooler without the **natural greenhouse effect** driven by water vapor, CO<sub>2</sub>, O<sub>3</sub> and clouds. These absorb surface infrared radiation and re-emit most of it back to the surface.


### We will learn about:

- characteristics of solar and terrestrial radiation,
- how solar radiation passes through the atmosphere,
- how terrestrial radiation passes through the atmosphere,
- why there is a net loss of energy in the atmosphere,
- how greenhouse gases, aerosols and clouds interact with radiation.

- **Radiation:** emission or transmission of energy in the form of waves or particles through space or through a material medium
- **Irradiance/Flux Density:** radiant flux (Watts) **received** by a **surface** per unit area, Symbol:  $F$ ; Unit:  $W\ m^{-2}$ 
  - **Spectral irradiance** is the irradiance of a surface per unit frequency or wavelength; Unit:  $W\ m^{-2}\ nm^{-1}$  or  $W\ m^{-2}\ Hz^{-1}$
  - **Insolation/solar irradiance:** The incident radiant energy emitted by the sun which reaches a unit area over a period of time, typically measured over a horizontal area at the Earth's surface or at the top of Earth's atmosphere. Units:  $W\ m^{-2}$
- **Radiance:** is the radiant flux emitted, reflected, transmitted or received by a given surface, per unit solid angle per unit projected area, Symbol:  $I$ ; Units:  $W\ m^{-2}\ sr^{-1}$ 
  - **Radiant intensity:** radiant flux emitted, reflected, transmitted or received, per unit solid angle, Units:  $W\ sr^{-1}$
  - **Spectral intensity:** is the radiant intensity per unit frequency or wavelength, Units:  $W\ sr^{-1}\ Hz^{-1}$  or  $W\ sr^{-1}\ nm^{-1}$




# Solar and terrestrial emission spectra



# Solar and terrestrial emission spectra

Why do they radiate?





# Blackbody radiation

Definition of blackbody: matter or object that absorbs all radiation incident on it, no reflection. Examples: coal, cave

**Blackbodies emit radiation.** The radiation *emitted* by a blackbody is *isotropic* (same intensity independent of location/direction) and depends only on the **temperature of the body**, not on the composition of the blackbody, nor on whether any radiation is incident on the blackbody. The blackbody is losing energy by emission of radiation, so heat must be supplied to keep its temperature constant.

Most objects are not blackbodies. Their emissions can nevertheless be approximated by using blackbody radiation laws.

Recent example of using blackbody radiation: temperature measurements as covid-19 check



# Blackbody radiation: Planck function

A critical climate feedback

Monochromatic intensity of a blackbody ( $B_\lambda$ )

$$(B_\lambda(T)) = \frac{2hc^2/\lambda^5}{e^{hc/\lambda kT} - 1} \quad (\text{W m}^{-2} \text{ sr}^{-1} \mu\text{m}^{-1})$$

$k$  = Boltzmann's constant,  $1.380649 \times 10^{-23} \text{ J K}^{-1}$

$h$  = Planck's constant,  $6.63 \times 10^{-34} \text{ J s}$

$c$  = speed of light,  $299,792,458 \text{ m s}^{-1}$

$T$  = Temperature of blackbody (K)

sr = solid angle (steradian)

$\lambda$  = wavelength

The Planck function states that the emission intensity of a blackbody depends on the body's temperature and the wavelength considered.




Fig. 4.6: Emission spectra of a blackbody at three different temperatures

## Peak wavelength

Obtained by differentiating  $B_\lambda(T)$  and setting equal to 0:


$$\frac{dB_\lambda(T)}{d\lambda} = 0 \quad \text{for } \lambda = \lambda_{\max} \rightarrow T = \text{const.} = 2.897 \times 10^{-3} \text{ (m K)}$$

$\lambda_{\max}$  = wavelength of maximum intensity

## Wien's displacement law

$$\lambda_{\max} (\mu\text{m}) = \frac{2897}{T}$$

With Wien's displacement law, we can calculate the temperature of a body, if we know its emission spectrum.



Integration over all wavelengths

$$B(T) = \int_0^{\infty} B_{\lambda}(T) d\lambda \sim T^4$$


Integration over all angles of a hemisphere covering a horizontal surface ( $B(T)$  is independent of direction): **total flux density (irradiance)  $F_B$**

$$F_B(T) = \pi B(T) = \pi \int B_{\lambda}(T) d\lambda = \sigma T^4$$


$\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$  Stefan Boltzmann constant

Knowing the flux density  $F_B$  of a blackbody, we can calculate the equivalent blackbody temperature or effective emission temperature ( $T_E$ ).  $T_E$  is the temperature a blackbody would need to emit radiation at, at the measured rate  $F$ . If the body is a blackbody then  $T_E$  and  $T$  of the body are equal.

In the exercises, you will calculate the equivalent blackbody temperature of Earth.



**Fig. 4.8** Radiation balance of the Earth. Parallel beam solar radiation incident on the Earth's orbit, indicated by the thin red arrows, is intercepted over an area  $\pi R_E^2$  and outgoing (blackbody) terrestrial radiation, indicated by the wide red arrows, is emitted over the area  $4\pi R_E^2$ .



Sphere with radius  $r$  has a surface area  $4\pi r^2$

So the flux density received by Earth is:  $4 \times 10^{26} / 4\pi r^2 \rightarrow$  **solar constant**


The solar constant is defined as the rate at which solar energy is received by a surface oriented perpendicular to the Sun's direction at the distance of the Earth's orbit. It is measured to be  $1370 \text{ W m}^{-2}$ .

$$S_0 = 1370 \text{ W m}^{-2}$$

Is it constant?

- Different for each planet.
- Earth's orbit is eccentric ( $r$  varies by  $\pm 1.75\%$ ). Closer to the sun in January, farthest in July. The constant is defined for the average distance.
- It varies as the sun rotates (29-day cycle), bringing sunspots groups across the Earth-facing side of the Sun.
- It varies by  $0.1\%$  over the 11-year solar cycle, with a maximum at sunspot maximum (next 2025).
- It increases as the Sun ages ( $1\%$  every 100 million years).


# Earth's Energy Balance




Solar constant is  $1370 \text{ W m}^{-2}$

Earth receives on average  $340 \text{ W m}^{-2}$

→ See exercises



# Transfer of solar radiation through the atmosphere




There is less radiation reaching the Earth's surface compared to the top of the atmosphere, meaning there must be interaction between solar radiation and Earth's atmosphere.

# Interaction of radiation with atmospheric constituents (gases, particles)


## Definitions

1. **Absorption:** A photon is destroyed. Its energy is distributed by collision into other forms of energy (vibration, rotation...). The frequency of subsequently emitted radiation is independent of the frequency of the absorbed photon.
2. **Scattering/Reflection:** The photon changes direction but not frequency.
3. **Emission:** A photon is created. The molecule emitting the photon loses internal energy (vibrational, rotational...) equivalent to the energy of the photon.



Processes have different importance:


Shortwave: Absorption + Scattering (mostly gases, particles, clouds)  
Longwave: Absorption + Emission (mostly greenhouse gases, clouds)



Absorption continua in **x-ray and UV range** caused by:

- Photoionization (remove electrons from atoms, extreme ultraviolet  $\lambda \leq 0.1 \mu\text{m}$ , happens in ionosphere, i.e upper part of thermosphere)
- Photodissociation (break molecules, ultraviolet  $\lambda \leq 0.31 \mu\text{m}$ , happens down to stratosphere,  $\text{O}_2$  breakup important for  $\text{O}_3$  production)

→ energy is converted into kinetic energy (temperature increase of gas)





Depth of penetration of solar UV radiation (Fig. 4.20, Wallace and Hobbs, 2006)

**Depletion of UV radiation in atmosphere, important for life!**


# Absorption by gases

Absorption lines, mostly in **visible and infrared** spectrum, caused by a change in internal energy state of a gas molecule.





More going on.

# Earth's Energy Balance




Reflection is also important.


## Spectrum of Solar Radiation (Earth)



# Absorption by gases




# At which wavelengths is the atmosphere mostly transparent?



Fundamental consideration:


- 1. Temperature** is a measure of the average energy of **molecular motion** in matter: to and fro translation, intramolecular vibration, and rotation. The sum of these motions' energies can be described as the “thermal energy”. Thermal energy and, hence, **temperature can change as** various forms of energy, including electromagnetic energy, interact with the sample and **change the average energy of motion**.
- 2. Molecular vibrations** (and some energetic rotations) have energy level spacings that **correspond to energies in the IR region** of the electromagnetic spectrum. Thus IR radiation absorbed by molecules causes **increased vibration**.
- 3. Collisions** between these energized molecules in the atmosphere transfer energy among all the molecules, which increases the average thermal energy and, hence, **raises the temperature**.

4. For molecular vibrations to absorb IR energy, the vibrational motions must change the dipole moment of the molecule. All molecules with three or more atoms meet this criterion and are IR absorbers.
5. While the Earth's (dry) atmosphere is predominantly composed of non-IR absorbers, N<sub>2</sub> (78%), O<sub>2</sub> (21%), and Ar (~0.9%), **the 0.1% of remaining trace gases contains many species that absorb IR.**
6. Earth's atmosphere is not dry and can contain several % of water vapor. **Water vapor is the main greenhouse gas.**
7. The presence of "natural" (i.e. not influenced by human action) water vapor, CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and other greenhouse gases are **responsible for the natural greenhouse effect.**



**Dipoles** means that a molecule has negative and positive charges, and the averaged centers of the charges do not co-incide. This leads to a dipole moment, capable of IR absorption. O<sub>2</sub> and N<sub>2</sub> are not dipoles.

# Atmospheric «windows»




The window is between 8-9 and 10 - 12 μm, roughly.

The result of the presence of greenhouse gases is that only a small part of the longwave surface emission makes it to space.


Different greenhouse gases work together to close “atmospheric windows” in the longwave part of the spectrum. They are located at the window edges.

Ozone absorbs at 9.6 μm.







1. Solar radiation is transmitted through Earth's nearly transparent atmosphere (transparent to shortwave radiation).
2. Earth's surface absorbs the solar radiation and warms.
3. The Earth emits thermal IR radiation (blackbody radiation).
4. The atmosphere is much less transparent to thermal infrared radiation and absorbs it.
5. The absorbed radiation leads to warming of the atmosphere, which in turn emits thermal IR radiation in all directions, but also partly downwards.
6. So the net thermal IR flux from the Earth (as blackbody) to space is greatly reduced. This diminishes the radiative cooling of the Earth's surface and leads to surface warming.



# Earth's Energy Balance



Net atmospheric loss of energy.



Climate model from the 60s (**Manabe** and Strickler (1964)


- Longwave & shortwave
- Average Earth cloudiness assumption
- 3 greenhouse gases

**Troposphere:** Cooling due to H<sub>2</sub>O, mainly, and CO<sub>2</sub>. Effect of H<sub>2</sub>O decreases with height along with smaller mixing ratio. Longwave CO<sub>2</sub> effect and shortwave H<sub>2</sub>O nearly cancel each other.


**Stratosphere:** Equilibrium due to warming by O<sub>3</sub> (UV absorption, and a little bit of IR absorption at 9.6  $\mu\text{m}$ ) and cooling by H<sub>2</sub>O and CO<sub>2</sub> (dominant cooling agent through longwave emission). Model assumes equilibrium in the stratosphere.

# Convective - radiative equilibrium

Further aloft in the troposphere, the re-emission of IR radiation by infrared-active gases and clouds leads to a net energy loss of the atmosphere.



Because the density of the atmosphere increases close to the surface, the layers become optically thicker and physically thinner. Hence, removing heat near the surface becomes difficult. If only radiative equilibrium were responsible for the energy balance, Earth's surface would be very hot!




In fact, convective transport of sensible and latent heat of water vapor cools the surface and warms the troposphere and leads to the observed lapse rate of -6.5 K / km.

equil. = equilibrium  
ajd. = adjustment

■ Catling, D., & Kasting, J. (2017). Principles of Planetary Atmospheres. In *Atmospheric Evolution on Inhabited and Lifeless Worlds* (pp. 1-168). Cambridge: Cambridge University Press. Wallace and Hobbs, 2006, Ch. 4.3.6

Greenhouse gases reduce the emission by Earth at the top of the atmosphere!



In this graph, the radiation emitted by the Earth's surface retrieved at the TOA (e.g. on a satellite) is shown. It is about 40 % less than emitted at the surface. The grey lines show blackbody emissions for different temperatures.

Where the "measured" spectrum is close to the curve of a high blackbody temperature, atmospheric "windows" are present.

Absorption bands show low brightness temperatures, because the maximum absorption (extinction) happens further up in the atmosphere, where it is colder.

The greenhouse effect is the difference between the blue and ~300 K curve.


# Summary of greenhouse effect processes

- In summary the greenhouse effect induced by infrared-active gases consists of three elements:
  1. Downwelling irradiance from the atmosphere to the surface, heating the surface and elevating the surface temperature;
  2. A decrease in upwelling irradiance at the top of the atmosphere; and
  3. Radiative cooling (and heating) of the atmosphere, affecting the vertical temperature structure of the atmosphere.


# Stratospheric cooling

- The stratosphere has been cooling as the troposphere is warming.
- Contributing effects are:
  - Depletion of the ozone layer. Ozone is a greenhouse gas. Less ozone means less IR absorption.
  - More CO<sub>2</sub> in the troposphere means that the flux of 15 μm IR to the stratosphere is diminished. So stratospheric CO<sub>2</sub> absorbs less IR while emitting the same amount.

Global Mean Temperature Changes in Model and Observed SSU and MSU Data



# Atmospheric absorption spectra



doi:10.5194/esd-7-697-2016

# What about the effect of particles and clouds?

