



Microorganisms in Diverse Environments

Diversity of growth conditions for microorganism:


- Temperature
- O₂ availability
- Light availability
- pH
- Water availability
- Nutrient availability
- Toxic contaminants



Gradients and extremes

Temperature Classes of Microorganisms


- Cardinal Temperatures
 - **Minimum, optimum, and maximum** temperatures at which an organism grows
 - Characteristic of any given microorganism
 - Differ dramatically between species
 - At optimum, all/most cellular components are functioning at maximum rate



Temperature Classes of Microorganisms

Broad classes related to growth temperature optima

- **Psychrophile:** low, found in cold environments
- **Psychrotroph:** low-mid can grow at cold temperatures
- **Mesophile:** midrange, most commonly studied
- **Thermophile:** high, found in hot environments
- **Hyperthermophile:** very high, found in extremely hot habitats such as hot springs and deep-sea hydrothermal vents



Extremophiles




Extremophiles




Microbial Life in the Cold

Cold Environments:

- Much of Earth's surface is cold
 - Oceans: 5°C
 - Arctic and Antarctic are permanently frozen or rarely unfrozen
- **Constantly** vs. **seasonally** cold



Antarctic Microbial Habitats and Microorganisms



(a) A core of frozen seawater from McMurdo, coloration due to pigmented microorganisms

(b) Phototrophic microorganisms within the the core, most are diatoms or green algae

(c) *Polaromonas*, a bacterium that lives in sea ice with optimal growth at 4°C



(d) Permanently ice-covered lake, Antarctica. The water column under the ice contains a diverse array of *Bacteria*, *Archaea*, and microbial eukaryotes


(e) Glaciers and subglacial lakes are full with of microbial life

Microbial Life in the Cold


- Psychrophilic and Psychrotolerant Microorganisms
 - **Psychrophiles:** optimal growth temperature $\leq 15^{\circ}\text{C}$, maximum $< 20^{\circ}\text{C}$, minimum $\leq 0^{\circ}\text{C}$
 - Constantly cold environments
 - Found in polar regions, permanent snowfields, glaciers
 - **Psychrotolerant:** can grow at 0°C but have optima of $20^{\circ}\text{C} - 40^{\circ}\text{C}$
 - More widely distributed in nature than psychrophiles
 - Isolated from soils and water in temperate climates and food at 4°C

Microbial Life in the Cold


- Psychrophilic algae and bacteria often grow within and under sea ice in polar regions
- They can also be found on the surfaces of permanent snowfields and glaciers
- This alga grows within the snow as a green-pigmented vegetative cell and then sporulates



- As the snow melts the spores become concentrated on the surface



Microbial Life in the Cold



Planococcus halocryophilus

Isolated from high Arctic permafrost, grows and divides at $-15\text{ }^{\circ}\text{C}$ and is metabolically active at $-25\text{ }^{\circ}\text{C}$

Microbial Life in the Cold





Cryptoendolith

Psychrotolerant microorganisms are more widely distributed in nature than are psychrophiles






Food preservation temperatures.






Molecular Adaptations to Life in the Cold

- Production of enzymes that function optimally in the cold
 - more α -helices than β -sheets \rightarrow greater flexibility for catalysis at cold temperatures
 - more polar and fewer hydrophobic amino acids




Molecular Adaptations to Life in the Cold

- Cytoplasmic membranes function at low temperatures.
 - higher unsaturated and shorter-chain fatty acid content
 - some polyunsaturated fatty acids, which remain flexible at very low temperatures





Molecular Adaptations to Life in the Cold

- Cold shock proteins (chaperones)




Molecular Adaptations to Life in the Cold

- Cryoprotectants - prevent formation of ice crystals
 - e.g., antifreeze proteins, certain solutes (glycerol, sugars, ethanol)
- Exopolysaccharide cell surface slime
 - extracellular carbohydrate polymers



Molecular Adaptations to Life in the Cold


- Exopolysaccharide cell surface slime
 - extracellular carbohydrate polymers



Microbial Life at High Temperatures

- Thermal Environments

- **Thermophiles:** growth temperature optima between 45°C and 80°C
- **Hyperthermophiles:** optima greater than 80°C
 - inhabit hot environments, including boiling hot springs and seafloor hydrothermal vents, that can experience temperatures in excess of 100°C



Microbial Life at High Temperatures

- Hyperthermophiles and Thermophiles
 - Thermophiles - optima 45–80°C
 - moderately or intermittently hot environments
 - Hyperthermophiles
 - inhabit boiling hot springs
 - chemoorganotrophic and chemolithotrophic species present
 - Fast generation times (can be 1h)
 - high prokaryotic diversity (both **Archaea** and **Bacteria** represented)



(a)

superheated




(b)

Hot Spring Microbes and Their Heat-Stable Enzymes



Nancy L. Spear

(a)



(b)


Presently Known Upper Temperature Limits for Growth of Living Organisms

Group	Upper temperature limits (°C)	
Macroorganisms		
Animals		
Fish and other aquatic vertebrates	38	
Insects	45–50	
Ostracods (crustaceans)	49–50	
Plants		
Vascular plants	45 (60 for one species)	
Mosses	50	
Microorganisms		
Eukaryotic microorganisms		
Protozoa	56	
Algae	55–60	
Fungi	60–62	
Bacteria and Archaea		
Bacteria		
Cyanobacteria	73	Above 65°C, only prokaryotic life forms thrive
Anoxygenic phototrophs	70–73	
Chemoorganotrophs/chemolithotrophs	95	
Archaea		
Chemoorganotrophs/chemolithotrophs	122	extensive diversity

Presently Known Upper Temperature Limits for Growth of Living Organisms



Back smokers emit hydrothermal vent fluid at 250–350°C or higher



Methanopyrus: most thermophilic, up to 122°C



Microbial Life at High Temperatures

Protein and Membrane Stability at High Temperatures

- Enzymes and proteins more heat-stable and function optimally at high temperatures
 - AA changes compared to mesophiles
 - Increased ionic bonding and highly hydrophobic interiors
- Thermophilic and hyperthermophilic enzymes commercially useful
 - Prolong shelf life (stability)
 - **e.g., Taq polymerase** for polymerase chain reaction [PCR]

Microbial Life at High Temperatures

- Heat naturally works to peel apart the lipid bilayer
- Cytoplasmic membranes must be heat stable
 - **Bacteria** have lipids rich in long-chain and saturated fatty acids, fewer unsaturated fatty acids
 - Most hyperthermophiles (**Archaea**) have C₄₀ hydrocarbons made of repeating isoprene units
 - membrane forms lipid monolayer rather than bilayer.







Oxygen and Microbial Growth

Microorganisms can be grouped according to their relationship with O₂

Oxygen Classes of Microorganisms:

- **Aerobes:** grow at full O₂ tension (~21%) and respire O₂
- **Microaerophiles:** can use O₂ only at levels reduced from that in air (microoxic) due to limited respiration or oxygen sensitivity
- **Facultative** organisms: can live with or without oxygen



Mycobacterium tuberculosis

Helicobacter pylori

Saccharomyces cerevisiae


Oxygen and Microbial Growth

Microorganisms can be grouped according to their relationship with O₂

Oxygen Classes of Microorganisms:

- **Anaerobes:** cannot respire oxygen
- **Aerotolerant anaerobes:** tolerate oxygen and grow in its presence even though they cannot respire
- **Obligate anaerobes:** inhibited or killed by oxygen, (e.g., Bacteria and Archaea, few fungi, and few protozoa)

Anoxic (oxygen free) habitats: mud, bogs, marshes, animal intestines, etc...



Oxygen Relationships of Microorganisms

Aerobes

Group	Relationship to O ₂	Type of metabolism	Example ^a	Habitat ^b
Obligate	Required	Aerobic respiration	Micrococcus luteus (B)	Skin, dust
Facultative	Not required, but growth better with O ₂	Aerobic respiration, anaerobic respiration, fermentation	Escherichia coli (B)	Mammalian large intestine
Microaerophilic	Required but at levels lower than atmospheric	Aerobic respiration	Spirillum volutans (B)	Lake water

Anaerobes


Group	Relationship to O ₂	Type of metabolism	Example ^a	Habitat ^b
Aerotolerant	Not required, and growth no better when O ₂ present	Fermentation	Streptococcus mutans (B)	Oral cavity
Obligate	Harmful or lethal	Fermentation or anaerobic respiration	Methanobacterium formicicum (A)	Sewage sludge, anoxic lake sediments

^aLetters in parentheses indicate phylogenetic status (B, **Bacteria**; A, **Archaea**). Representatives of either domain of prokaryotic cells are known in each category. Most eukaryotes are obligate aerobes, but facultative aerobes (for example, yeast) and obligate anaerobes (for example, certain protozoa and fungi) are known.

^bListed are typical habitats of the example organism; many others could be listed.



Culture Techniques for Aerobes and Anaerobes

- Aerobes need extensive aeration (e.g., shaking, bubbling).




Culture Techniques for Aerobes and Anaerobes

- Anaerobes need oxygen to be excluded
 - **reducing agents:** added to culture media to reduce oxygen to water
 - Flush or consume oxygen (e.g., glove bag)



(b)


Coy Laboratory Products



(a)

Deborah O. Jung and M.T. Madigan

Culture Techniques for Aerobes and Anaerobes



Four-electron reduction of O₂ to H₂O by stepwise addition of electrons

Why is oxygen toxic?

Why are anaerobic microorganisms inhibited in their growth or even killed by oxygen?

Reactants	Products
$O_2 + e^- \rightarrow$	O_2^- (superoxide)
$O_2^- + e^- + 2 H^+ \rightarrow$	H_2O_2 (hydrogen peroxide)
$H_2O_2 + e^- + H^+ \rightarrow$	$H_2O + OH^\bullet$ (hydroxyl radical)
$OH^\bullet + e^- + H^+ \rightarrow$	H_2O (water)


Outcome:



- Molecular oxygen (O₂) is not toxic.
- Exposure to oxygen yields toxic byproducts

- **superoxide anion** (O₂⁻)
- **hydrogen peroxide** (H₂O₂)
- **hydroxyl radical** (OH[•])

Reactive Oxygen Species (ROS)
by-products of the reduction of O₂ to H₂O in respiration. Electron carriers found in virtually all cells can catalyse these reactions.



Superoxide Dismutase and Other Enzymes That Destroy Toxic Oxygen

Keeping toxic oxygen molecules under control




Universal across life in O₂ environments



Superoxide Dismutase and Other Enzymes That Destroy Toxic Oxygen

Enzymes destroy O₂⁻ and H₂O₂


- Catalase and peroxidase convert H₂O₂ to O₂ and H₂O
- Superoxide dismutase converts 2 O₂⁻ to H₂O₂ and O₂
- **Superoxide reductase** in some strict anaerobes converts O₂⁻ to H₂O₂ without producing O₂




(a) **Catalase**




(b) **Peroxidase**



(c) **Superoxide dismutase**




(d) **Superoxide dismutase/catalase in combination**



(e) **Superoxide reductase**

Presence of Catalase Test




T.D. Brock



Effects of pH on Microbial Growth

- pH expresses acidity or alkalinity of a solution
- pH 7 = neutral
- **acidic** pH < 7
- **alkaline** pH > 7
- Each microbe has a pH range ~2–3 pH units within which growth is possible
- Each organism shows a well-defined pH optimum
- Most natural environments are pH 3–9



Relationships of Microorganisms to pH


Physiological class (optima range)	Approximate pH optimum for growth	Example organism ^a
Neutrophile (pH > 5.5 and < 8)	7	<i>Escherichia coli</i>
Acidophile (pH < 5.5)	5	<i>Rhodospila globiformis</i>
	3	<i>Acidithiobacillus ferrooxidans</i>
	1	<i>Picrophilus oshimae</i>
Alkaliphile (pH ≥ 8)	8	<i>Chloroflexus aurantiacus</i>
	9	<i>Bacillus firmus</i>
	10	<i>Natronobacterium gregoryi</i>

Although some microorganisms can live at very low or very high pH, the cell's internal pH remains near neutrality

^a*Picrophilus* and *Natronobacterium* are Archaea; all others are Bacteria.

Acidophiles


- Grow best at low pH (< 5.5)
- Different classes growing best at moderately acidic or very low pH
 - Many fungi and bacteria grow best at pH 5
 - Restricted number grow best below pH 3
 - Few grow best below pH 2
 - pH optima below 1 are extremely rare
- Most acidophiles cannot grow at pH 7 and many cannot grow at pH values more than two units above their optimum



Ferroplasma acidiphilum

Acidophiles

- Governed by stability of cytoplasmic membrane
- At neutral pH, membranes of strong acidophiles lyse; protons required for stability



Isolated from dry, extremely acidic soil (pH < 0.5) that was heated by solfataric gases to about 55C

Picrophilus

grows optimally at pH 0.7 and 60°C


- One of the smallest genomes
- High coding density in the genome
- Unknown DNA repair mechanism
- Large number of transport proteins:
transmembrane proton gradient for transport

permanent huge Δ pH
exists across the
membrane of acidophiles

Acidophiles & Alkaliphiles

- Cytoplasmic pH
 - Optimal pH refers to extracellular only
 - The **intracellular** pH must stay relatively close to neutral (pH 5–9), consistent with macromolecule stability
 - Extreme acidophiles maintain cytoplasmic pH near neutrality
- Highly impermeable cell membranes to prevent proton (H^+) influx
- Active proton pumps to export any entering protons
- Cation transporters (K^+ uptake) to maintain electrochemical balance and create a positive membrane potential that repels H^+

Highly impermeable cell membranes of Acidophiles




Article | [Open access](#) | Published: 27 November 2012

A proton shelter inspired by the sugar coating of acidophilic archaea

[Xiumei Wang](#), [Bei'er Lv](#), [Guixin Cai](#), [Long Fu](#), [Yuanzi Wu](#), [Xiang Wang](#), [Bin Ren](#) & [Hongwei Ma](#)

[Scientific Reports](#) 2, Article number: 892 (2012) | [Cite this article](#)



Alkaliphiles

- High pH optima for growth, $\text{pH} \geq 8$
 - sometimes as high as $\text{pH} 10$
- Found in highly alkaline habitats: soda lakes and high-carbonate soils
- Used commercially (**e.g.**, secreted proteases and lipases that are added to laundry detergents to remove proteins and fats)
- Some have sodium (Na^+) motive force rather than proton motive force.

Bacillus firmus unusually broad range for growth, from $\text{pH} 7.5$ to 11

Natronobacterium gregoryi 20% and optimum pH of 10



Osmolarity and Microbial Growth




Water availability

Water is the solvent of life, and water availability is an important factor affecting the growth of microorganisms.




Osmolarity and Microbial Growth

- Water availability depends on:
 - environmental moisture/dryness
 - concentration of solutes – bind water
- Water availability: Water activity (a_w)
 - varies from zero (no free water) to one (pure water)
- **Osmosis:** Water diffuses from high to low concentrations.
 - Cell cytoplasm typically have higher solute concentration than environment




Cell in isotonic solution.



Under these conditions, the solute concentration in the cell is equivalent to a solute concentration of 0.85% sodium chloride (NaCl).

Plasmolyzed cell in hypertonic solution.



If the concentration of solutes such as NaCl is higher in the surrounding medium than in the cell (the environment is hypertonic), water tends to leave the cell. Growth of the cell is inhibited.


Water Activity of Substances


Water activity (a_w)	Material	Example organisms ^a
1.000	Pure water	Caulobacter, Spirillum
0.995	Human blood	Streptococcus, Escherichia
0.980	Seawater	Pseudomonas, Vibrio
0.950	Bread	Most gram-positive rods
0.900	Maple syrup, ham	Gram-positive cocci such as Staphylococcus
0.850	Salami	Saccharomyces rouxii (yeast)
0.800	Fruit cake, jams	Zygosaccharomyces bailii (yeast), Penicillium (fungus)
0.750	Salt lakes, salted fish	Halobacterium, Halococcus
0.700	Cereals, candy, dried fruit	Xeromyces bisporus and other xerophilic fungi

^aSelected examples of **Bacteria** and **Archaea** or fungi capable of growth in culture media adjusted to the stated water activity.

Halophiles

- Seawater contains ~3% NaCl
 - Freshwater is <1%
- **Halophiles:** grow best at $a_w = 0.98$ (seawater); have a specific requirement for NaCl
- **Halotolerant:** tolerate some dissolved solutes but generally grow best in the absence of added solute
- **Extreme halophiles:** require very high levels (15-30%) of NaCl; often unable to grow at lower concentrations






Aliivibrio fischeri

Related organisms

- Organisms able to live in environments high in sugar: Osmophiles
- Organisms able to live in very dry environments: Xerophiles



Wallemia sebi



Zygosaccharomyces rouxii




Compatible Solutes of Microorganisms


- Maintenance of positive water balance
- increase of internal solute concentration
- Pumping solutes into the cell from the environment
- Synthesizing cytoplasmic solutes
 - Must not inhibit biochemical pathways of the cell

Compatible solutes:

- highly water-soluble organic molecules
- sugars, alcohols, and amino acid derivatives







Glycine



Glycine betaine

Compatible Solutes of Microorganisms

Organism group and example	Major cytoplasmic compatible solute(s)	Minimum a_w for growth ^c
Most nonphototrophic Bacteria (<i>Escherichia</i>) and freshwater cyanobacteria (<i>Anabaena</i>)	Amino acids (mainly glutamate or proline ^a)/sucrose, trehalose ^b	0.98 
Marine cyanobacteria (<i>Synechococcus</i>)	α -Glucosylglycerol ^b	0.92
Marine algae (<i>Phaeocystis</i>)	Mannitol, ^b various glycosides, dimethylsulfoniopropionate	0.92 
Halotolerant Bacteria (<i>Staphylococcus</i>)	Amino acids	0.90
Salt lake cyanobacteria (<i>Aphanothece</i>)	Glycine betaine	0.75 
Halophilic phototrophic purple Bacteria (<i>Halorhodospira</i>)	Glycine betaine, ectoine, trehalose ^b	0.75 

Organism group and example	Major cytoplasmic compatible solute(s)	Minimum a_w for growth ^c
Extremely halophilic Archaea (Halobacterium) and some Bacteria (Salinibacter)	KCl	0.75
Halophilic green algae (Dunaliella)	Glycerol	0.75 <div style="text-align: center; margin-top: 10px;"> $\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CHOH} \\ \\ \text{CH}_2\text{OH} \\ \text{Glycerol} \end{array}$ </div>
Haloalkaliphilic Archaea (Natrinema)	KCl	0.68
Xerophilic and osmophilic yeasts (Zygosaccharomyces)	Glycerol	0.62
Xerophilic filamentous fungi (Xeromyces)	Glycerol	0.605

Why study extremophiles?

