

**Exercise 1** (Ex 1.1 page 7).

Let

$$F(x, y, z) = (y^2 \sin(xz), e^y \cos(x^2 + z), \ln(2 + \cos(xy))) = (F_1, F_2, F_3).$$

Compute:

1.  $\text{grad } F_1, \text{grad } F_2, \text{grad } F_3$
2.  $\text{div } F$
3.  $\text{rot } F$ .

**Exercise 2** (Ex 1.2 page 7).

Which of the following expressions are correct in the case that  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  is of class  $C^1(\mathbb{R}^3)$  and  $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is of class  $C^1(\mathbb{R}^3; \mathbb{R}^3)$ ?

- a)  $\text{grad } f$     b)  $f \text{ grad } f$     c)  $F \cdot \text{grad } f$     d)  $\text{div } f$   
 e)  $\text{div}(fF)$     f)  $\text{rot}(fF)$     g)  $\text{rot } f$     h)  $f \text{ rot } F$   
 i)  $\text{rot div } F$ .

**Exercise 3** (Ex 1.3 page 5).

Let  $x = (x_1, \dots, x_n)$ ,  $a = (a_1, \dots, a_n)$ , and  $r$  such that  $r = \sqrt{\sum_{i=1}^n (x_i - a_i)^2}$ . Let  $f$  be a scalar field defined by  $f(x) = 1/r$ . Compute  $\Delta f$ .

**Exercise 4** (Ex 1.6 and 1.7 page 8).

Let  $\Omega \subset \mathbb{R}^3$  be an open domain. Verify that:

1. If  $f \in C^1(\Omega)$  and  $g \in C^2(\Omega)$ , then:

$$\text{div}(f \text{ grad } g) = f \Delta g + \text{grad } f \cdot \text{grad } g$$

2. If  $f, g \in C^1(\Omega)$ , then:

$$\text{grad}(fg) = f \text{ grad } g + g \text{ grad } f$$

3. If  $f \in C^1(\Omega)$  and  $F \in C^1(\Omega, \mathbb{R}^3)$  then:

$$\text{div}(fF) = f \text{ div } F + F \cdot \text{grad } f$$

4. If  $F \in C^2(\Omega, \mathbb{R}^3)$ , then:

$$\operatorname{rot} \operatorname{rot} F = -\Delta F + \operatorname{grad} \operatorname{div} F,$$

where  $\Delta F = (\Delta F_1, \Delta F_2, \Delta F_3)$  for a given vector field  $F = (F_1, F_2, F_3)$ .

5. If  $f \in C^1(\Omega)$  and  $F \in C^1(\Omega, \mathbb{R}^3)$ , then:

$$\operatorname{rot}(fF) = \operatorname{grad} f \times F + f \operatorname{rot} F$$

**Exercise 5** (Ex 1.4 page 7).

Let  $f \in C^2(\Omega)$ , where:

$$\Omega = \{(x, y) \in \mathbb{R}^2 : x, y > 0\}.$$

1. In the case where we have:

$$g(r, \theta) := f(r \cos \theta, r \sin \theta) = f(x, y),$$

show that:

$$\frac{\partial^2 g(r, \theta)}{\partial r^2} + \frac{1}{r} \frac{\partial g(r, \theta)}{\partial r} + \frac{1}{r^2} \frac{\partial^2 g(r, \theta)}{\partial \theta^2} = \frac{\partial^2 f(x, y)}{\partial x^2} + \frac{\partial^2 f(x, y)}{\partial y^2} = \Delta f(x, y).$$

2. Compute  $\Delta f$  when:

$$f(x, y) := \sqrt{x^2 + y^2} + \left( \arctan \frac{y}{x} \right)^2.$$