

TABLE DE TRANSFORMÉES DE FOURIER

	$f(y)$	$\mathfrak{F}(f)(\alpha) = \hat{f}(\alpha)$
1	$f(y) = \begin{cases} 1 & \text{si } y < b \\ 0 & \text{sinon} \end{cases}$	$\hat{f}(\alpha) = \sqrt{\frac{2}{\pi}} \frac{\sin(b \alpha)}{\alpha}$
2	$f(y) = \begin{cases} 1 & \text{si } b < y < c \\ 0 & \text{sinon} \end{cases}$	$\hat{f}(\alpha) = \frac{e^{-ib\alpha} - e^{-ic\alpha}}{i\alpha\sqrt{2\pi}}$
3	$f(y) = \begin{cases} e^{-\omega y} & \text{si } y > 0 \\ 0 & \text{sinon} \end{cases} \quad (\omega > 0)$	$\hat{f}(\alpha) = \frac{1}{\sqrt{2\pi}} \frac{1}{\omega + i\alpha}$
4	$f(y) = \begin{cases} e^{-\omega y} & \text{si } b < y < c \\ 0 & \text{sinon} \end{cases}$	$\hat{f}(\alpha) = \frac{1}{\sqrt{2\pi}} \frac{e^{-(\omega+i\alpha)b} - e^{-(\omega+i\alpha)c}}{(\omega + i\alpha)}$
5	$f(y) = \begin{cases} e^{-i\omega y} & \text{si } b < y < c \\ 0 & \text{sinon} \end{cases}$	$\hat{f}(\alpha) = \frac{1}{i\sqrt{2\pi}} \frac{e^{-i(\omega+\alpha)b} - e^{-i(\omega+\alpha)c}}{\omega + \alpha}$
6	$f(y) = \frac{1}{y^2 + \omega^2} \quad (\omega \neq 0)$	$\hat{f}(\alpha) = \sqrt{\frac{\pi}{2}} \frac{e^{- \omega\alpha }}{ \omega }$
7	$f(y) = \frac{e^{- \omega y }}{ \omega } \quad (\omega \neq 0)$	$\hat{f}(\alpha) = \sqrt{\frac{2}{\pi}} \frac{1}{\omega^2 + \alpha^2}$
8	$f(y) = e^{-\omega^2 y^2} \quad (\omega \neq 0)$	$\hat{f}(\alpha) = \frac{1}{\sqrt{2} \omega } e^{-\frac{\alpha^2}{4\omega^2}}$
9	$f(y) = ye^{-\omega^2 y^2} \quad (\omega \neq 0)$	$\hat{f}(\alpha) = \frac{-i\alpha}{2\sqrt{2} \omega ^3} e^{-\frac{\alpha^2}{4\omega^2}}$
10	$f(y) = \frac{4y^2}{(\omega^2 + y^2)^2} \quad (\omega \neq 0)$	$\hat{f} = \sqrt{2\pi} \left(\frac{1}{ \omega } - \alpha \right) e^{- \omega\alpha }$