


## GROUP THEORY 2024 - 25, EXERCISE SHEET 2

**Exercise 1.** (easy) *Warm up exercise*

Consider the natural action of  $S_4$  on the set  $X$  of all 2-elements subsets of  $\{1, 2, 3, 4\}$ . This action can be seen as permuting the edges of the following diagram



- (1) Determine the orbit of  $\{1, 2\} \in X$ ;
- (2) Find the stabilizer of  $\{1, 2\} \in X$ ;
- (3) Verify that the Orbit-Stabilizer theorem holds by recovering  $|S_4|$  from the previous points.

**Exercise 2.** (medium) *Action on cosets*

Let  $G$  be a group and  $H \leq G$  a subgroup. We define an action of  $G$  on  $G/H$  by

$$\begin{aligned} G \times G/H &\rightarrow G/H \\ (g, g'H) &\mapsto gg'H \end{aligned}$$

- (1) Show that the above operation is well-defined and that it indeed defines a  $G$ -action;
- (2) For  $gH \in G/H$ , find the stabilizer  $\text{Stab}_G(gH)$ ;

If  $X, Y$  are sets endowed with  $G$ -actions, we say that  $X$  and  $Y$  are isomorphic as  $G$ -sets if there exists a bijective function  $f : X \rightarrow Y$  such that  $f(g \cdot x) = g \cdot f(x)$  for all  $g \in G$  and  $x \in X$ .

**Exercise 3.** (medium) Let  $H$  and  $K$  be subgroups of a group  $G$ . Define the action of  $G$  on the sets  $G/H$  and  $G/K$  by left multiplication (see Exercise 2). Show that  $G/H$  and  $G/K$  are isomorphic as  $G$ -sets if and only if  $H$  and  $K$  are conjugate subgroups of  $G$ , i.e. there exists  $g \in G$  such that  $gHg^{-1} = K$ .

**Exercise 4.** (medium) Let  $V = \mathbb{F}_2^3$  and  $G = GL(V)$  the group of linear automorphisms of  $V$  (its elements are bijective linear maps  $V \rightarrow V$ ).

- (1) Define a natural action of  $G$  on

$$X = \{W \subset V \mid \dim(W) = 2\}.$$

- (2) Show the action defined above is transitive;
- (3) Determine the cardinality  $|X|$ .

**Exercise 5.** (medium)

- (1) Let  $G \leq S_n$ . Consider its natural action on  $\Omega = \{1, 2, \dots, n\}$ . Show that if  $G$  acts transitively on  $\Omega$ , then  $n$  divides  $|G|$ .
- (2) Let  $G$  be a group and  $X = \{H \subseteq G\}$  be the set of subgroups of  $G$ . Show that  $G$  acts on  $X$  by

$$G \times X \rightarrow X$$
$$(g, H) \mapsto gHg^{-1}.$$

For  $H \in X$ , what is the stabilizer  $\text{Stab}_G(H)$  ?

**Exercise 6.** (hard) *p*-group actions and fixed points

Use the orbit-stabiliser theorem to prove the following result. Let  $p > 0$  be a prime number and let  $G$  be a group of order  $p^n$  for some  $n \geq 1$  which acts on a set  $X$  with  $p \nmid |X|$ . Show that there exists  $x \in X$  such that  $g \cdot x = x$  for all  $g \in G$ .