

## GROUP THEORY 2024 - 25, SOLUTION SHEET 10

**Exercise 1.** To do yourself. Ask the assistant if something is unclear.

**Exercise 2.** Let  $G/H$  be the set of left co-sets of  $H$  in  $G$ . Then  $|G/H| = p$  and hence there is an induced homomorphism  $\varphi : G \rightarrow S_p$ . Let  $K$  denote the kernel of  $\varphi$  and consider the following two lemmas:

**Lemma 2.1:** The cardinality of  $G/K$  is  $p$ .

**Lemma 2.2:** We have an inclusion of subgroups  $K \subseteq H$ .

Assuming the lemmas, since the index of both  $K$  and  $H$  in  $G$  is  $p$  and  $K \subseteq H$  we can conclude that  $H = K$ . The fact that  $K$  is a kernel of a homomorphism yields that  $H = K$  is a normal subgroup. We leave the proof of Lemma 2.2 to the reader and prove Lemma 2.1.

**Proof of Lemma 2.1:**

Let  $q$  be a prime factor of  $|G/K|$ , then since  $p$  is assumed to be the minimum prime dividing  $|G|$  and  $|G/K| \mid |G|$ , we have that  $q \geq p$ . By the first isomorphism theorem applied to  $\varphi$  we obtain that  $G/K$  is isomorphic to a subgroup of  $S_p$ . Therefore  $q \mid |G/K| \mid p!$  and hence  $q \leq p$ . So we obtain  $q = p$ . So  $|G/K| = p^n$  but  $|G/K| \mid p!$  also implies that  $n = 1$ . Hence  $|G/K| = p$ .  $\square$

**Exercise 3.** (1) By an exercise of a preceding series, a  $p$ -group of order  $n$  has normal subgroups of order  $p^k$  for all  $1 \leq k \leq n$ , which proves the claim.

(2) Without loss of generality suppose that  $p > q$ . By the Sylow theorems, the number  $n_p$  of Sylow  $p$ -subgroups of the group divides  $q$  and has residue 1 modulo  $p$ . As  $p > q$ ,  $n_p$  has to be 1 and by an exercise of series 9 the unique Sylow  $p$ -subgroup is normal.

(3) If  $q < p$ , then the index of a Sylow  $p$ -subgroup  $P$  is equal to  $q$ , the smallest prime that divides the order of the group. By exercise 2,  $P$  is normal in  $G$ . Now suppose that  $p < q$ . It cannot be that case that  $p$  has residue 1 modulo  $q$ , so the number of Sylow  $q$ -subgroups, which we denote by  $n_q$  should obey  $n_q = 1$  or  $n_q = p^2$ . If  $n_q = 1$ , then the unique Sylow  $q$ -subgroup is normal and  $G$  is not simple, so we assume that  $n_q = p^2$ . Since a Sylow  $q$ -subgroup has order  $q$  and two distinct Sylow  $q$ -subgroups intersect trivially (since  $Q \cap Q'$  is a subgroup of  $Q$ , its order must divide the prime  $q$ , hence either  $Q = Q'$  or  $Q \cap Q' = 1$ ),  $G$  has  $p^2(q - 1)$  elements of order  $q$ . Therefore, a Sylow  $p$ -subgroup contains all of the remaining  $p^2$  elements of  $G$ . In this case, we conclude that the Sylow  $p$ -subgroup is unique, so it is normal in  $G$ .

(4) Without loss of generality suppose that  $p < q < r$ . If the number  $n_s$  of Sylow  $s$ -subgroup is 1 for  $s = p, q$  or  $r$ , then the (unique) Sylow  $s$ -subgroup is normal. So we suppose

now that  $n_p, n_q$  and  $n_r$  are all strictly bigger than 1. Using Sylow's theorems, we deduce that  $n_p \geq q, n_q \geq r$  and  $n_r \geq pq$ . Since for  $s = p, q, r$  Sylow  $s$ -subgroups are intersect trivially (as in the preceding point), we can count the elements in those Sylow  $s$ -subgroups (those elements are of order  $s$ ) to find that:

$$\begin{aligned} |G| &\geq n_p(p-1) + n_q(q-1) + n_r(r-1) \\ &\geq q(p-1) + r(q-1) + pq(r-1) \\ &= qp - q + rq - r + pqr - pq = pqr + r(q-1) - q \\ &\geq |G| + q(q-2) \\ &> |G| \end{aligned}$$

a contradiction.

**Exercise 4.** Note that every positive integer less than 60 can be written in one of the following forms:

1.  $p^n$  for a prime  $p$  and  $n \geq 0$ .
2.  $p^a q^b$  for distinct primes  $p, q$  and  $a > 0, b > 0$ .
3.  $pqr$  for distinct primes  $p, q, r$ .

Let  $G$  be a non-abelian group of order  $n$  such that  $n < 60$ . If  $n$  is of the form  $p^n, pqr$  then  $G$  is not simple by exercise 3. If  $n$  is of the form  $p^a q^b$ , then  $G$  is solvable by Burnside's Theorem. But then solvability of  $G$  implies that  $H := [G, G]$  is a normal subgroup of  $G$  with  $G \neq H$ . Since  $G$  is not abelian we also have that  $H$  is not trivial. Hence  $G$  is not simple.

**Exercise 5.** By the Sylow theorems, the number of Sylow 2-subgroups must be either 1 or 3. In the former case, this subgroup is normal and we are done. In the latter case, we have a group homomorphism  $G \rightarrow S_3$  given by the action of  $G$  on the set of Sylow 2-subgroups. If  $G$  is simple, this must be injective, which means that  $G$  has at most 6 elements, which contradicts the hypothesis  $n \geq 2$ .

**Exercise 6.** (1) Let  $\sigma \in \text{Aut}(K)$  such that  $\sigma\varphi_1(L)\sigma^{-1} = \varphi_2(L)$ . Let  $x \in L$  be a generator of the cyclic group  $L$ . Then there exists  $a \in \mathbb{N}$  such that  $\sigma \circ \varphi_1(x) \circ \sigma^{-1} = \varphi_2(x^a) = \varphi_2(x)^a$ . Now for every  $l \in L$  there exists  $b \in \mathbb{N}$  such that  $l = x^b$ , which implies that

$$\begin{aligned} \sigma \circ \varphi_1(l) \circ \sigma^{-1} &= \sigma \circ \varphi_1(x^b) \circ \sigma^{-1} \\ &= (\sigma \circ \varphi_1(x) \circ \sigma^{-1})^b \\ &= (\varphi_2(x)^a)^b \\ &= \varphi_2(x^b)^a \\ &= \varphi_2(l)^a \end{aligned}$$

(1)

as suggested by the hint. We now define

$$\psi : K \rtimes_{\varphi_1} L \rightarrow K \rtimes_{\varphi_2} L$$

by  $\psi(k, l) = (\sigma(k), l^a)$ . We let the reader verify that  $\psi$  is a group homomorphism. To construct an inverse  $\phi : K \rtimes_{\varphi_2} L \rightarrow K \rtimes_{\varphi_1} L$  of  $\psi$ , we just change the role of  $\varphi_1$  and  $\varphi_2$  above. By the same argument, there exists an integer  $b$  such that

$$(2) \quad \sigma^{-1} \circ \varphi_2(l) \circ \sigma = \varphi_1(l)^b.$$

Hence we know that  $\phi : (k, l) \mapsto (\sigma^{-1}(k), l^b)$  is a group homomorphism. Combining the two equations (1) and (2) we obtain that  $\varphi_2(l^{ab}) = \varphi_2(l)$  and  $\varphi_1(l^{ab}) = \varphi_1(l)$ . It is now straightforward to check that  $\phi$  and  $\psi$  are inverses of each other.

(2) Let  $\varphi_1, \varphi_2 : L = \mathbb{Z}/p\mathbb{Z} \rightarrow \text{Aut}(\mathbb{Z}/p^2\mathbb{Z})$  be non trivial group homomorphisms. By the hint we know that  $\text{Aut}(\mathbb{Z}/p^2\mathbb{Z}) \cong \mathbb{Z}/m\mathbb{Z}$  for some integer  $m$ . Since the two homomorphisms are non trivial, their kernel must be trivial and  $\varphi_1$  and  $\varphi_2$  are thus injective. It follows that  $\varphi_1(L)$  and  $\varphi_2(L)$  are subgroups of order  $p$  in  $\mathbb{Z}/m\mathbb{Z}$ . But cyclic groups have *unique* subgroups of each order, hence  $\varphi_1(L) = \varphi_2(L)$ . In particular they are conjugate, and we conclude by using the first part of the exercise.

(3) Let  $\varphi_1, \varphi_2 : L = \mathbb{Z}/p\mathbb{Z} \rightarrow \text{Aut}(\mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z} = K)$  be non trivial group homomorphisms. To identify the codomain, we note that every automorphism  $f : K \rightarrow K$  is a  $L$ -vector space linear automorphism. This is because for  $\alpha \in L$  we have that  $f(\alpha \cdot (a, b)) = \alpha \cdot f(a, b)$ . It follows that  $L$ -automorphisms are in bijections with invertible matrices with coefficients in  $L$ , and therefore

$$|\text{Aut}(K)| = |\text{GL}_2(\mathbb{Z}/p\mathbb{Z})| = (p^2 - 1)(p^2 - p) = p(p^3 - p^2 - p - 1) = p \cdot r$$

for some even number  $r \in \mathbb{N}$ . It follows that Sylow  $p$ -subgroups are of order  $p$ , and hence all subgroups of order  $p$  are conjugate (by Sylow's theorem). As in the previous point the groups  $\varphi_1(L)$  and  $\varphi_2(L)$  are subgroups of  $\text{Aut}(K)$  of order  $p$ , and hence are conjugate. We conclude by the first point.