

Exercise Sheet #5

Course Instructor: Ethan Ackelsberg
Teaching Assistant: Felipe Hernández

P1. Let (X, \mathcal{B}, μ) σ -finite measure space. Show that

- (a) μ is s -finite.
- (b) μ is semi-finite.

P2. Let (X, \mathcal{B}, μ) be a measure space. Define $\tilde{\mathcal{B}} = \{E \subseteq X \mid \forall F \in \mathcal{B}, \mu(F) < \infty \Rightarrow E \cap F \in \mathcal{B}\}$.

- (a) Prove that $\tilde{\mathcal{B}}$ is a sigma algebra.
- (b) Define $\tilde{\mu}$ on M by $\tilde{\mu}(E) = \mu(E)$ for $E \in \mathcal{B}$ and $\tilde{\mu}(E) = \infty$ otherwise. Prove that $\tilde{\mu}$ is a saturated measure on $\tilde{\mathcal{B}}$.

P3. Let \mathcal{P} be a π -system that contains X and \mathcal{F} a family of functions from X to \mathbb{R} such that

- (a) $A \in \mathcal{P} \implies \mathbf{1}_A \in \mathcal{F}$,
- (b) \mathcal{F} is a real vector space: $f, g \in \mathcal{F}$ and $c \in \mathbb{R} \implies cf + g \in \mathcal{F}$,
- (c) if $(f_n)_{n \in \mathbb{N}}$ is a non-decreasing sequence of positive functions in \mathcal{F} and $f = \lim_{n \rightarrow \infty} f_n$ bounded, then $f \in \mathcal{F}$.

Show that \mathcal{F} contains the set $\{f : X \rightarrow \mathbb{R} \mid f \text{ is a bounded } \sigma(\mathcal{P})\text{-measurable function}\}$.

P4. We will show that if (X, \mathcal{B}, μ) is a non-atomic probability space, then for all $t \in [0, 1]$, there is $E \in \mathcal{B}$ such that $\mu(E) = t$. For this:

- (a) Show that for every $s \in (0, 1)$, there is $E \in \mathcal{B}$ such that $\mu(E) \in (0, s)$.
- (b) Fix $t \in (0, 1)$. Construct a family of disjoint sets $(E_n)_{n \in \mathbb{N}} \subseteq \mathcal{B}$ such that:
 - i) For each $n \in \mathbb{N}$, $\mu(\bigcup_{i=1}^n E_i) < t$.
 - ii) If it is possible, for each $n \in \mathbb{N}$, E_n is chosen such that $\mu(E_n) \geq \frac{1}{n}$.

Show that $\mu(\bigcup_{n \in \mathbb{N}} E_n) = t$.

Hint: If the latter is not true, then find $F \in \mathcal{B}$ such that $0 < \mu(F) < t - \mu(\bigcup_{n \in \mathbb{N}} E_n)$. What does this imply for condition ii) of the definition of the $(E_n)_{n \in \mathbb{N}}$?

P5. Verify if the following are examples of π -system and/or λ -systems:

- (a) The collection $\mathcal{P} = \{(a, b] : a, b \in \mathbb{R}\}$ of half-open intervals in \mathbb{R}
- (b) Given two measurable spaces (X, \mathcal{B}) and (Y, \mathcal{C}) , the family $\mathcal{P} = \{B \times C : B \in \mathcal{B}, C \in \mathcal{C}\}$ of “rectangles” in $X \times Y$.
- (c) For two probability measures μ, ν on a measurable space (X, \mathcal{B}) , the family $\mathcal{L} = \{E \in \mathcal{B} : \mu(E) = \nu(E)\}$.