

Combinatorial Number Theory

LECTURE NOTES

Contents

1	Ramsey's Theorem	3
1.1	Ramsey's Theorem for graphs	3
1.2	The compactness principle for colorings	5
1.3	Ramsey's Theorem for 2-sets	6
1.4	Schur's Theorem	7
1.5	Ramsey's Theorem for k -sets	10
1.6	Ramsey's Theorem for hypergraphs	11
1.7	Erdős-Szekeres' Theorem on convex polygons	12
1.8	Erdős-Szekeres' Theorem on monotone paths	15
2	van der Waerden's Theorem	17
2.1	Notions of largeness	17
2.2	Syndetic sets and thick sets	19
2.3	van der Waerden's Theorem – equivalent forms	21
2.4	Proof of van der Waerden's Theorem	23
2.5	Gallai's Theorem	24

Chapter 1

Ramsey's Theorem

1.1. Ramsey's Theorem for graphs

Definition 1. A graph $G = (V, E)$ is a set V of points, called *vertices*, and a set E of distinct pairs of vertices, called *edges*.

Definition 2. A subgraph $G' = (V', E')$ of a graph $G = (V, E)$ is a graph such that $V' \subseteq V$ and $E' \subseteq E$.

Figure 1.1 below depicts a graph G with four vertices $V = \{V_1, V_2, V_3, V_4\}$ and four edges $E = \{e_1, e_2, e_3, e_4\}$, where $e_1 = \{V_1, V_2\}$, $e_2 = \{V_2, V_3\}$, $e_3 = \{V_3, V_4\}$, and $e_4 = \{V_2, V_4\}$. Note that edges are *unordered* pairs of vertices, meaning that $\{V_1, V_2\}$ and $\{V_2, V_1\}$ refer to the same edge. Next to it is a graph $G' = (V', E')$ with $V' = V = \{V_1, V_2, V_3, V_4\}$ and $E' = \{e_1, e_3\}$. Since $V' \subseteq V$ and $E' \subseteq E$, we deduce that G' is a subgraph of G .




Figure 1.1: A graph G and one of its subgraphs G' .

Definition 3. Given $n \in \mathbb{N}$, a *complete graph on n vertices*, denoted by K_n , is a graph with n vertices and the property that every pair of distinct vertices is connected by an edge.




Figure 1.2: A depiction of K_n for $n = 2, 3, 4, 5$, and 6.

Definition 4. An *edge-coloring* of a graph $G = (V, E)$ is an assignment of a color to each edge of the graph. A graph that has been edge-colored is called *monochromatic* if all of its edges are the same color.

An edge-coloring of a graph can also be viewed as a function where the domain is the set of edges of the graph and the codomain is the set of colors. For example, suppose one has a graph with edges $E = \{e_1, e_2, e_3\}$ and a set of colors $C = \{\text{red}, \text{blue}\}$. A valid coloring of this graph can be seen as a function $\chi: E \rightarrow C$, where, for instance, $\chi(e_1) = \text{red}$, $\chi(e_2) = \text{blue}$, and $\chi(e_3) = \text{red}$.

Ramsey's Theorem for graphs. For any $n, m \in \mathbb{N}$ there exists $R = R(n, m) \in \mathbb{N}$ such that any edge-coloring of K_R with at most m colors contains a monochromatic copy of K_n as a subgraph.

Let us illustrate the content of Ramsey's Theorem for graphs by looking at an example. If the edge-coloring consists only of two colors, say **red** and **blue**, and we assume $n = 3$, then Ramsey's Theorem asserts that there exists a number $R(3, 2)$ such that any edge-coloring of a complete graph on $R(3, 2)$ vertices admits a monochromatic triangle. Note that $R(3, 2)$ cannot equal 5, because Figure 1.3 below shows a 2-coloring of K_5 containing no monochromatic triangle. However, taking




Figure 1.3: An edge-coloring of K_5 containing no monochromatic copy of K_3 .

$R(3,2) = 6$ already works. Indeed, through some trial-and-error, one quickly realizes that it is impossible to find an edge-coloring of K_6 using only 2 colors that avoids monochromatic triangles. For instance, Figure 1.4 below shows a complete graph on 6 vertices where all but one edge have been colored either red or blue. As can be seen from the picture, it is impossible to complete the coloring without creating either a red or a blue triangle.




Figure 1.4: An almost-complete edge-coloring of K_6 that cannot be completed without creating a monochromatic copy of K_3 . This example illustrates that it is impossible to color K_6 using two colors without producing a monochromatic copy of K_3 .

The best possible value for $R(n,m)$ is called the *Ramsey number* for (n,m) . Below is a list of Ramsey numbers known to date:

(n,m)	Ramsey Number
$(3,2)$	6
$(4,2)$	18
$(3,3)$	17
$(3,4)$	30
$(5,2)$	unknown
$(3,5)$	unknown
$(4,3)$	unknown
\vdots	

1.2. The compactness principle for colorings

Compactness Theorem for finite colorings. Let Y be an infinite set, let $m \in \mathbb{N}$, and let \mathcal{F} be a collection of finite subsets of Y . The following are equivalent:

- (i) For any coloring of Y using no more than m colors there exists $F \in \mathcal{F}$ such that all elements in F have the same color.
- (ii) There exists a finite set $Z \subseteq Y$ such that for any finite coloring of Z using no more than m colors there exists $F \in \mathcal{F}$ with $F \subseteq Z$ and such that all elements in F have the same color.

Proof. The implication (ii) \implies (i) is immediate, so it only remains to prove (i) \implies (ii). We can view a coloring of Y that uses no more than m colors as a function $\chi: Y \rightarrow \{1, \dots, m\}$ simply by associating a number from 1 to m with each color. This means the space of all possible colorings of Y can be identified with the product space $\{1, \dots, m\}^Y$. Note that the finite set $\{1, \dots, m\}$, endowed with the discrete topology, is a compact Hausdorff space. By Tychonoff's theorem, $\{1, \dots, m\}^Y$ endowed with the product topology is therefore also a compact Hausdorff space.

For any finite non-empty set $Z \subseteq Y$ let \mathcal{C}_Z be the set of all colorings in $\{1, \dots, m\}^Y$ for which there is monochromatic $F \in \mathcal{F}$ with $F \subseteq Z$. Then \mathcal{C}_Z is an open set in the product topology on $\{1, \dots, m\}^Y$. Moreover, in light of statement (i), we have

$$\bigcup_{\substack{Z \subseteq Y \\ 0 < |Z| < \infty}} \mathcal{C}_Z = \{1, \dots, m\}^Y.$$

By compactness, the above cover admits a finite subcover, or in other words, there exist finite non-empty sets $Z_1, \dots, Z_\ell \subseteq Y$ such that $\mathcal{C}_{Z_1} \cup \dots \cup \mathcal{C}_{Z_\ell} = \{1, \dots, m\}^Y$. Taking $Z = Z_1 \cup \dots \cup Z_\ell$ and noting that $\mathcal{C}_{Z_1} \cup \dots \cup \mathcal{C}_{Z_\ell} \subseteq \mathcal{C}_{Z_1 \cup \dots \cup Z_\ell}$, it follows that $\mathcal{C}_Z = \{1, \dots, m\}^Y$, completing the proof. \square

1.3. Ramsey's Theorem for 2-sets

Definition 5. A *2-set* is a set consisting of exactly two elements. Given a set X , a *2-subset* of X is any subset of X that is a 2-set. We will use $X^{(2)}$ to denote the set of all 2-subsets of X .

We have already seen examples of 2-subsets in the previous section. Indeed, the set of edges E of a graph $G = (V, E)$ consists of 2-subsets of the set of vertices V . In other words, $E \subseteq V^{(2)}$. Note that a graph $G = (V, E)$ is a complete graph if and only if $E = V^{(2)}$.

Definition 6. Let X be a set. A *coloring of $X^{(2)}$* is an assignment of a color to each 2-subset of X . We call $X^{(2)}$ *monochromatic* if all elements in $X^{(2)}$ have the same color.

The following can be viewed as an “infinitary” version of Ramsey's Theorem for graphs.

Ramsey's Theorem for 2-sets. *Let X be an infinite set. Then for any finite coloring of $X^{(2)}$ there exists an infinite subset $Y \subseteq X$ such that $Y^{(2)}$ is monochromatic.*

Proof. Fix an arbitrary element $x_1 \in X$ and note that any 2-set of the form $\{x_1, x\}$ for $x \in X \setminus \{x_1\}$ has a certain color. Since the number of colors is finite but the set $X \setminus \{x_1\}$ is infinite, there exists an infinite subset $X_1 \subseteq X \setminus \{x_1\}$ such that all 2-sets of the form $\{x_1, x\}$ for $x \in X_1$ have the same color. Now fix an arbitrary element $x_2 \in X_1$

and let us repeat the same procedure. Any 2-set of the form $\{x_2, x\}$ for $x \in X_1 \setminus \{x_2\}$ has a certain color. For the same reason as before, since the number of colors is finite but the set $X_1 \setminus \{x_2\}$ is infinite, there exists an infinite subset $X_2 \subseteq X_1 \setminus \{x_2\}$ such all 2-sets of the form $\{x_2, x\}$ for $x \in X_2$ have the same color. Continuing this procedure produces an infinite sequence of distinct elements x_1, x_2, x_3, \dots and a nested family of infinite sets $X \supseteq X_1 \supseteq X_2 \supseteq X_3 \supseteq \dots$ such that for all $i \in \mathbb{N}$ we have $x_{i+1} \in X_i$ and the set $\{\{x_i, x\} : x \in X_i\}$ is monochromatic.

Let c_i denote the color of elements in the set $\{\{x_i, x\} : x \in X_i\}$. Then c_1, c_2, c_3, \dots is an infinite sequence of colors. Since there are only finitely many different colors, one color must appear infinitely often in this sequence. In other words, there exists a color c and an infinite sequence $i_1 < i_2 < i_3 < \dots \in \mathbb{N}$ such that $c_{i_k} = c$ for all $k \in \mathbb{N}$.

To finish the proof, define $Y = \{x_{i_k} : k \in \mathbb{N}\}$ and observe that any 2-subset of Y is of the form $\{x_{i_k}, x_{i_\ell}\}$ for $k < \ell \in \mathbb{N}$. Since $x_{i_\ell} \in X_{i_{\ell-1}}$ and $X_{i_{\ell-1}} \subseteq X_{i_k}$, the 2-set $\{x_{i_k}, x_{i_\ell}\}$ has the color c . Hence all 2-subsets of Y have the color c , which proves that $Y^{(2)}$ is monochromatic. \square

Proposition 7. *Ramsey's Theorem for 2-sets implies Ramsey's Theorem for graphs.*

Proof. Fix $n, m \in \mathbb{N}$. Let $\mathbb{N}^{(2)}$ denote the set of all 2-element subsets of \mathbb{N} , and define

$$\mathcal{F} = \{F^{(2)} : F \subseteq \mathbb{N}, |F| = n\}.$$

By Ramsey's Theorem for 2-sets, any coloring of $\mathbb{N}^{(2)}$ with at most m colors admits a monochromatic set of the form $F^{(2)}$ for some $F \subseteq \mathbb{N}$ with $|F| = n$.

By the Compactness Theorem for finite colorings, there exists a finite set $Z \subseteq \mathbb{N}^{(2)}$ such that any coloring of Z with at most m colors already contains a monochromatic $F^{(2)}$ with $F \subseteq \mathbb{N}$ and $|F| = n$. Enlarging Z if necessary, we may assume $Z = H^{(2)}$ for some finite $H \subseteq \mathbb{N}$.

In other words, for every m -coloring of $H^{(2)}$, there exists a subset $F \subseteq H$ of size n such that $F^{(2)}$ is monochromatic. Define $R = R(n, m) := |H|$. Identifying $H^{(2)}$ with the edge set of the complete graph K_R , the subset $F^{(2)}$ corresponds to a copy of K_n inside K_R . Thus we have shown that any edge-coloring of K_R with at most m colors contains a monochromatic copy of K_n . This completes the proof. \square

1.4. Schur's Theorem

Fermat's Last Theorem states that for $m \geq 3$ the equation

$$x^m + y^m = z^m \tag{1.4.1}$$

has no positive integer solutions $x, y, z \in \mathbb{N}$. For centuries, this remained one of the biggest open problems in mathematics, and one whose intriguing nature captivated many mathematicians. Among them was also Issai Schur, who investigated a

natural, localized version of Fermat's Last Theorem. More precisely, he wondered whether for any $m \geq 2$ the congruence equation

$$x^m + y^m \equiv z^m \pmod{p} \quad (1.4.2)$$

possesses non-trivial solutions for all but finitely many primes p . Note that any non-trivial solution to Fermat's equation $x^m + y^m = z^m$ also offers a non-trivial solution to Schur's equation $x^m + y^m \equiv z^m \pmod{p}$ for all primes p satisfying $p > z^m$, but not the other way around. In order to address (1.4.2), Schur proved a theorem that is often regarded as the earliest result in Ramsey Theory:

Schur's Theorem ([Sch17]). *For any $m \in \mathbb{N}$ there exists $S = S(m) \in \mathbb{N}$ such that if the set $\{1, \dots, S\}$ is colored using at most m colors then there exist monochromatic $x, y, z \in \{1, \dots, S\}$ with $x + y = z$.*

Proof. Take $S = R(3, m)$, where $R(3, m)$ is the Ramsey number for $(3, m)$. Let K_S denote the complete graph on S vertices and denote the vertices of K_S by V_1, V_2, \dots, V_S . Any coloring of the set $\{1, \dots, S\}$ induces an edge-coloring on K_S by assigning to each edge $\{V_i, V_j\}$ the color of the number $|i - j| \in \{1, \dots, S\}$. According to Ramsey's Theorem for graphs, K_S contains a monochromatic triangle. Let V_a, V_b , and V_c , for $a < b < c$, be the vertices of this monochromatic triangle. By setting

$$x = b - a, \quad y = c - b, \quad \text{and} \quad z = c - a,$$

it is then easy to check that x, y, z have the same color and satisfy $x + y = z$. \square

The smallest possible positive integer $S(m)$ for which the conclusion of Schur's Theorem holds is referred to as the *Schur number* for m . The known Schur numbers to date are:

m	Schur Number
2	5
3	14
4	45
5	161
6	unknown
7	unknown
\vdots	

Here is an example from Schur's original paper [Sch17] of a 3-coloring of $\{1, \dots, 13\}$ admitting no monochromatic solution to the equation $x + y = z$:

color 1: {2, 3, 11, 12}

color 2: {5, 6, 8, 9}

color 3: {1, 4, 7, 10, 13}

More examples along these lines can be found here: <https://oeis.org/A030126>.

The proof that the Schur number for 5-colorings equals 161 took up 2 petabytes of space. Even though every 5-coloring of $\{1, \dots, 161\}$ admits a monochromatic solution to $x + y = z$, there are 2447113088 many 5-colorings of $\{1, \dots, 160\}$ admitting no monochromatic solution to $x + y = z$.

With the help of the above theorem, Schur was able to show that, contrary to Fermat's equation (1.4.1), its "local" counterpart (1.4.2) does possess non-trivial solutions.

Theorem 8. *Let $m \in \mathbb{N}$. There exists $F = F(m)$ such that for all prime numbers $p > F$ there exist $x, y, z \in \{1, \dots, p-1\}$ with $x^m + y^m \equiv z^m \pmod{p}$.*

For the proof of Theorem 8, we will need the following basic fact from algebra, the proof of which is left to the interested reader.

Lemma 9. *Let $(K, +, \cdot)$ be a field and $f(x) \in K[x]$ a polynomial of degree $\deg(f) = m$ with coefficients in K . Then the number of roots of $f(x)$ is at most m .*

Let us now see the proof of Theorem 8.

Proof of Theorem 8. Take $F = S(m)$, where $S(m)$ is as guaranteed by Schur's Theorem. Let p be any prime number bigger than F . The set $\mathbb{F}_p = \{0, 1, \dots, p-1\}$ of congruence classes modulo p naturally forms a field $(\mathbb{F}_p, +, \cdot)$ under the modular arithmetic operations $+$ and \cdot . Let $\mathbb{F}_p^\times = \mathbb{F}_p \setminus \{0\}$ and consider the set

$$C := \{x^m : x \in \mathbb{F}_p^\times\}.$$

Note that C is a subgroup of the multiplicative group $(\mathbb{F}_p^\times, \cdot)$. This means that \mathbb{F}_p^\times can be covered by cosets of C . More precisely, there exist coset representatives $g_1, g_2, \dots, g_r \in \mathbb{F}_p^\times$ such that

$$\mathbb{F}_p^\times = g_1 C \cup g_2 C \cup \dots \cup g_r C. \quad (1.4.3)$$

Any element of \mathbb{F}_p^\times is a root of the polynomial $x^m - a$ for some $a \in C$, or equivalently,

$$\mathbb{F}_p^\times = \bigcup_{a \in C} \{x : x^m - a \equiv 0 \pmod{p}\}.$$

But since each such polynomial has at most m roots due to Lemma 9, we have

$$|\{x : x^m - a \equiv 0 \pmod{p}\}| \leq m.$$

It follows that

$$|\mathbb{F}_p^\times| = \left| \bigcup_{a \in C} \{x : x^m - a \equiv 0 \pmod{p}\} \right| \leq \sum_{a \in C} |\{x : x^m - a \equiv 0 \pmod{p}\}| \leq m \cdot |C|.$$

So C can have at most m cosets, or in other words, $r \leq m$. Since $p > F$, the set $\{1, \dots, F\}$ is a subset of $\mathbb{F}_p^\times = \{1, \dots, p-1\}$ and hence (1.4.3) yields a partition of the set $\{1, \dots, F\}$ involving r disjoint cells. We can think of this partition as a coloring of $\{1, \dots, F\}$ using r colors. Since $F = S(m)$ and $r \leq m$, it follows from Schur's Theorem that there exist monochromatic $\tilde{x}, \tilde{y}, \tilde{z} \in \{1, \dots, F\}$ for which $\tilde{x} + \tilde{y} = \tilde{z}$. Since $\tilde{x}, \tilde{y}, \tilde{z}$ have

the same color, they all belong to the same coset. In other words, there exists a coset representative $g_i \in \{g_1, \dots, g_r\}$ such that $\tilde{x}, \tilde{y}, \tilde{z} \in g_i C$. Take any $x, y, z \in \mathbb{F}_p^\times$ for which

$$\tilde{x} \equiv g_i x^m \pmod{p}, \quad \tilde{y} \equiv g_i y^m \pmod{p}, \quad \text{and} \quad \tilde{z} \equiv g_i z^m \pmod{p},$$

which is possible because $\tilde{x}, \tilde{y}, \tilde{z} \in g_i C$. Then we have

$$g_i x^m + g_i y^m \equiv g_i z^m \pmod{p},$$

from which it follows that

$$x^m + y^m \equiv z^m \pmod{p},$$

because $g_i \not\equiv 0 \pmod{p}$. □

1.5. Ramsey's Theorem for k -sets

Definition 10. A k -set is a set consisting of exactly k elements. Given a set X , a k -subset of X is any subset of X that is a k -set. We will use $X^{(k)}$ to denote the set of all k -subsets of X .

We have already seen Ramsey's Theorem for 2-sets. Here is Ramsey's result in full generality.

Ramsey's Theorem for k -sets ([Ram30]). *Let X be an infinite set and $k \geq 2$. Then for any finite coloring of $X^{(k)}$ there exists an infinite subset $Y \subseteq X$ such that $Y^{(k)}$ is monochromatic.*

Proof. Let us use a proof by induction on k . The base case of the induction, when $k = 2$, follows from Ramsey's Theorem for 2-sets established in Section 1.3. To prove the inductive step, assume $k \geq 3$ and Ramsey's Theorem has already been proven for $(k - 1)$ -sets. Let $Y_0 = X$ and fix an arbitrary element $y_1 \in Y_0$. Note that any k -set of the form $\{y_1, x_2, \dots, x_k\}$ for $\{x_2, \dots, x_k\} \in (Y_0 \setminus \{y_1\})^{(k-1)}$ has a certain color, which induces a finite coloring on $(Y_0 \setminus \{y_1\})^{(k-1)}$. Applying Ramsey's Theorem for $(k - 1)$ -sets, we can find an infinite subset $Y_1 \subseteq Y_0 \setminus \{y_1\}$ such that all k -sets of the form $\{y_1, x_2, \dots, x_k\}$ for $\{x_2, \dots, x_k\} \in Y_1^{(k-1)}$ are monochromatic. Next, fix an arbitrary element $y_2 \in Y_1$ and repeat the same procedure. The given coloring of k -sets of the form $\{y_2, x_2, \dots, x_k\}$ for $\{x_2, \dots, x_k\} \in (Y_1 \setminus \{y_2\})^{(k-1)}$ induces a finite coloring of $(Y_1 \setminus \{y_2\})^{(k-1)}$. Applying Ramsey's Theorem for $(k - 1)$ -sets once more yields an infinite subset $Y_2 \subseteq Y_1 \setminus \{y_2\}$ such that all k -sets of the form $\{y_2, x_2, \dots, x_k\}$ for $\{x_2, \dots, x_k\} \in Y_2^{(k-1)}$ are monochromatic. Continuing this procedure produces an infinite sequence of distinct elements y_1, y_2, y_3, \dots and a nested family of infinite sets $X = Y_0 \supseteq Y_1 \supseteq Y_2 \supseteq Y_3 \supseteq \dots$ such that for all $i \in \mathbb{N}$ the set $\{\{y_i, x_2, \dots, x_k\} : \{x_2, \dots, x_k\} \in Y_i^{(k-1)}\}$ is monochromatic. Moreover, we have $y_{i+1} \in Y_i$ for all $i \in \mathbb{N}$.

Let c_i denote the color of elements in the set $\{y_i, x_2, \dots, x_k\} : \{x_2, \dots, x_k\} \in Y_i^{(k-1)}\}$. Since the sequence c_1, c_2, c_3, \dots is infinite but the number of colors is finite, one color must appear infinitely often in this sequence. In other words, there exists a color c and an infinite subsequence $c_{i_1}, c_{i_2}, c_{i_3}, \dots \in \mathbb{N}$ such that $c_{i_\ell} = c$ for all $\ell \in \mathbb{N}$. To finish the proof, define $Y = \{y_{i_k} : k \in \mathbb{N}\}$ and observe that any k -subset of Y is of the form $\{y_{i_{\ell_1}}, \dots, y_{i_{\ell_k}}\}$ for $\ell_1 < \dots < \ell_k \in \mathbb{N}$. Since $\{y_{i_{\ell_2}}, \dots, y_{i_{\ell_k}}\} \in Y_{i_{\ell_1}}$ because $\ell_1 < \ell_2 < \dots < \ell_k$, the k -set $\{y_{i_{\ell_1}}, \dots, y_{i_{\ell_k}}\}$ has the color c . Hence all k -subsets of Y have the color c , which proves that $Y^{(k)}$ is monochromatic. \square

1.6. Ramsey's Theorem for hypergraphs

A hypergraph is a generalization of a graph in which an edge can join multiple vertices at once.

Definition 11. Let $k \in \mathbb{N}$. A k -uniform hypergraph is a pair $G = (V, E)$ where V is a set of points, called *vertices*, and $E \subseteq V^{(k)}$ is a set of k -subsets of V , called *hyperedges*.




Figure 1.5: Here is an example of a 3-uniform hypergraph with vertices $V = \{7, 13, 17, 23, 53, 73, 97, 103, 137, 193\}$, where three vertices are connected by a hyperedge if and only if their squares form a 3-term arithmetic progression. For example, $\{7, 13, 17\}$ is an edge, because $7^2, 13^2, 17^2$ are in an arithmetic progression.

Given $k, n \in \mathbb{N}$ with $k \leq n$, a *complete k -uniform hypergraph on n vertices* is a k -uniform hypergraph $G = (V, E)$ where the set of vertices has cardinality n and where every set of k distinct vertices in V is connected by an edge. In other words, $G = (V, E)$ is a complete k -uniform hypergraph on n vertices if $|V| = n$ and $E = V^{(k)}$.

Ramsey's Theorem for hypergraphs. For any $n, m, k \in \mathbb{N}$ there exists a number $R = R_k(n, m) \in \mathbb{N}$ such that any edge-coloring of a complete k -uniform hypergraph on R vertices with at most m colors admits a monochromatic copy of a complete k -uniform hypergraph on n vertices.

Proof. Let $n, m, k \in \mathbb{N}$ be given. It follows from Ramsey's Theorem for k -sets that for any m -coloring of $\mathbb{N}^{(k)}$ there exists a set $S \subseteq \mathbb{N}$ with $|S| = n$ such that $S^{(k)}$ is monochromatic. If we now apply the Compactness Theorem for finite colorings to this statement (with $Y = \mathbb{N}^{(k)}$ and $\mathcal{F} = \{S^{(k)} : S \subseteq \mathbb{N}, |S| = n\}$), it follows that there exists some integer $R = R_k(n, m)$ such that for any m -coloring of $\{1, \dots, R\}^{(k)}$ there exists a set $S \subseteq \{1, \dots, R\}$ with $|S| = n$ such that $S^{(k)}$ is monochromatic. Note that $\{1, \dots, R\}^{(k)}$ can be identified with a complete k -uniform hypergraph on R vertices, and $S^{(k)}$ with a complete k -uniform hypergraph on n vertices. This finishes the proof. \square

1.7. Erdős-Szekeres' Theorem on convex polygons

Definition 12. A non-empty set $C \subseteq \mathbb{R}^2$ is called *convex* if for any $\vec{x}, \vec{y} \in C$ and $\lambda \in [0, 1]$ one has $\lambda\vec{x} + (1 - \lambda)\vec{y} \in C$.

The point $\lambda\vec{x} + (1 - \lambda)\vec{y}$ is usually referred to as a *convex combination* of the points \vec{x} and \vec{y} . Also observe that the set $\{\lambda\vec{x} + (1 - \lambda)\vec{y} : \lambda \in [0, 1]\}$ is just an algebraic description for the line segment joining the points \vec{x} and \vec{y} .



Figure 1.6: A convex polygon (left) and a non-convex polygon (right).

Definition 13. The *convex hull* of a non-empty set $K \subseteq \mathbb{R}^2$ is the smallest convex set that contains K .

Since the intersection of convex sets is again a convex set, it follows that the convex hull of K equals the intersection of all convex sets that contain K . The convex hull can also be described algebraically as the set of all finite convex combinations

of elements in the set. More precisely, if K is a subset of \mathbb{R}^2 and we use $\text{conv}(K)$ to denote its convex hull, then

$$\text{conv}(K) = \{w_1\vec{z}_1 + \dots + w_\ell\vec{z}_\ell : \ell \in \mathbb{N}, \vec{z}_1, \dots, \vec{z}_\ell \in K, w_1, \dots, w_\ell \in [0, 1], w_1 + \dots + w_\ell = 1\}. \quad (1.7.1)$$

Mind that the convex hull of K should not be confused with the *closed convex hull* of K , which is defined as the smallest closed convex set that contains K , and is usually denoted by $\overline{\text{conv}}(K)$ instead of $\text{conv}(K)$.

Definition 14. A non-empty set of points $K \subseteq \mathbb{R}^2$ is said to be in *convex position* if no point $\vec{x} \in K$ belongs to the convex hull of $K \setminus \{\vec{x}\}$.

For example, a finite set $K \subseteq \mathbb{R}^2$ is in convex position if and only if its elements are the corners of a convex polygon.

Definition 15. A set $K \subseteq \mathbb{R}^2$ is called *discrete* if it has no accumulation points.

Note that the notion of a discrete set introduced in Definition 15 is more commonly referred to in the literature as a “closed discrete set”, since the term “discrete” by itself usually does not include closedness. For brevity, we will simply use the term discrete.

Erdős-Szekeres’ Theorem on points in convex position. *Let K be an infinite discrete set of points in \mathbb{R}^2 . Then either there is an infinite subset of K whose points lie on a straight line or there is an infinite subset of K whose points are in convex position.*

For the proof of Erdős-Szekeres’ Theorem on points in convex position we will need the following classical result from convex geometry.

Carathéodory’s theorem. *Let $K \subseteq \mathbb{R}^2$ with $|K| \geq 4$ be given. Then K is in convex position if and only if any four distinct points from K form a convex quadrilateral.*

Proof. Clearly, if K is in convex position then any quadrilateral formed using points from K is convex. To prove the converse, we will show that if K is not in convex position then there exist four points in K such that one of these points lies within the triangle spanned by the others.

Suppose K is not in convex position. Then there exists a point $\vec{x} \in K$ lying in the convex hull of $K' = K \setminus \{\vec{x}\}$. In light of (1.7.1), this means that we can write \vec{x} as

$$\vec{x} = w_1\vec{z}_1 + \dots + w_\ell\vec{z}_\ell, \quad (1.7.2)$$

where $\vec{z}_1, \dots, \vec{z}_\ell \in K'$ and $w_1, \dots, w_\ell \in [0, 1]$ with $w_1 + \dots + w_\ell = 1$. Note that we can assume without loss of generality that $\vec{z}_1, \dots, \vec{z}_\ell$ are in convex position. Indeed, if for example \vec{z}_ℓ belongs to the convex hull of $\vec{z}_1, \dots, \vec{z}_{\ell-1}$ then we can express \vec{z}_ℓ as a convex combination of $\vec{z}_1, \dots, \vec{z}_{\ell-1}$ and substitute this representation in (1.7.2), allowing us to represent \vec{x} as a convex combination of $\vec{z}_1, \dots, \vec{z}_{\ell-1}$ instead of $\vec{z}_1, \dots, \vec{z}_\ell$. Thus, invoking induction on ℓ , we may assume that $\vec{z}_1, \dots, \vec{z}_\ell$ are in convex position. This implies that $\vec{z}_1, \dots, \vec{z}_\ell$ form the corners of a convex polygon. Since \vec{x} lies inside




Figure 1.7: A convex polygon divided into triangles.

this polygon and since convex polygons decompose into triangles (as illustrated in Figure 1.7), there exists $i < j < k \in \{1, \dots, \ell\}$ such that \vec{x} lies in the triangle spanned by $\vec{z}_i, \vec{z}_j, \vec{z}_k$, finishing the proof. \square

Proof of Erdős-Szekeres' Theorem on points in convex position. Let $K \subseteq \mathbb{R}^2$ be infinite. We begin by coloring $K^{(3)}$ by assigning the color red to $\{\vec{x}, \vec{y}, \vec{z}\} \in K^{(3)}$ if the points $\vec{x}, \vec{y}, \vec{z}$ are collinear and the color blue otherwise. According to Ramsey's Theorem for k -sets, there exists an infinite set $L \subseteq K$ such that all 3-sets in $L^{(3)}$ have the same color. If this color is red, then any three distinct points in L are collinear. This can only happen if all the points in L lie on a straight line, in which case we are done.

It remains to deal with the case when all elements in $L^{(3)}$ are blue, i.e., when no three points in L are collinear. In this situation, we need to apply Ramsey's Theorem one more time. Note that L is a discrete set. This implies that for any three points $\vec{x}, \vec{y}, \vec{z} \in L$ the triangle $\Delta\vec{x}\vec{y}\vec{z}$ contains only finitely many points from L . Color all elements in $L^{(3)}$ by assigning the color red to the 3-set $\{\vec{x}, \vec{y}, \vec{z}\} \in L^{(3)}$ if the triangle $\Delta\vec{x}\vec{y}\vec{z}$ contains an even number of points from L , and the color blue otherwise. By Ramsey's Theorem for k -sets there exists an infinite set $C \subseteq L$ such that $C^{(3)}$ is monochromatic. We claim that C is in convex position. Indeed, if C were not in convex position then, in view of Carathéodory's theorem, there exist four points $\vec{w}, \vec{x}, \vec{y}, \vec{z} \in C$ such that \vec{w} lies inside the triangle $\Delta_0 = \Delta\vec{x}\vec{y}\vec{z}$. Note that Δ_0 splits into three smaller triangles, $\Delta_1 = \Delta\vec{w}\vec{y}\vec{z}$, $\Delta_2 = \Delta\vec{w}\vec{x}\vec{z}$, and $\Delta_3 = \Delta\vec{w}\vec{x}\vec{y}$, as seen in Figure 1.8. For $i = 0, 1, 2, 3$ let $\#\Delta_i$ denote the number of points from L inside the triangle Δ_i . Since no three points from L are collinear, there are no points on the boundary of any of these triangles aside from their corners. This means that the number of points from L inside Δ_0 equals the combined number of points inside the three smaller triangles plus the point \vec{w} , or in other words,

$$\#\Delta_0 = \#\Delta_1 + \#\Delta_2 + \#\Delta_3 + 1. \quad (1.7.3)$$

Recall that $C^{(3)}$ is monochromatic. If all elements in $C^{(3)}$ are red then the quantities $\#\Delta_0, \#\Delta_1, \#\Delta_2,$ and $\#\Delta_3$ are even numbers. This would imply that the left hand side of (1.7.3) is an even number whereas the right hand side is an odd number, a contradiction. Similarly, if all elements in $C^{(3)}$ are blue then $\#\Delta_0, \#\Delta_1, \#\Delta_2, \#\Delta_3$ are odd numbers, implying that the left hand side of (1.7.3) is odd whereas the right




Figure 1.8

hand side is even. Either way, we have obtained a contradiction, which means that C is in convex position. \square

The following is a big open conjecture at the interface of convex geometry and Ramsey theory, posed by Erdős and Szekeres in 1960.

Conjecture (Erdős-Szekeres convex polygon problem). *Let $n \geq 3$. Any set of $2^{n-2} + 1$ points in the plane, no three of which are collinear, contains a subset of n points in convex position.*

1.8. Erdős-Szekeres' Theorem on monotone paths

Erdős-Szekeres' Theorem on monotone paths. *Fix $n, m \in \mathbb{N}$. Any sequence of distinct real numbers of length at least $nm + 1$ admits either a monotonically increasing subsequence of length $n + 1$ or a monotonically decreasing subsequence of length $m + 1$.*

Proof. Let $x_1, x_2, \dots, x_{nm+1}$ be a sequence of distinct real numbers. Label each element x_i in the sequence with the pair (a_i, b_i) , where a_i is the length of the longest monotonically increasing subsequence ending with x_i and b_i is the length of the longest monotonically decreasing subsequence ending with x_i . Note that any two elements in the sequence are labeled with a different pair: if $i < j$ and $x_i < x_j$ then $a_i < a_j$, and on the other hand if $x_i > x_j$ then $b_i < b_j$. If $a_i \leq n$ and $b_i \leq m$ for all i then there are only nm possible labels, contradicting the fact that there are $nm + 1$ elements in the sequence each with a unique label. It follows that either $a_i > n$ or $b_i > m$ for some i , yielding either an increasing sequence of length at least $n + 1$ or a decreasing sequence of length at least $m + 1$. \square

Chapter 2

van der Waerden's Theorem

2.1. Notions of largeness

The goal of this section is to develop a general framework for dealing with notions of largeness for sets. In what follows, let X be a set and \mathcal{P} a family of subsets of X . Since any reasonable notion of largeness is closed under supersets, the following definition will be very useful for our purposes.

Definition 16. We call \mathcal{P} *upward closed* if for all $A \subseteq B \subseteq X$ we have $A \in \mathcal{P} \implies B \in \mathcal{P}$.

Natural examples of upward closed families include the set of all infinite subsets and the set of all cofinite subsets of a given infinite set X ,

$$\mathcal{P}_{\text{inf}} = \{A \subseteq X : A \text{ is infinite}\} \quad \text{and} \quad \mathcal{P}_{\text{cofin}} = \{A \subseteq X : A \text{ is cofinite}\}.$$

Another example of an upward closed family is the collection of all sets that share a common point,

$$\mathcal{P}_{\{x\}} = \{A \subseteq X : x \in A\}$$

where $x \in X$ is fixed.

Definition 17. The *dual family* of \mathcal{P} , denoted by \mathcal{P}^* , is defined as

$$\mathcal{P}^* = \{A \subseteq X : A \cap B \neq \emptyset \text{ for all } B \in \mathcal{P}\}.$$

The families \mathcal{P}_{inf} and $\mathcal{P}_{\text{cofin}}$ are mutually dual, meaning that $\mathcal{P}_{\text{inf}}^* = \mathcal{P}_{\text{cofin}}$ and $\mathcal{P}_{\text{cofin}}^* = \mathcal{P}_{\text{inf}}$, whereas the family $\mathcal{P}_{\{x\}}$ is self-dual in the sense that $\mathcal{P}_{\{x\}} = \mathcal{P}_{\{x\}}^*$. Note that if \mathcal{P} is upward closed then its dual \mathcal{P}^* is also upward closed. Also, if \mathcal{P} is upward closed then we have the following two convenient properties:

- For any set $A \subseteq X$,

$$A \in \mathcal{P}^* \iff A^c \notin \mathcal{P}, \tag{2.1.1}$$

where $A^c = X \setminus A$ denotes the complement of A in X .

- $\mathcal{P}^{**} = \mathcal{P}$.

Definition 18. The family \mathcal{P} is called *partition regular* if for any finite coloring of a set $A \in \mathcal{P}$ there exists a monochromatic subset of A that belongs to \mathcal{P} .

Using a standard “color blindness” argument, we deduce that any upward closed family \mathcal{P} is partition regular if and only if for any disjoint $A, B \subseteq X$ with $A \cup B \in \mathcal{P}$ either $A \in \mathcal{P}$ or $B \in \mathcal{P}$. With some additional work, one can even remove the word “disjoint” from this statement.

Definition 19. We say a family of sets \mathcal{P} is *closed under finite intersections* if for any $A_1, \dots, A_k \in \mathcal{P}$ we have $A_1 \cap \dots \cap A_k \in \mathcal{P}$.

Coming back to our previous examples, we see that the family \mathcal{P}_{inf} is partition regular but not closed under finite intersections, whereas the family $\mathcal{P}_{\text{cofn}}$ is not partition regular but closed under finite intersections. In contrast, the family $\mathcal{P}_{\{x\}}$ is simultaneously partition regular and closed under finite intersections. These observations are explained by the next proposition.

Proposition 20. Let \mathcal{P} be an upward closed family of subsets of a set X . Then \mathcal{P} is partition regular if and only if \mathcal{P}^* is closed under finite intersections.

Proof. The proof is divided into two parts, addressing the forward and backward implications of the equivalence.

(\Rightarrow) Suppose \mathcal{P} is partition regular, let $A_1, \dots, A_k \in \mathcal{P}^*$, and define $C_i = A_i^c$ for $i = 1, \dots, k$. In view of (2.1.1) we have $C_1, \dots, C_k \notin \mathcal{P}$. As \mathcal{P} is partition regular, it follows from $C_1, \dots, C_k \notin \mathcal{P}$ that $\bigcup_{i=1}^k C_i \notin \mathcal{P}$. Using (2.1.1) once more we get

$$\left(\bigcup_{i=1}^k C_i \right)^c = \bigcap_{i=1}^k A_i \notin \mathcal{P}^*.$$

This proves that \mathcal{P}^* is closed under finite intersections.

(\Leftarrow) Assume \mathcal{P}^* is closed under finite intersections, let $C_1, \dots, C_k \in \mathcal{P}$, and assume $\bigcup_{i=1}^k C_i \in \mathcal{P}$. Define $A_i = C_i^c$ for $i = 1, \dots, k$ and note that from (2.1.1) and $\bigcup_{i=1}^k C_i \in \mathcal{P}$ we have

$$\bigcap_{i=1}^k A_i \notin \mathcal{P}^*.$$

Since \mathcal{P}^* is closed under finite intersections, it follows that for some $i \in \{1, \dots, k\}$ we must have $A_i \notin \mathcal{P}^*$. By (2.1.1) we conclude that $C_i \in \mathcal{P}$, showing that \mathcal{P} is partition regular. \square

Proposition 21. Let \mathcal{P} be upward closed. Then the family $\mathcal{P} \wedge \mathcal{P}^* = \{A \cap B : A \in \mathcal{P}, B \in \mathcal{P}^*\}$ is partition regular.

Proof. Suppose $C \in \mathcal{P} \wedge \mathcal{P}^*$. It suffices to show that if $C = C_1 \cup C_2$ with $C_1 \cap C_2 = \emptyset$ then either $C_1 \in \mathcal{P} \wedge \mathcal{P}^*$ or $C_2 \in \mathcal{P} \wedge \mathcal{P}^*$ (see the remark after Definition 18). Pick

$A \in \mathcal{P}$ and $B \in \mathcal{P}^*$ such that $C = A \cap B$, and define $D = C_1 \cup A^c$. If $D \in \mathcal{P}^*$ then $C_1 = A \cap D$ belongs to $\mathcal{P} \wedge \mathcal{P}^*$ and we are done. On the other hand, if $D \notin \mathcal{P}^*$ then $D^c \in \mathcal{P}$ (by (2.1.1)) and $C_2 = D^c \cap B$, which implies $C_2 \in \mathcal{P} \wedge \mathcal{P}^*$ and we are also done. \square

2.2. Syndetic sets and thick sets

In what follows, let $A - n = \{m \in \mathbb{N} : m + n \in A\}$.

Definition 22. A set of positive integers $S \subseteq \mathbb{N}$ is called *syndetic* if there exists $h \in \mathbb{N}$ such that $S \cup (S - 1) \cup \dots \cup (S - h) = \mathbb{N}$.

Observe that syndetic sets are characterized by the property that the distance between consecutive elements is bounded. In other words, if $s_1 < s_2 < \dots$ is an increasing enumeration of elements in S , then S is syndetic if and only if $\sup_{k \in \mathbb{N}} (s_{k+1} - s_k) < \infty$. For this reason, syndetic sets are sometimes also referred to as *sets with bounded gaps*.

Definition 23. A set of positive integers $T \subseteq \mathbb{N}$ is called *thick* if for every $h \in \mathbb{N}$ the intersection $T \cap (T - 1) \cap \dots \cap (T - h)$ is non-empty.

Thick sets are characterized by the property that they contain arbitrarily long blocks of consecutive integers, i.e., a set $T \subseteq \mathbb{N}$ is thick if and only if for every $h \in \mathbb{N}$ there exists $n \in \mathbb{N}$ such that $\{n, n + 1, \dots, n + h\} \subseteq T$.

Let us use \mathcal{P}_{syn} to denote the family of all syndetic subsets of \mathbb{N} and $\mathcal{P}_{\text{thick}}$ for the family of all thick subsets of \mathbb{N} .

Proposition 24. The families \mathcal{P}_{syn} and $\mathcal{P}_{\text{thick}}$ are dual, i.e., $\mathcal{P}_{\text{syn}}^* = \mathcal{P}_{\text{thick}}$ and $\mathcal{P}_{\text{thick}}^* = \mathcal{P}_{\text{syn}}$.

Proof. Since any syndetic set has bounded gaps, it must have non-empty intersection with any thick set, because thick sets contain arbitrarily long intervals. From this, it follows that $\mathcal{P}_{\text{syn}} \subseteq \mathcal{P}_{\text{thick}}^*$. On the other hand, if a set intersects every thick set then its complement cannot be thick. If the complement is not thick then the set itself must have bounded gaps, i.e., it is syndetic. This implies $\mathcal{P}_{\text{thick}}^* \subseteq \mathcal{P}_{\text{syn}}$. In conclusion, we have $\mathcal{P}_{\text{syn}} = \mathcal{P}_{\text{thick}}^*$, which implies $\mathcal{P}_{\text{syn}}^* = \mathcal{P}_{\text{thick}}^{**} = \mathcal{P}_{\text{thick}}$ as desired. \square

Definition 25. Sets belonging to $\mathcal{P}_{\text{syn}} \wedge \mathcal{P}_{\text{thick}}$ are called *piecewise syndetic* sets.

Piecewise syndetic sets are characterized by the property that they have bounded gaps on arbitrarily large intervals. Here is a more intuitive explanation of what this means. Let A be a subset of \mathbb{N} and let a_n denote the n -th element of A , so that a_1, a_2, a_3, \dots becomes an increasing enumeration of elements in A . Then A is piecewise syndetic if and only if there exists some number $h \in \mathbb{N}$ with the following property: Somewhere in $A = \{a_1, a_2, a_3, \dots\}$ there are two consecutive

elements a_n, a_{n+1} whose distance $a_{n+1} - a_n$ is at most h . Somewhere else in A there are three consecutive elements a_m, a_{m+1}, a_{m+2} such that the distance between the first and the second $a_{m+1} - a_m$ and the distance between the second and the third $a_{m+2} - a_{m+1}$ are at most h . Then, somewhere else in the set, there exist four consecutive elements $a_k, a_{k+1}, a_{k+2}, a_{k+3}$ such that the distances $a_{k+1} - a_k, a_{k+2} - a_{k+1}, a_{k+3} - a_{k+2}$ are all at most h . And so on. This is another way of characterizing piecewise syndeticity.

Corollary 26. *Piecewise syndetic sets are partition regular.*

Proof. This follows by combining Proposition 21 and Proposition 24. \square

Proposition 27. *Let $A \subseteq \mathbb{N}$ be piecewise syndetic. Then there exists a syndetic set L such that for any finite, non-empty $F \subseteq L$ the intersection*

$$\bigcap_{n \in F} (A - n) \tag{2.2.1}$$

is piecewise syndetic.

Proof. By definition, if $A \subseteq \mathbb{N}$ is piecewise syndetic, then there exists $h \in \mathbb{N}$ such that the set $T_0 = A \cup (A - 1) \cup \dots \cup (A - h)$ is thick. Since T_0 is thick, for each $n \in \mathbb{N}$ there exists $t_n \in \mathbb{N}$ such that

$$\{t_n + 1, \dots, t_n + n\} \subseteq T_0.$$

Define

$$T := \bigcup_{n \in \mathbb{N}} \{t_{2n} + 1, \dots, t_{2n} + n\}.$$

Then for every $m \in \mathbb{N}$, the translate $T_0 - m$ contains all but finitely many elements of T , because it contains $\bigcup_{n \geq m} \{t_{2n} + 1, \dots, t_{2n} + n\}$. This property will be used later in the proof.

We now construct a nested sequence of piecewise syndetic sets

$$B_0 \supseteq B_1 \supseteq B_2 \supseteq \dots$$

as follows. Set $B_0 = T$. Suppose B_{n-1} has already been defined. Since B_{n-1} is piecewise syndetic, by the partition regularity of piecewise syndetic sets (Corollary 26), at least one of

$$B_{n-1} \cap (A - n) \quad \text{or} \quad B_{n-1} \setminus (A - n)$$

is piecewise syndetic. Let B_n be whichever of the two is piecewise syndetic. This construction ensures that for each $n \in \mathbb{N}$, we have either

$$B_n \cap (A - n) = \emptyset \quad \text{or} \quad B_n \subseteq (A - n).$$

Define $L := \{n \in \mathbb{N} : B_n \subseteq (A - n)\}$. By construction, for any finite nonempty $F \subseteq L$, the

intersection

$$\bigcap_{m \in F} (A - m)$$

contains $B_{\max F}$, and is therefore piecewise syndetic.

It remains to show that L is syndetic. Fix $n \in \mathbb{N}$. Note that $T_0 - n$ contains all but finitely many elements of B_{n+h} , because $B_{n+h} \subseteq T$. Since $T_0 - n = (A - n) \cup (A - n - 1) \cup \dots \cup (A - n - h)$, there must exist some $j \in \{0, 1, \dots, h\}$ such that $B_{n+h} \cap (A - n - j) \neq \emptyset$. But $B_{n+h} \subseteq B_{n+j}$, so we must have $B_{n+j} \cap (A - n - j) \neq \emptyset$. By construction of the sequence $B_0 \subseteq B_1 \supseteq B_2 \supseteq \dots$, this forces $B_{n+j} \subseteq (A - n - j)$, i.e. $n + j \in L$. Since n was arbitrary, we conclude that

$$L \cup (L - 1) \cup \dots \cup (L - h) = \mathbb{N},$$

which shows that L is syndetic. □

From Proposition 27 we immediately obtain the following interesting corollary.

Corollary 28. *For any piecewise syndetic $A \subseteq \mathbb{N}$ there exist infinitely many $n \in \mathbb{N}$ such that $A \cap (A - n)$ is piecewise syndetic.*

2.3. van der Waerden's Theorem – equivalent forms

van der Waerden's Theorem is one of the key results in Combinatorial Number Theory.

van der Waerden's Theorem ([vdW28]). *For any $k \in \mathbb{N}$ and any finite coloring of \mathbb{N} there exists a monochromatic k -term arithmetic progression.*




Figure 2.1: An example of a 3-coloring of the set $\{1, \dots, 26\}$. Can you find a monochromatic arithmetic progression of length 3?

Proposition 29. *Fix $k \in \mathbb{N}$. The following are equivalent:*

- (i) *(van der Waerden's Theorem – infinitary version). For any finite coloring of \mathbb{N} there exists a monochromatic k -term arithmetic progression.*
- (ii) *(van der Waerden's Theorem – finitary version). For any $r \in \mathbb{N}$ there exists $W = W(r, k) \in \mathbb{N}$ such that if the set $\{1, \dots, W\}$ is colored using at most r colors then there exists a monochromatic k -term arithmetic progression in $\{1, \dots, W\}$.*
- (iii) *Any syndetic set $S \subseteq \mathbb{N}$ contains a k -term arithmetic progression.*

(iv) For any piecewise syndetic $A \subseteq \mathbb{N}$ there exists $d \in \mathbb{N}$ and a piecewise syndetic set $B \subseteq \mathbb{N}$ such that for all $b \in B$ we have $\{b + d, b + 2d, \dots, b + kd\} \subseteq A$.

Let us now provide a proof of Proposition 29.

Proof of (i) \iff (ii). This equivalence follows immediately from the Compactness Theorem for finite colorings (see Section 1.2) applied to the set $Y = \mathbb{N}$ and the family $\mathcal{F} = \{\{a, a + d, \dots, a + (k - 1)d\} : a, d \in \mathbb{N}\}$. \square

Proof of (i) \implies (iii). Let $S \subseteq \mathbb{N}$ be syndetic. By definition, this means there exists $h \in \mathbb{N}$ such that $S \cup (S - 1) \cup \dots \cup (S - h)$ covers \mathbb{N} . We can interpret this finite partitioning of \mathbb{N} as a finite coloring of \mathbb{N} using at most h colors. According to (i), one of the cells of the partition, say $S - j$, contains a k -term arithmetic progression. But if $S - j$ contains a k -term arithmetic progression then shifting this progression by j shows that S also contains a k -term arithmetic progression. \square

Proof of (iii) \implies (iv). Let $A \subseteq \mathbb{N}$ be piecewise syndetic. Using Proposition 27 we can find a syndetic set L such that for any finite, non-empty $F \subseteq L$ the intersection

$$\bigcap_{n \in F} (A - n) \tag{2.3.1}$$

is piecewise syndetic. According to part (iii), the syndetic set L contains a k -term arithmetic progression, i.e., there exist $a, d \in \mathbb{N}$ such that $\{a, a + d, \dots, a + (k - 1)d\} \subseteq L$. In view of (2.3.1), the set $B' = (A - a) \cap (A - a - d) \cap \dots \cap (A - a - (k - 1)d)$ is piecewise syndetic. This implies that the set

$$B = (A - d) \cap \dots \cap (A - kd)$$

is also piecewise syndetic, because $B = B' - d + a$. It is now easy to check that for all $b \in B$ we have $\{b + d, b + 2d, \dots, b + kd\} \subseteq A$ as desired. \square

Proof of (iv) \implies (i). If \mathbb{N} is colored using finitely many colors then, according to Corollary 26, there exists a monochromatic piecewise syndetic set. By part (iv), any piecewise syndetic set contains a k -term arithmetic progression. It follows that there exists a monochromatic k -term arithmetic progression. \square

The smallest possible number $W(r, k)$ in part (ii) of Proposition 29 is called the *van der Waerden number* for (r, k) . Below is a table of known van der Waerden numbers (or best known lower bounds):

The best known upper bound on van der Waerden numbers that holds for all $r, k \geq 2$ is

$$W(r, k) \leq 2^{2^{r-2^{k+9}}}$$

k/r	2 Colors	3 Colors	4 Colors	5 Colors	6 Colors
3 – Term	9	27	76	> 170	> 225
4 – Term	35	293	> 1,048	> 2,254	> 9,778
5 – Term	178	> 2,173	> 17,705	> 98,740	> 98,748
6 – Term	1132	> 11,191	> 91,331	> 540,025	> 816,981
7 – Term	> 3,703	> 48,811	> 420,217	> 2,941,519	> 20,590,633
8 – Term	> 11,495	> 238,400	> 2,388,317	> 16,718,219	> 117,027,533
9 – Term	> 41,265	> 932,745	> 10,898,729	> 79,706,009	> 557,942,063
10 – Term	> 103,474	> 4,173,724	> 76,049,218	> 542,694,970	> 3,798,864,790
11 – Term	> 193,941	> 18,603,731	> 329,263,781	> 3,621,901,591	> 39,840,917,501
12 – Term	> 638,727	> 79,134,144	> 1,536,435,264	> 16,900,787,904	> 185,908,666,944
13 – Term	> 1,642,309	> 251,282,317	> 5,683,410,589	> 73,884,37,657	> 960,496,389,541

2.4. Proof of van der Waerden's Theorem

Color Focusing Lemma. *Let $k \in \mathbb{N}$ and suppose van der Waerden's Theorem has already been proven for k . Then for any finite coloring of \mathbb{N} and any $r \in \mathbb{N}$ there exist monochromatic piecewise syndetic sets $A_0, A_1, \dots, A_r \subseteq \mathbb{N}$ such that for all $0 \leq i < j \leq r$ there exists $u \in \mathbb{N}$ with*

$$\{a + u, a + 2u, \dots, a + ku : a \in A_j\} \subseteq A_i. \quad (2.4.1)$$

Proof. We proceed by induction on r . It follows from Corollary 26 that there exists a monochromatic piecewise syndetic set $A_0 \subseteq \mathbb{N}$. If A_0, \dots, A_{r-1} have already been found then A_r is constructed as follows. According to part (iv) of Proposition 29, there exists a piecewise syndetic set $B \subseteq \mathbb{N}$ and some $d \in \mathbb{N}$ such that for all $b \in B$ we have $\{b + d, b + 2d, \dots, b + kd\} \subseteq A_{r-1}$. The finite coloring of \mathbb{N} induces a finite partition of B . Hence, using Corollary 26 once more, we can find a monochromatic piecewise syndetic set $A_r \subseteq B$. Thus

$$\{a + d, a + 2d, \dots, a + kd : a \in A_r\} \subseteq A_{r-1}. \quad (2.4.2)$$

Let $0 \leq i < j \leq r$. If $j < r$ then (2.4.1) follows from the induction hypothesis. If $j = r$ then we can first use the induction hypothesis to find some $\tilde{u} \in \mathbb{N}$ such that

$$\{a + \tilde{u}, a + 2\tilde{u}, \dots, a + k\tilde{u} : a \in A_{r-1}\} \subseteq A_i. \quad (2.4.3)$$

Then, taking $u = \tilde{u} + d$ and combining (2.4.2) and (2.4.3), we obtain $\{a + u, a + 2u, \dots, a + ku : a \in A_r\} \subseteq A_i$ as desired. \square

Proof of van der Waerden's Theorem. We proceed by induction on k . If $k = 2$ then van der Waerden's Theorem is trivial. So let us assume that $k \geq 2$ and that van der Waerden's Theorem has already been proven for k . We want to show that any finite coloring of \mathbb{N} admits a monochromatic $(k + 1)$ -term arithmetic progression.

Suppose \mathbb{N} is colored using m colors. By applying the Color Focusing Lemma with $r = m$ we can find monochromatic piecewise syndetic sets $A_0, A_1, \dots, A_m \subseteq \mathbb{N}$ such that for all $0 \leq i < j \leq m$ there exists $u \in \mathbb{N}$ with

$$\{a + u, a + 2u, \dots, a + ku : a \in A_j\} \subseteq A_i. \quad (2.4.4)$$

Figure 2.2: Since $N(3, 3) = 27$, there exists no 3-coloring of the set $\{1, \dots, 27\}$ without a monochromatic 3-term arithmetic progression. But there exist 48 distinct colorings of the set $\{1, \dots, 26\}$ without a monochromatic 3-term arithmetic progression. A complete list of these 48 colorings, denoted by p_1, \dots, p_{48} , is depicted above.

Since there are $m + 1$ sets A_0, A_1, \dots, A_m but only m colors, two of the sets must have the same color. In other words, there exist $0 \leq i < j \leq m$ such that A_i and A_j have the same color. Take any $u \in \mathbb{N}$ for which (2.4.4) is satisfied and take any $a \in A_j$. Then the $(k + 1)$ -term arithmetic progression $a, a + u, \dots, a + ku$ is monochromatic, finishing the proof. \square

2.5. Gallai's Theorem

What if instead of finitely coloring the positive integers \mathbb{N} as in Schur's Theorem or van der Waerden's Theorem, one colors the integer lattice points in the plane \mathbb{N}^2 . This begs the following natural question.

Question 30. Is it possible to find for any finite coloring of \mathbb{N}^2 a monochromatic square $(a, b), (a + h, b), (a, b + h), (a + h, b + h)$?

An affirmative answer to Question 30 is provided by Gallai's Theorem, which can be viewed as a higher-dimensional generalization of van der Waerden's Theorem. We need the following definition.




Figure 2.3: Can you find a monochromatic square in this two-coloring of the 14×14 grid?

Definition 31. Let $V, W \subseteq \mathbb{Z}^d$. We say that W is *homothetic* to V if V can be shifted and dilated to become W , i.e., there exist $\vec{u} \in \mathbb{Z}^d$ and $\lambda \in \mathbb{Z} \setminus \{0\}$ such that $W = \lambda V + \vec{u}$.




Figure 2.4: Two homothetic pyramids.

Gallai's Theorem. Let V be a finite subset of \mathbb{Z}^d . For any finite coloring of \mathbb{Z}^d there exists a monochromatic set of points homothetic to V .

We can reduce Gallai's Theorem to the following.

Theorem 32. For any finite coloring of \mathbb{Z}^d there exist $(a_1, \dots, a_d) \in \mathbb{Z}^d$ and $h \in \mathbb{N}$ such that the d -dimensional "cube"

$$\{(a_1 + \varepsilon_1 h, \dots, a_d + \varepsilon_d h) : \varepsilon_1, \dots, \varepsilon_d \in \{0, 1\}\}$$

is monochromatic.

Proof that Theorem 32 implies Gallai's Theorem. Let $V = \{\vec{v}_1, \dots, \vec{v}_r\}$ be a finite subset of \mathbb{Z}^d and suppose $\chi: \mathbb{Z}^d \rightarrow \{1, \dots, m\}$ is a coloring of \mathbb{Z}^d using at most m colors. Define a coloring $\tilde{\chi}: \mathbb{Z}^r \rightarrow \{1, \dots, m\}$ of \mathbb{Z}^r as

$$\tilde{\chi}(n_1, \dots, n_r) = \chi(n_1 \vec{v}_1 + \dots + n_r \vec{v}_r), \quad \forall (n_1, \dots, n_r) \in \mathbb{Z}^r.$$

By Theorem 32, there exist $(a_1, \dots, a_r) \in \mathbb{Z}^r$ and $h \in \mathbb{N}$ such that $\{(a_1 + \varepsilon_1 h, \dots, a_r + \varepsilon_r h) : \varepsilon_1, \dots, \varepsilon_r \in \{0, 1\}\}$ is monochromatic. Define

$$\vec{u} = a_1 \vec{v}_1 + \dots + a_r \vec{v}_r \quad \text{and} \quad \lambda = h.$$

Then the set $\lambda V + \vec{u}$ is homothetic to V and monochromatic with respect to the coloring χ . \square

Proof that Gallai's Theorem implies Theorem 32. Suppose $\chi: \mathbb{Z}^d \rightarrow \{1, \dots, m\}$ is a finite coloring of \mathbb{Z}^d . Let

$$H = \{(\varepsilon_1, \dots, \varepsilon_d) : \varepsilon_1, \dots, \varepsilon_d \in \{0, 1\}\}$$

denote the unit cube in \mathbb{Z}^d . By Gallai's Theorem, we can find a homothetic image of H that is monochromatic with respect to χ , finishing the proof. \square

The proof of Gallai's Theorem is omitted.

Bibliography

- [Ram30] F. P. RAMSEY, On a Problem of Formal Logic, *Proceedings of the London Mathematical Society* **s2-30** (1930), 264–286. <http://dx.doi.org/10.1112/plms/s2-30.1.264>.
- [Sch17] I. SCHUR, Über die Kongruenz $x^m + y^m \equiv z^m \pmod{p}$, *Jahresbericht der Deutschen Mathematiker-Vereinigung* **25** (1917), 114–116. Available at <http://eudml.org/doc/145475>.
- [vdW28] B. L. VAN DER WAERDEN, Beweis einer Baudetschen Vermutung, *Nieuw. Arch. Wisk.* **15** (1928), 212–216.