

### Exercises – week 13

**Exercise 1.** *Using a short exact sequence.*

Let  $\iota: D \rightarrow X$  be an effective Cartier divisor on an integral scheme  $X$ . As there is a short exact of sheaves

$$0 \rightarrow \mathcal{O}(-D) \rightarrow \mathcal{O}_X \rightarrow \iota_*\mathcal{O}_D \rightarrow 0.$$

In consequence there is a long exact sequence in cohomology,

$$(\dots) \rightarrow H^i(X, \mathcal{O}(-D)) \rightarrow H^i(X, \mathcal{O}_X) \rightarrow H^i(D, \mathcal{O}_D) \rightarrow (\dots)$$

Let  $k$  be a field. Consider  $\mathbb{P}_k^5 = \text{Proj}(k[x_0, \dots, x_5])$ . We consider the closed subscheme

$$X = V_+(x_0^2 + x_1x_2).$$

You can freely use that  $X$  is a Cartier divisor in  $\mathbb{P}_k^5$  with ideal sheaf isomorphic to  $\mathcal{O}_{\mathbb{P}_k^5}(-2)$ .

(1) Show that

$$H^i(X, \mathcal{O}_X) = 0$$

if  $i > 0$  and that  $H^0(X, \mathcal{O}_X) = k$ .

(2) Show that for  $1 \leq j \leq 3$  we have

$$H^i(X, \mathcal{O}_X(-j)) = 0$$

for all  $i \geq 0$ .

**Exercise 2.** *Stability properties of (very-)ample sheaves under tensor product.* Let  $X$  be a Noetherian scheme. Let  $\mathcal{L}$  and  $\mathcal{M}$  be invertible sheaves on  $X$ .

- (1) If  $\mathcal{L}$  is ample and  $\mathcal{M}$  is globally generated, show that  $\mathcal{L} \otimes \mathcal{M}$  is ample.
- (2) If  $\mathcal{L}$  is ample and  $\mathcal{M}$  is arbitrary, deduce that there is a  $n$  such that  $\mathcal{L}^n \otimes \mathcal{M}$  is ample.
- (3) Show that if  $\mathcal{L}$  and  $\mathcal{M}$  are ample, then  $\mathcal{L} \otimes \mathcal{M}$  is ample.

Now suppose that  $X$  is an  $A$ -scheme where  $A$  is a Noetherian ring.

- (4) If  $\mathcal{L}$  is  $A$ -very ample and  $\mathcal{M}$  is globally generated, then  $\mathcal{L} \otimes \mathcal{M}$  is  $A$ -very ample.
- (5) If  $\mathcal{L}$  is ample, then there is a  $n_0 > 0$  such that  $\mathcal{L}^n$  is  $A$ -very-ample for all  $n \geq n_0$ .

**Exercise 3.** *A Čech cohomology computation.* Let  $k$  be a field. Let  $U = \mathbb{A}_k^2 \setminus 0$ . Compute the cohomology of  $\mathcal{O}_U$  on  $U$ . After showing that  $\mathcal{O}_U$  is ample, deduce that Serre vanishing does not hold for  $U$ .

**Exercise 4.** *Curves in  $\mathbb{P}_k^2$ .* Let  $k$  be a field. Let  $C = V_+(F)$  for  $F \in \mathcal{O}_{\mathbb{P}_k^2}(d)(\mathbb{P}_k^2)$  for a  $d \geq 1$ .

- (1) Show that  $H^0(C, \mathcal{O}_C) \cong k$ .
- (2) Deduce that any  $C_1$  and  $C_2$  of the above form intersect.
- (3) Deduce that  $H^1(C, \mathcal{O}_C)$  is a  $k$ -vector space of dimension  $\frac{(d-1)(d-2)}{2}$  using the long-exact sequence from exercise 1.

**Remark.** We say that  $\frac{(d-1)(d-2)}{2}$  is the *arithmetic genus* of  $C$ . Curves of degree 3 have arithmetic genus 1. Smooth ones are called *elliptic curves*. Any smooth curve  $C$  over an algebraically closed field  $k$  with  $H^1(E, \mathcal{O}_E) = 1$  can be realized as a smooth cubic in  $\mathbb{P}_k^2$ , see for example Hartshorne III,4.6.

**Exercise 5.** *Blow-ups, revisited.* Let  $X$  be a scheme and  $\mathcal{I}$  a quasi-coherent ideal sheaf. We define the *blow-up* of  $X$  at  $\mathcal{I}$  to be

$$\pi: \text{Bl}_{\mathcal{I}}(X) = \underline{\text{Proj}}\left(\bigoplus_{n \geq 0} \mathcal{I}^n\right) \rightarrow X$$

where  $\underline{\text{Proj}}$  denotes the relative Proj of the graded  $\mathcal{O}_X$ -algebra  $\bigoplus_{n \geq 0} \mathcal{I}^n$ . Note that for every open affine  $\text{Spec}(A) \subset X$ , where  $\mathcal{I}$  corresponds to  $I$ , the pullback of the above is the blow-up  $\text{Proj}(\bigoplus_{n \geq 0} I^n)$ . The affine blow-up previously introduced in the exercises.

- (1) Show that  $\mathcal{O}(1)$  of this relative Proj is the ideal sheaf corresponding to the exceptional divisor (the pullback of  $V(\mathcal{I})$  along  $\pi$ ). Note then that by construction, the exceptional divisor of a blow-up is always Cartier. We denote  $\mathcal{O}(1)$  accordingly by  $\mathcal{O}(-E)$  below. We also recall that from the Proj construction we have a canonical surjective map of sheaves

$$\pi^* \mathcal{I} \rightarrow \mathcal{O}(1) = \mathcal{O}(-E).$$

- (2) *Resolving indeterminacies.* Suppose now that  $X$  is an  $S$ -scheme. Let  $\mathcal{L}$  be a line bundle on  $X$ . Let  $s_0, \dots, s_n \in \mathcal{L}(X)$  be global sections. Let

$$U = \bigcup_{i=0}^n D(s_i) \rightarrow \mathbb{P}_S^n$$

be the induced morphism.

- (a) *Base locus.* Show that there is a unique ideal sheaf  $\mathcal{I} \subset \mathcal{O}_X$  such that  $\mathcal{I}\mathcal{L}$  is the image of the natural map

$$\mathcal{O}_X^{\oplus(n+1)} \xrightarrow{(s_0, \dots, s_n)} \mathcal{L}.$$

The associated closed subscheme  $V(\mathcal{I})$  is denoted by  $V(s_0, \dots, s_n)$  and is called the *base locus* of  $s_0, \dots, s_n$ . Furthermore, realize (because  $\mathcal{L}$  is locally free), that the natural map  $\mathcal{I} \otimes \mathcal{L} \rightarrow \mathcal{I}\mathcal{L}$  is an isomorphism. Show also that

$$X \setminus V(\mathcal{I}) = \bigcup_{i=0}^n D(s_i).$$

- (b) *Resolving maps.* Show that you can extend the partially defined map  $X \dashrightarrow \mathbb{P}_S^n$  (totally defined on  $U$ ) to a map

$$\begin{array}{ccc} \mathrm{Bl}_{\mathcal{I}}(X) & & \\ \pi \downarrow & \searrow & \\ X & \dashrightarrow & \mathbb{P}_S^n \end{array}$$

*Hint: In this case, show that  $\mathcal{O}(-E) \otimes \pi^* \mathcal{L}$  has  $n + 1$  induced global generating sections  $\pi^* s_i$  as summarized below*

$$\pi^* \mathcal{O}_X^{\oplus(n+1)} \xrightarrow{(\pi^* s_0, \dots, \pi^* s_n)} \pi^*(\mathcal{I}\mathcal{L}) \cong \pi^* \mathcal{I} \otimes \pi^* \mathcal{L} \rightarrow \mathcal{O}(-E) \otimes \pi^* \mathcal{L}.$$

- (3) *Application to curves.* Let  $C$  be a regular curve over  $k$ . Let  $C \dashrightarrow \mathbb{P}_k^n$  be a partially defined map on a non-empty open set of  $C$ . Show that there is a unique extension of the map to  $C \rightarrow \mathbb{P}_k^n$ . *Hint: show that blowing-up a Cartier divisor is an isomorphism.*

**Exercise 6.** *An example of resolution.* Consider  $k$  an algebraically closed field and  $X = \mathbb{P}_k^2$ . Let  $x_0, x_1 \in \mathcal{O}_{\mathbb{P}_k^2}(1)$ . Then, it defines a partially defined map

$$U = \mathbb{P}_k^2 \setminus V_+(x_0, x_1) \rightarrow \mathbb{P}_k^1.$$

- (1) Show that on  $k$ -rational points, this map is to be understood as  $[\lambda_0 : \lambda_1 : \lambda_2] \mapsto [\lambda_0, \lambda_1]$ .

Consider the blow-up of  $\mathbb{P}_k^2$  at  $\mathcal{I}$  the ideal sheaf defining  $V_+(x_0, x_1) = [0 : 0 : 1]$  – this is the ideal sheaf of the base locus of  $x_0, x_1$ , see Exercise 5.b.a. We denote this blow-up by  $\pi: B \rightarrow \mathbb{P}_k^2$ .

- (2) Write  $\frac{x_0}{x_2} = x$  and  $\frac{x_1}{x_2} = y$ . Show that the exceptional divisor  $E$  of this blow-up is isomorphic to  $\mathbb{P}_k^1$ . More precisely, by computing the blow-up locally in

$$D_+(x_2) = \mathbb{A}_{x,y}^2,$$

understand that the exceptional divisor naturally identifies to the projective space of lines through the origin of  $\mathbb{A}_{x,y}^2$ .<sup>1</sup> In particular, any  $k$ -rational point of the exceptional correspond to such a line.

- (3) Using Exercise 5, consider the resolved map

$$B \rightarrow \mathbb{P}_k^1.$$

Show that a  $k$ -rational point of the exceptional corresponding to a line  $L$  through  $[0 : 0 : 1]$ , say given by  $[\mu \cdot \lambda_1 : \mu \cdot \lambda_2 : 1]$  for  $\mu \in k$  and  $(\lambda_1, \lambda_2) \in k^2 \setminus 0$  is sent to  $[\lambda_1 : \lambda_2]$  by this resolved map.<sup>2</sup>

*Hint: work locally on  $D_+(x_2)$ . Then understand explicitly the map on standard blow-up charts. Another method is to show that the intersection of  $E$  with the closure of  $\pi^{-1}(L \setminus [0 : 0 : 1])$  in  $B$  is*

<sup>1</sup>Note also that the  $\mathbb{P}_k^1$  where we project is naturally understood as  $\mathbb{A}_{x,y}^2 \setminus 0 / \mathbb{G}_{m,k}$ .

<sup>2</sup>One could summarize as follows: the line in the exceptional is sent to itself by the resolved map. This is explained by the fact that the resolved map from the blow-up of  $\mathbb{A}_k^2$  at 0 to  $\mathbb{P}_k^1$  (resolving the quotient map  $\mathbb{A}_k^2 \setminus 0 \rightarrow \mathbb{P}_k^1$ ) is equal to the tautological line bundle of  $\mathbb{P}_k^1$ .

*precisely the point corresponding to  $L$ . Then, because the projection is constant along  $L$ , conclude.*