

Most flows are unstable...


1. Intro-definitions
2. Rayleigh-Taylor
3. Rayleigh Plateau (destabilization through surface tension)
4. Rayleigh-Benard (convection)
5. Benard-Marangoni
6. Taylor Couette-Centrifugal instability
7. Kelvin-Helmholtz
8. Inflection point theorem Rayleigh, Orr sommerfeld
9. Orr sommerfeld, transient growth
10. Spatial growth

SPATIALLY DEVELOPING SHEAR FLOWS


Flat plate boundary layer



Mixing layer




Cylinder wake



Inviscid instability
(inflection point criterion)
Stabilized by viscosity
Transient growth on top

Plane channel flow




Inviscid stability
(inflection point criterion)
Destabilized by viscosity
Transient growth important


2D jet



Hagen Poiseuille




Couette



Linear stability
Transient growth essential

Viscosity has limited stabilizing influence on K-H instability




$$U_B(y) = \tanh y$$

$$Re := \Delta U \delta / \nu$$

$$\omega_{i,max} \frac{\delta}{\Delta U} \approx \sqrt{\frac{0,2}{1 + \text{constant}/0,2Re}}$$


Viscosity has stabilizing influence on K-H instability



PARALLEL FLOW CONCEPTS

Viscous instabilities

Hyperbolic tangent mixing layer



2D PARALLEL FLOW CONCEPTS

Dispersion relation

2D vorticity equation

$$\left(\frac{\partial}{\partial t} + \frac{\partial \Psi}{\partial y} \frac{\partial}{\partial x} - \frac{\partial \Psi}{\partial x} \frac{\partial}{\partial y} \right) \nabla^2 \Psi = \frac{1}{Re} \nabla^4 \Psi$$

Basic flow + perturbation

$$\Psi(x, t) = \int U(y) dy + \psi(x, y, t)$$

Linear vorticity equation

$$\left(\frac{\partial}{\partial t} + U(y) \frac{\partial}{\partial x} \right) \nabla^2 \psi - U''(y) \frac{\partial \psi}{\partial x} = \frac{1}{Re} \nabla^4 \psi$$

Dispersion relation

$$D(k, \omega) = 0$$

Temporal approach:

k is real; ω is complex

Perturbation grow and decay in time!

Spatial approach:

ω is real; k is complex

Perturbations grow and decay in space!

Shear layer is inviscidly unstable!


Hyperbolic tangent mixing layer

$$U(y) = \bar{U} + \frac{\Delta U}{2} \tanh\left(\frac{2y}{\delta_\omega}\right)$$

$$\delta_\omega(x) \equiv \frac{(U_1 - U_2)}{(dU/dy)_{\max}}$$


Velocity ratio

$$R \equiv \frac{U_1 - U_2}{U_1 + U_2} = \frac{\Delta U}{2\bar{U}}$$



Why?

Only a necessary condition for instability!
Remember: Influence of confinement



$$R = 1$$

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

$$U(y; R) = 1 + R \tanh y$$


$$U_1(y) = \tanh y$$

Dispersion relation


$$\omega(k; R) = k + R \omega_1(k)$$

PARALLEL FLOW CONCEPTS


Effect of velocity ratio




$R = 0$



$0 < R < 1$



$R = 1$



$R > 1$

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Temporal approach

$$\omega_1(k) = i \omega_{1,i}(k)$$

$$\omega_i(k; R) = R \omega_{1,i}(k)$$

$$c_r = \omega_r / k = 1$$

Temporal approach: k is real; ω is complex

2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Spatial approach

$$k + R \omega_1(k) = \omega$$

$$R \ll 1$$

$$-k_i(\omega, R) \sim R \omega_{1,i}(\omega)$$

Gaster transformation

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer


$$U(y) = \begin{cases} U_1, & y > \delta_\omega/2 \\ (U_1 + U_2)/2 + (U_1 - U_2)y/\delta_\omega, & |y| < \delta_\omega/2 \\ U_2, & y < -\delta_\omega/2 \end{cases}$$

$$\phi'' - k^2 \phi = 0$$

$$\phi_1(y) = A_1 e^{-ky}, \quad y > \delta_\omega/2,$$

$$\phi_2(y) = B_2 e^{ky}, \quad y < -\delta_\omega/2,$$

$$\phi_0(y) = A_0 e^{-ky} + B_0 e^{ky}, \quad |y| < \delta_\omega/2$$



2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$A_1 e^{-k\delta_\omega/2} = A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2},$$

$$B_2 e^{-k\delta_\omega/2} = A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2},$$

$$\begin{aligned} -k(U_1 - c)A_1 e^{-k\delta_\omega/2} &= k(U_1 - c)(-A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2}) \\ &\quad - \frac{\Delta U}{\delta_\omega}(A_0 e^{-k\delta_\omega/2} + B_0 e^{k\delta_\omega/2}), \end{aligned}$$

$$\begin{aligned} k(U_2 - c)B_2 e^{-k\delta_\omega/2} &= k(U_2 - c)(-A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2}) \\ &\quad - \frac{\Delta U}{\delta_\omega}(A_0 e^{k\delta_\omega/2} + B_0 e^{-k\delta_\omega/2}). \end{aligned}$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$-\frac{\Delta U}{\delta_\omega} A_0 e^{-k\delta_\omega/2} + \left[2k(U_1 - c) - \frac{\Delta U}{\delta_\omega} \right] B_0 e^{k\delta_\omega/2} = 0$$
$$\left[2k(U_2 - c) + \frac{\Delta U}{\delta_\omega} \right] A_0 e^{k\delta_\omega/2} + \frac{\Delta U}{\delta_\omega} B_0 e^{-k\delta_\omega/2} = 0$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer

$$4(k\delta_\omega)^2(c - \bar{U})^2 - \left[(k\delta_\omega - 1)^2 - e^{-2k\delta_\omega} \right] \Delta U^2 = 0$$

$$k\delta_\omega \mapsto 2k, c/\bar{U} \mapsto c$$

$$4k^2(c - 1)^2 - R^2 \left[(2k - 1)^2 - e^{-4k} \right] = 0$$



$$c \equiv \frac{\omega}{k} = 1 \pm \frac{R}{2k} \left[(2k - 1)^2 - e^{-4k} \right]^{1/2}$$

2D PARALLEL FLOW CONCEPTS

Broken-line profile mixing layer



Temporal approach

$$2k_n - 1 = e^{-2k_n}$$



Spatial approach



$$R = 0.5$$





2D PARALLEL FLOW CONCEPTS

Hyperbolic tangent mixing layer

Temporal



Spatial



Michalke (1964, 65)

Solving a spatial instability problem

ex: Rayleigh equation

Back to temporal stability analysis!

How to solve Rayleigh equation for real k and complex ω ?

We fix k , we need to find all ω and ψ such that

$$k \left(U \left(\frac{d^2}{dy^2} - k^2 \right) - U''(y) \right) \psi = \omega \left(\frac{d^2}{dy^2} - k^2 \right) \psi$$
$$\psi(-L) = \psi(L) = 0$$

Formally,

$$\mathcal{A}\psi = c\mathcal{E}\psi$$

$$c = \omega/k$$


Discretize

$$\mathbf{A}\Psi = c\mathbf{E}\Psi$$

Generalized eigenvalue problem

How to solve Rayleigh equation for real k and complex ω ?

Finite differences of order 1



$$\Psi = \begin{pmatrix} \psi(y_1) \\ \psi(y_2) \\ \vdots \\ \psi(y_N) \\ \psi(y_{N+1}) \end{pmatrix} = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \vdots \\ \psi_N \\ \psi_{N+1} \end{pmatrix} \quad \Psi'' = \begin{pmatrix} \psi''(y_1) \\ \psi''(y_2) \\ \vdots \\ \psi''(y_N) \\ \psi''(y_{N+1}) \end{pmatrix}$$

How to solve Rayleigh equation for real k and complex ω ?

Finite differences

$$\begin{pmatrix} \psi_2'' \\ \psi_3'' \\ \psi_4'' \\ \vdots \\ \psi_{N-3}'' \\ \psi_{N-2}'' \\ \psi_{N-1}'' \end{pmatrix} = \begin{pmatrix} -2 & 1 & 0 & \dots & \dots & \dots & 0 \\ 1 & -2 & 1 & 0 & \dots & \dots & 0 \\ 0 & 1 & -2 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 & -2 & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 & -2 & 1 \\ 0 & \dots & \dots & \dots & 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} \psi_2 \\ \psi_3 \\ \psi_4 \\ \vdots \\ \psi_{N-3} \\ \psi_{N-2} \\ \psi_{N-1} \end{pmatrix}$$

Sparse matrix but low order!

How to solve Rayleigh equation for complex k and real ω ?

We fix ω , we need to find all k and ψ such that

$$k \left(U \left(\frac{d^2}{dy^2} - k^2 \right) - U''(y) \right) \psi = \omega \left(\frac{d^2}{dy^2} - k^2 \right) \psi$$
$$\psi(-L) = \psi(L) = 0$$


Formally,

$$(A_0(\omega, y) + kA_1(\omega, y) + k^2A_2(\omega, y) + k^3A_3(\omega, y)) \psi = 0$$

Polynomial eigenvalue problem

Many more eigenvalues (for Rayleigh equation: 3 x more!)

$$U=1+0.9*\tanh(y); \omega=0.4; L=5$$



Which of these waves are unstable?


$\text{Im}(k) < 0?$

$\text{Im}(k) > 0?$

Recall : $\exp(i(kx - \omega t))$

The stability of a spatial wave can be only determined if one knows in which direction it propagates!

k^+ waves propagate towards positive x




k^- waves propagate towards negative x



However, determining this direction of propagation is particularly difficult, except in the case of a temporally stable flow.

The addition of a positive imaginary offset to the frequency makes the temporal problem stable!

This separates the spatial waves into k^+ and k^- waves.




offset spatial stability

spatial stability: $\text{Im}(\omega)=0$

The branches are then followed by continuity

Validity of Gaster transformation?

$R=0.4$




$$R \ll 1$$

$$-k_i(\omega, R) \sim R\omega_{1,i}(\omega)$$

Validity of Gaster transformation?

R=0.9

x: Gaster transformation



$$-k_i(\omega, R) \sim R\omega_{1,i}(\omega)$$

Neutral curve

Schmid and Henningson 2002

