


COMPOSANTS SEMI-CONDUCTEURS

IV) Solutions S4

P.A. Besse

EPFL


E 4.2: Frog Oocyte: planar model



C. Massin et al., "Planar microcoil-based magnetic resonance imaging of cells",
 Proceedings Transducer's 2003, pp. 967-970

E 4.2: Start situation: all channels closed

Pompe par ATPase



Concentrations when all the channels are closed.

Constant concentrations in each region


	Oocyte	Outside	V_{nernst}
Na+	10 mM	90 mM	+ 56 mV
K+	90 mM	10 mM	- 56 mV
Cl- and anions-	100 mM	100 mM	0 mV

$$1 \text{ mM} \equiv 10^{-3} \text{ mole / liter} = 10^{-3} \cdot N_A \cdot \frac{\text{molecule}}{1000 \text{ cm}^3} = 6 \cdot 10^{17} \text{ molecule / cm}^3$$

Nernst voltage:
$$V_{\text{nernst}} \equiv \frac{RT}{F} \text{Ln} \left(\frac{[C_{\text{out}}]}{[C_{\text{in}}]} \right)$$

$R \equiv k \cdot N_A$
 $F \equiv |q_i| \cdot N_A$

K-channels open



$$V_{bi} = -\frac{kT}{q} \text{Ln} \left(\frac{[K^+]_{out}}{[K^+]_{in}} \right) = + 56 \text{ mV}$$

- Tension de Nernst

4 unknown: E, [K+], [Na+] and [Cl-]

4 equations:


Poisson:
$$\Delta\varphi = -\frac{1}{\epsilon_0\epsilon} \rho = -\frac{1}{\epsilon_0\epsilon} q \cdot ([Na^+] + [K^+] - [Cl^-])$$

$$\vec{E} = -\overrightarrow{\text{grad}}\varphi$$

Continuity:

$\frac{\partial [Na^+]}{\partial t} = \overset{\text{ATPase}}{G_{Na^+} - R_{Na^+}} - \frac{1}{q} \text{div}(\vec{j}_{Na^+})$	<div style="display: flex; justify-content: space-around; font-weight: bold; color: red; font-size: 0.8em;"> Drift Diffusion </div> $\vec{j}_{Na^+} = [Na^+] q \mu_{Na^+} \cdot \vec{E} - kT \mu_{Na^+} \overrightarrow{\text{grad}}[Na^+]$
$\frac{\partial [K^+]}{\partial t} = \overset{\text{ATPase}}{G_{K^+} - R_{K^+}} - \frac{1}{q} \text{div}(\vec{j}_{K^+})$	$\vec{j}_{K^+} = [K^+] q \mu_{K^+} \cdot \vec{E} - kT \mu_{K^+} \overrightarrow{\text{grad}}[K^+]$
$\frac{\partial [Cl^-]}{\partial t} = \overset{=0}{G_{Cl^-}} - \overset{=0}{R_{Cl^-}} + \frac{1}{q} \text{div}(\vec{j}_{Cl^-})$	$\vec{j}_{Cl^-} = [Cl^-] q \mu_{Cl^-} \cdot \vec{E} + kT \mu_{Cl^-} \overrightarrow{\text{grad}}[Cl^-]$

K and Na channels open



$$\frac{\mu_{Na^+}}{\mu_{K^+}} = 10$$



$$V_{bi} = -\frac{kT}{q} \ln \left(\frac{[K^+]_{out} + \frac{\mu_{Na^+}}{\mu_{K^+}} [Na^+]_{out}}{[K^+]_{in} + \frac{\mu_{Na^+}}{\mu_{K^+}} [Na^+]_{in}} \right) = -40 \text{ mV}$$

- Tension de Nernst

Na channels opening and closing


$$\frac{\mu_{Na^+}}{\mu_{K^+}} = 10$$

$$V_{bi} = -\frac{kT}{q} \text{Ln} \left(\frac{[K^+]_{out}}{[K^+]_{in}} \right) = 56 \text{ mV}$$




$$V_{bi} = -\frac{kT}{q} \text{Ln} \left(\frac{\mu_{K^+} [K^+]_{out} + \mu_{Na^+} [Na^+]_{out}}{\mu_{K^+} [K^+]_{in} + \mu_{Na^+} [Na^+]_{in}} \right) = -40 \text{ mV}$$

Na⁺ channels closed




Na⁺ channels open



Electric rays

about 500 cells in series in electric rays (*Torpedinidae*)




Y. Tanaka et al. «An electric generator using living Torpedo electric organs
Controlled by fluid pressure-based alternative nervous systems»


www.nature.com/scientificreports , May 2016

Source de tension «gels ioniques»

Gels séparés




Gels en contact (pression mécanique)





T. B. H. Schroeder, et al., «An electric-eel-inspired soft power source from stacked hydrogels»,
Nature, Vol. 552, p. 214-218, 14 dec. 2017

Source de tension «gels ioniques»



2449 gels (612 cells de 180 mV)
→ 110 V



T. B. H. Schroeder, et al., «An electric-eel-inspired soft power source from stacked hydrogels», Nature, Vol. 552, p. 214-218, 14 dec. 2017