









d. (2-(butan-2-yl)-1,7-dihydroxyoctan-3-yl) 2-hydroxy-3-methylbutanamide

e. (3,3,6-trichlorohex-4-yn-2-yl)(*E*)-3-(4-hydroxy-3-methoxyphenyl)prop-2-enoate


f. 1-benzyl-5-phenoxy-3-phenyl-2-(propan-2-yl)benzene

2. Give the IUPAC names for the following compounds. Include the isomerism of the double bond where necessary.





3. This question is about L-ascorbic acid, known as vitamin C (structure shown below).




a. Mark all chiral atoms with an asterisk and determine their absolute configurations.


b. Draw the enantiomer of ascorbic acid.

c. Draw all diastereomers of ascorbic acid.

d. Even though ascorbic acid does not contain carboxylic functional groups, it is an acidic compound. The proton of the hydroxylic group labeled in red is the most acidic proton in this structure. Using the resonance structures of the anion obtained by deprotonation of ascorbic acid explain why this proton is acidic.



4. Give the exact structural relationship (same compounds, different compounds, isomers – specify which type of isomerism) between the compounds in the following pairs:



### Reading Suggestions:

Clayden, Greeves, Warren, *Oxford University Press*, **2012**.

Organic Chemistry, John McMurry, *Thomson Brooks/Cole*, **2008**.

Chimie Organique, Les Grands Principes, John McMurry, *Dunod Editeur*, **2009**.

Chimie Organique, Paul Arnaud, *Dunod Editeur*, **2009**.