


# General Physics: Mechanics




**EPFL**


Swiss  
Plasma  
Center

**PHYS-101(en)**  
Lecture 10b: Collisions

Dr. Marcelo Baquero  
[marcelo.baquero@epfl.ch](mailto:marcelo.baquero@epfl.ch)  
November 18<sup>th</sup>, 2025




# 2D elastic collisions, same mass



We saw yesterday that

$$0 = 2 v_{Af} v_{Bf} \cos(\theta_A + \theta_B)$$

# 2D elastic collisions, same mass



- To solve for the 4 unknowns ( $v_{Af}$ ,  $v_{Bf}$ ,  $\theta_A$ ,  $\theta_B$ ), we need a fourth equation
- Knowing the “impact parameter”  $d$ , allows you to directly determine  $\theta_B$

# Conceptual question

---

Cart A is at rest. An identical cart B is moving to the right and collides **elastically** with cart A. After the collision, which of the following is true

- A. Carts A and B are both at rest.
- B. Cart B stops and cart A moves to the right with speed equal to the original speed of cart B.
- C. Cart A remains at rest and cart B bounces back with speed equal to its original speed.
- D. Cart A moves to the right with a speed slightly less than the original speed of cart B and cart B moves to the right with a very small speed.

# Conceptual question


---

Cart A is at rest. An identical cart B, moving to the right, collides **inelastically** with cart A. They stick together. After the collision, which of the following is true.

- A. Carts A and B are both at rest.
- B. Carts A and B move to the right with a speed less than cart B's original speed.
- C. Carts A and B moves to the right with speed greater than Cart B's original speed.
- D. Cart B stops and cart A moves to the right with speed equal to the original speed of cart B.

# Example: Fusion reactions


In a fusion reactor, isotopes of hydrogen are confined as they are heated to temperatures exceeding those of the sun (i.e. 100 million Celsius = 0.01 MeV) in order to enable the fusion reaction



This reaction is inelastic as it releases  $Q = 17.6$  MeV of energy, which is carried away in the form of the kinetic energy of the products. If the  ${}^3\text{H}$  particle is at rest and the  ${}^2\text{H}$  particle has 0.01 MeV of kinetic energy, how much energy does the neutron end up with?

Hints:

- You can still write down a version of conservation of energy, but you need to consider all the types of energy involved
- Look for an approximation that simplifies the problem significantly and barely affects the final answer



# Example: Fusion reactions

---

# Conceptual question

---

An explosion splits an object initially at rest into two pieces of unequal mass. Which piece has the greater kinetic energy?

- A. The less massive piece
- B. The more massive piece
- C. They both have the same kinetic energy
- D. There is not enough information to tell