

# Plasma Physics I

Series 6 (October 15, 2025)

Prof. Christian Theiler

*Swiss Plasma Center (SPC)*

*École Polytechnique Fédérale de Lausanne (EPFL)*

## Exercise 1

In the lecture we have neglected the effect of collisions in the two-fluid model used to derive the dispersion relation of a wave in a magnetised plasma.

Consider a cold unmagnetized fluid plasma ( $T = 0$ ,  $\mathbf{B}_0 = 0$ ).

- a.) Derive the dispersion relation of waves in such plasma keeping the collision term in the momentum equation for the electrons.
- b.) Show that in this case longitudinal waves (*Langmuir* waves) are damped.

## Exercise 2

An antenna can detect frequencies around  $f = 80$  MHz and is used to measure the wave coming from a pulsar producing a broad electromagnetic spectrum.

Due to the dispersion of the group velocity caused by the interstellar plasma, the measured frequency during a pulse drift varies according to  $df/dt = -5$  MHz  $\cdot$  s $^{-1}$ .

- a.) Considering  $\omega^2 \gg \omega_p^2$  and neglecting the magnetic field in the interstellar plasma, demonstrate that:

$$\frac{df}{dt} \approx -\frac{c}{x} \frac{f^3}{f_p^2}$$

where  $f_p = \omega_p/2\pi$  and  $x$  is the distance of the pulsar.

- b.) Find the distance of the pulsar in *parsec* (1 parsec =  $3 \times 10^{16}$  m) considering a mean electron density in space of  $2 \times 10^6$  m $^{-3}$ .