

# Quantum Field Theory

## Set 3

### Exercise 1: Classical Electromagnetism

Consider the classical electromagnetic fields  $\vec{E}(\vec{x}, t)$ ,  $\vec{B}(\vec{x}, t)$ .

- Write the Maxwell equations in presence of external an source.

Define the *field strength*  $F_{\mu\nu}$  as the  $4 \times 4$  matrix

$$F_{\mu\nu} = \begin{bmatrix} 0 & E^1 & E^2 & E^3 \\ -E^1 & 0 & -B^3 & B^2 \\ -E^2 & B^3 & 0 & -B^1 \\ -E^3 & -B^2 & B^1 & 0 \end{bmatrix} \quad F^{\mu\nu} = \begin{bmatrix} 0 & -E^1 & -E^2 & -E^3 \\ E^1 & 0 & -B^3 & B^2 \\ E^2 & B^3 & 0 & -B^1 \\ E^3 & -B^2 & B^1 & 0 \end{bmatrix}$$

- Recalling the definitions of the field in term of the vector potential  $A_\mu = (A_0, A_i)$  show that

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

- Introduce the *Lagrangian density* of the electromagnetic field in the presence of an external source:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - A_\mu J^\mu$$

where  $J^\mu$  is the current associated to the source. .

Find the Euler Lagrange equations and show that they correspond to the inhomogeneous Maxwell equations.

- Show that the additional two Maxwell equations follow from the *Bianchi identity*

$$\epsilon_{\mu\nu\rho\sigma}\partial^\mu F^{\rho\sigma} = 0$$

- Find the static solution (independent of time)  $A_\mu(\vec{x})$  for the current corresponding to a particle at rest  $J^\mu = (e\delta^3(\vec{x} - \vec{x}_0), \vec{0})$ .

### Exercise 2: Affine transformations

Consider the two parameter Lie group acting on the real numbers defined by the transformation

$$U(\alpha, \beta) : x \rightarrow x' = e^\alpha x + \beta$$

for any  $(\alpha, \beta) \in \mathbb{R}^2$ .

Show that this transformation defines a group. Is it an abelian group?

### Exercise 3: Dilation symmetry

Consider a particle with position  $q(t)$  and mass  $m$  obeying the differential equation

$$m\ddot{q}(t) + kq^2(t) = 0$$

with  $k$  a constant. What is the dimension of  $k$ ?

Find the value of the constant  $p$  such that the equation is symmetric under the dilation transformation

$$q(t) \rightarrow q'(t') = \lambda^{-p}q(t), \quad t' = \lambda t$$

Could you have guessed the result from the dimension of  $k$ ?

#### Exercise 4 [optional]

The Large Hadron Collider (LHC) started its operations on 10th September 2008, completing the first entire revolution of the 27 km long ring at 10:28 a.m. The proton beam coming from the SPS has been injected inside the LHC with an energy of 450 GeV per proton. Compute the velocity of the beam in units of  $c$  and the Lorentz  $\gamma$  factor.

At full performance at LHC, proton beams circulate in opposite directions with an energy of 7 TeV each. There are approximately  $2.8 \times 10^{14}$  protons per beam. Find the velocity in km/h of a running TGV with mass 400 tons, so that its kinetic energy is the same of that of a proton beam in LHC.

Suppose that the LHC, instead of being a collider with two proton beams circulating in opposite directions with an energy of 7 TeV each, were a collider with fixed target and an incoming proton beam with an energy  $E_a$  in the laboratory frame. What should  $E_a$  be in order to have the same energy of LHC in the center of mass frame?

Given that the power lost by a charged particle in a circular trajectory of radius  $r$  is given by

$$P = \frac{2e^2}{3r^2}(\beta\gamma)^4$$

and that we can give approximately 1 GeV of energy per particle every lap, what is the minimum radius of this accelerator to reach the center of mass energy of 1.5 TeV?