




Degenerative disease of the hip and knee, THA and TKA

Dr. Antoniadis

Unil
UNIL | Université de Lausanne



Terminology



The hip



The hip joint

- Deep joint
- Force and stability
- Weight bearing articulation




The hip joint:

- Proximal femur

Femoral head

Small
trochanter

Great
trochanter



The hip joint:


Acetabulum (etymology)



From acēt(um) (“vinegar”) + -bulum (“a vessel for”)



The hip joint:

Acetabular cavity




The labrum


- Ring of fibrocartilage
- deepens cavity
- Suction effect



The hip articulation





The hip



Anteversion of the femoral neck



The hip:

stability




The hip:

stability



The hip:

mobility




The knee



The Knee

- superficial joint
- stability
- weight bearing articulation



The knee:

Articular surfaces


- Distal femur
 - Femoral condyles



The knee:

Articular surfaces


- Proximal tibial
 - Tibial condyles



The knee:

Articular surfaces


- Proximal tibial
 - Tibial condyles



The knee:



Articular surfaces

- Femoro-tibial incogruency




The knee:

Articular surfaces




The knee:

Stability antero-posterior




- capsule



- Patellar ligament

Le Genou:

Stability antero-posterieure




- Lig. cruciate anterior


- Lig. cruciate posterior

The Knee:

Stability frontal




Ligt collatéral latéral



Ligt collatéral médial


The Knee:

Stability frontal





The knee:

mobility



The knee:

mobility




Osteoarthritis

- most frequent musculoskeletal disease
- high morbidity in developed countries
- second cause of invalidity after cardiovascular diseases





The articular cartilage

= hyaline cartilage



Normal

Osteoarthritis



Osteoarthritis :

definition macroscopic & histologic



- Focal alterations of the hyaline cartilage

fissures

erosions



subchondral bone



Osteoarthritis :

definition macroscopic and histologic

- Focal modifications of the articular cartilage

fissures

erosions





Exposition of the subchondral bone



Osteoarthritis :

definition macroscopic & histologic


- Modification of adjacent structures



Osteoarthritis :

Definition macroscopic and histologic

- + modifications of adjacent structures:
 - Subcondral bone
 - Synovia



The articulation is affected globally

Osteoarthritis (OA) : classification

- Topographie : Coxarthrosis (hip)
 Gonarthrosis (knee)

All the articulations can be affected by OA!

+/-

- Causal reasons:
 - primary OA
 - secondary OA

Osteoarthritis of the hip

COXARTHROSIS

- The most common reason of a chronic coxodynia (>6 mois)
- Disease of the articulation:
 - Wear of the articulation by progressive destruction of the cartilage
 - secondarily, unspecific synovial reaction and alteration of the subchondral bone


 *Bone remodeling on the weight-bearing zone*

Primary OA of the hip

- Estimated prevalence: 5 - 10 % >55 ans
 - mean age: 61 ans
 - bilateral: 40 —50%

COXARTHROSIS :

- Biomechanical factors
 - Obesity
 - Conflit fémoro-acétabulaire



COXARTHROSIS :

- Biomechanical factors
 - Professional and athletic activity



COXARTHROSIS

- Clinical signs :
 - mechanical pain (typically anterior)
 - functional impairment:
 - limping
 - restriction of mobility

NO direct radio-clinical correlation


COXARTHROSIS : diagnostic

- The most common symptoms include:
 - pain
 - stifness
 - age > 40 ans
 - onset progressiv
 - mechanic character

Diagnostic radiologic :

• *Signes Rx:*


1. Joint space narrowing
2. sclerosis
3. Bone cysts
4. osteophytes



Diagnostic radiologic :

• *Signes Rx:*

1. Joint space narrowing
2. sclerosis
3. Bone cysts
4. osteophytes



Analysis Rx : Dg etiologique coxahrosis secondary

- Coxometrie : Hip dysplasia ?

Other secondary reasons:

- Avascular necrosis
- Post-traumatique
- Sequels of ostéochondrite
- Sequels of epiphysiolysis
- Sequels of septic arthritis

Analyse Rx : coxometrie

Femoral neck-shaft angle :

Caput-collum-diaphyseal **angle** (CCD angle)

The **angle** between the longitudinal axes of the **femoral** neck and shaft


Normal values of CCD:

- 125-130°




$125^{\circ} < N < 135^{\circ}$

Analyse Rx : coxometrie



$< 125^\circ$ COXAVARA



$> 135^\circ$ = COXAVALGA


Analyse Rx : coxometrie

Coverage of the femoral head:

Angle between the center of femoral head and lateral acetabular border

Wiberg or CE (center-edge) angle


Normal value: 25-35°



Analysis Rx : coxometrie

Anterior coverage angle:

False profile view of the hip



- angle between the center of the femoral head and the anterior-most aspect of the acetabulum




Analysis Rx : coxometrie



Acetabular index (Tönnis angle)

radiographic measurement of femoral head bony coverage by the acetabulum


- decreased: $<3^{\circ}$
 - in severe overcoverage, there may be a negative value
- normal range: $3-13^{\circ}$ ²
- increased: $>13^{\circ}$
 - in acetabular dysplasia






Coxometrie : structural anomaly ?



Anomalie architecturale = coxarthrose secondaire



Anomalie architecturale = coxaûhrose secondaire



COXARTHROSIS : Traitement

- Therapeutic strategy:

Is the cause of coxarthrosis correctable ?:

- *dysplasia or congenital dislocation*

Hip conservative surgery:

âge <45 ans, optimal conditions


COXARTHROSIS : Traitement

**Peri-acetabular
osteotomy (PAO)**



COXARTHROSIS : Traitement

FAI-surgery



Therapeutic options

Total hip arthroplasty



THA



« The operation of the century: total hip replacement »

Learmonth et al. *The Lancet*, volume 370, Issue 9597, Oct 2007

History



Wiles' acetabular cup and resurface of head of the femur, 1938



Smith-Petersen mould, 1938




Judet brothers' acrylic femur head-replacement, late 1940s



Moore intramedullary endoprosthesis, 1950s –



Charnley low-friction arthroplasty, 1958 –



Exeter total hip replacement, 1970 –





History



McMinn Birmingham resurfacing, 1991 –



History



Modern THA



PTH Basic principles



Surgical approach


Fixation modes

Choice of implants
depending on different
conditions:




- the indications
- The age





Surgical approaches



EXEMPLE D'ABORD TRANSGLUTEAL




Installation en décubitus latéral



Posterior approach

Exposition de la hanche par voie postérieure

Ouvrir la capsule pour exposer la hanche





Luxation de la hanche en rotation interne

Section de la tête à la base du col


Préparation de la cavité fémorale et du cotyle

Direct anterior approach

- Pat in supine position
- with or without traction table
- true intermuscular plane



Choice of implants



Choice of implants



Choice of implants



Metal on Polyethylene




Metal on Metal




Ceramic on Ceramic


Double-mobility




Preoperative planification




Preoperative planification




Cup Implantation





Reaming



cement mantel



Acetabular preparation



Cup implantation without cement

Implantation of the femoral stem













Preparation of the femoral cavity
using tailored rasps

Press-fit impaction


Or cement introduction

The prosthetic head is impacted on
the cone





Postoperative x-ray



Osteoarthritis of the knee

GONARTHROSIS : classification

- Topographie: OA
 - 1, 2 ou 3 compartiments :
 - Medial OA
 - Lateral OA
 - Patellofem. OA
- +/- causal factors:
 - primary OA
 - Secondary OA




Osteoarthritis of the knee

- Knee OA is primarily provoked by mechanical factors:
 - the deformation femoro-tibial
 - modifications of the articular surface
 - sequels of traumatisme
 - post meniscectomie
 - ligamentous ruptures(LCA)

Osteoarthritis of the knee

- deformation of the lower limb (morphotype)



Medial osteoarthritis (varus knee)



- The deviation of the axis leads to an accelerated wear
- The wear of the cartilage worsens the deformation

Lateral osteoarthritis (valgus knee)



5-10%

The major deformation is mostly on the femur

KNEE OA

- Symptoms:
 - pain
 - functional deficit
 - limping
 - intraarticular effusion
 - instability



Knee Osteoarthritis: diagnostic

Clinical exam:

- - morphotype (genu varum +++ : 90%)
- +/- déformation globale du genou
- Pain on palpation of the joint line
- Intraarticular effusion
- Popliteal cyst (Baker cyst)
- ROM limitation

Knee Osteoarthritis: diagnostic

RADIOLOGIE:

- x ray knee weight bearing a.p./lateral
- Schuss
- Skyline patella




Standing AP View



PA Flexion View



Lateral View



Sunrise view

RADIOLOGIE

4 stages (Ahlbäck):



Stage 1 :

narrowing < 50%

Stage 2 :

narrowing > 50%



Stage 3 :

Attrition < 5mm

Stage 4 :

attrition > 10mm

canton de
vaud



Treatment


The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE


Arthroscopic Partial Meniscectomy versus Sham Surgery for a Degenerative Meniscal Tear

Raine Sihvonen, M.D., Mika Paavola, M.D., Ph.D., Antti Malmivaara, M.D., Ph.D.,
Ari Itälä, M.D., Ph.D., Antti Joukainen, M.D., Ph.D., Heikki Nurmi, M.D.,
Juha Kalske, M.D., and Teppo L.N. Järvinen, M.D., Ph.D.,
for the Finnish Degenerative Meniscal Lesion Study (FIDELITY) Group

- Treatment of early medial varus OA
 - Valgisation osteotomy
- High tibial open wedge osteotomy




- Treatment of early medial varus OA
 - Valgisation osteotomy
 - High tibial closing wedge osteotomy





TREATMENT

- Treatment of early lateral valgus OA
 - Varisation osteotomy
 - High tibial closing wedge osteotomy





TREATMENT

- Treatment of early lateral valgus OA
 - Varisation osteotomy
 - High tibial closing wedge osteotomy




TREATMENT

- Treatment of early lateral valgus OA
 - Varisation osteotomy
 - Distal femoral closing wedge osteotomy




TREATMENT

- Treatment of advanced isolated OA
 - Unicompartmental arthroplasty (medial, lateral or patellofemoral)




TREATMENT

- Treatment of advanced OA
 - Total knee arthroplasty (tricompartimental resurfacing)





TREATMENT

- Surgical technique



- Mechanical alignment





Tibial resection

- Tibial cut





Tibial resection

- Tibial cut






Femoral resection

- Distal femoral cut






Ligament balancing

- Balance in flexion and extension




Ligament balancing


- Balance in flexion and extension



Navigation



Robotic assisted surgery



Merci