

Reducing agents assisted fed-batch fermentation to enhance ABE yields

Vijaya Chandgude ^a, Teemu Välijalami ^a, Juha Linnekoski ^a, Tom Granström ^b, Bruna Pratto ^a, Tero Eerikäinen ^a, German Jurgens ^c, Sandip Bankar ^a*

^a Department of Bioproducts and Biosystems, School of Chemical Engineering, Aalto University, P.O. Box 16100, FI-00076, Espoo, Finland

^b VTT Technical Research Centre of Finland Ltd., P.O. Box 1000, FI-02044, Espoo, Finland

^c Alimetrics Ltd., Koskelontie 19, 02920, Espoo, Finland



ARTICLE INFO

Keywords:

ABE fermentation
Butanol dehydrogenase
Clostridium acetobutylicum
Fed-batch
NADH
Reducing agents

ABSTRACT

Acetone-butanol-ethanol (ABE) fermentation process is a promising bioenergy option amid rising concerns over the environmental impact of fossil fuel usage. However, the commercialization of the ABE process has been marred by challenges of low product yield and titer, thereby non-competitive process economics. Here, we coupled cost competitive reducing agents with a controlled feeding strategy to improve both product titer and yield. Reducing agents promote cofactor dependent butanol production, while fed-batch operation enhances glucose consumption, final ABE titer, and partly mitigates product toxicity. The effects of ascorbic acid, L-cysteine, and dithiothreitol (DTT) on ABE fed-batch production using *Clostridium acetobutylicum* was investigated in current study. NADH, ATP, extracellular amino acid secretion, and NADH-dependent butanol dehydrogenase (BDH) assays were performed to study the metabolic modifications triggered by reducing agents. Incidentally, L-cysteine and DTT improved ABE solvent titer by 2-fold, producing 24.33 and 22.98 g/L ABE with solvent yields of 0.38 and 0.37 g/g, respectively. Elevated NADH, BDH, and ATP levels in fermentation broth reflected in enhanced ABE titer and yield. Furthermore, histidine secretion emerged as an important factor in *Clostridial* acid stress tolerance in this study. The results demonstrate that addition of reducing agents in fed-batch ABE fermentation operation enables efficient utilization of glucose with significant improvement in solvent production.

1. Introduction

Anthropogenic activities that emit atmospheric greenhouse gases (GHG) have been linked to climate change [1]. The climate change impact warrants reduction in carbon emission, and the scale-up of renewable energy supplies [2]. Among biofuels, biobutanol has attracted considerable attention because of its superior properties (such as high octane number, energy density) which are comparable to gasoline [3]. Moreover, it performs well with Jet A and diesel blends and produces lower emissions of nitrogen oxides (NOx) and carbon monoxide (CO) [4,5].

Traditionally biobutanol has been generated via the process of acetone–butanol–ethanol (ABE) fermentation of starch feedstocks using *Clostridia* [6]. In the ABE pathway (Fig. 1), the carbon flow from acetyl Co-A leads to the formation of acids and solvents via its major branches. ABE fermentation is a biphasic process. In the first phase of acidogenesis,

acetic and butyric acids are produced from sugars. These acids are then secreted into the medium. Acids lower the external pH and serve as inducers for the biosynthesis of solvent generating enzymes. In the second phase, acids assimilate back into *Clostridia* and act as co-substrates to generate neutral solvents (ABE) accompanied by further sugar consumption [7]. However, substantial practical problems of low product titer and yield, unwanted by-products, and butanol toxicity make the ABE process challenging for industrial-scale production of biobutanol [8]. To render the ABE process commercially viable, several process and genetic engineering approaches are being studied, including strain improvement [9], use of lignocellulosic substrates [10,11], and overexpression of targeted functional genes [12].

Production of butanol involves a sequence of reduction steps which utilize intracellular reducing equivalent, NADH (Fig. 1). Electron supply is essential to regenerate consumed reducing equivalents and thereby balance redox metabolism [14]. Redox balance can be manipulated to

* Corresponding author.

E-mail addresses: vijaya.chandgude@aalto.fi (V. Chandgude), teemu.valisalmi@aalto.fi (T. Välijalami), juha.linnekoski@aalto.fi (J. Linnekoski), tom.granstrom@vtt.fi (T. Granström), brunapratto@hotmail.com (B. Pratto), tero.eerikainen@aalto.fi (T. Eerikäinen), gera.jurgens@gmail.com (G. Jurgens), sandip.bankar@aalto.fi (S. Bankar).

improve butanol production via electron receptor addition [15], and bioelectrolysis (BEC) [16]. As the majority of *Clostridia* lack electroactivity, bioelectrical production of butanol needs a redox mediator [14]. Another option is the use of low-cost reducing agents to enhance ABE fermentation.

Addition of reducing agents improve end-product formation in both aerobic and anaerobic fermentation processes. In heterotrophic microbes, most enzymes preferentially utilize NADH as their cofactor [17,18]. These agents affect the parameters of redox/oxidation or redox potential (ORP), ATP, and cofactors, such as NADH. In turn, they influence the expression of certain genes and activities of specific enzymes. They donate electrons to regenerate NADH from NAD⁺ and facilitate the metabolic flux towards the desired metabolite [19]. The addition of neutral red in ABE fermentation resulted in enhanced butanol production with decreased acetone production [20]. Similarly, viologen dyes altered the electron flow towards NADH generation, thereby promoted alcohol formation [21]. However, considering the costs of viologens and neutral red, alternate less expensive reducing agents are essential to make this approach more feasible.

The threshold of substrate level is necessary for sustained solventogenesis [22]. Low substrate levels after the exponential phase (typically after 96 h) in *Clostridial* batch fermentation lead to cessation of ABE production despite the presence of residual substrate, resulting in low yields [23]. Hence, fed-batch operation, which maximizes substrate utilization [24] was considered in the current study. We hypothesized that channeling reducing equivalent towards butanol generation with the aid of reducing agents and maintaining a threshold of substrate level necessary for solvent generation throughout the fermentation would increase the solvent titer, yield, and consequently the efficacy of substrate utilization. Besides, intermittent feeding partially mitigates ABE solvent toxicity [25]. Therefore, the current work involves a comprehensive study of the role of reducing agents in conjunction with the fed-batch operation to enhance ABE production. It provides extensive insights into metabolism dynamics of energy and reducing equivalents, extracellular amino acid secretion under stress, and butanol dehydrogenase in presence of L-cysteine, DTT, and ascorbic acid in fed-batch operation mode. The study sheds light on the possible mechanisms of action of these reducing agents in ABE pathway. The findings

demonstrate an efficient strategy to enhance ABE production in a simple set-up using reducing agents. Furthermore, findings of stress-responsive amino acid secretion can be useful in the development of acid-tolerant *Clostridial* strain for industrial applications.

This study aimed to establish an efficient ABE production system by employing inexpensive reducing agents with *Clostridium acetobutylicum* NRRL B-527 strain. We investigated the effect of L-cysteine, DTT, and ascorbic acid on ABE solvent production in the fed-batch regimen. To study metabolic modifications induced by reducing agents, we analyzed NADH, ATP, extracellular amino acid secretion, and NADH dependent BDH assays.

2. Material and methods

2.1. Bacterial culture and medium

Clostridium acetobutylicum strain NRRL B-527 culture was a gift for research work from ARS (Agricultural Research Services) Culture Collection, USA. It was germinated by inoculating in RCM (reinforced Clostridium medium) containing (g/L): glucose (30), meat extract (10), peptone (5.0), sodium chloride (5.0), yeast extract (3.0), sodium acetate (3.0), starch (1.0), L-cysteine (0.50), and 5 M potassium hydroxide was used to adjust pH to 6.8 ± 0.2. Spore suspension for future use was prepared by inoculating germinated *Clostridial* cells (10% v/v) into starch solution (6% w/v) and stored at room temperature. Spores were activated by adding 2.5% v/v spore suspension into RCM and heat-shocked at 80 °C for 10 min [26]. The pre-culture was grown for 20–24 h until OD 600 reached 1.5. Then ABE fermentation batches were inoculated with 5% v/v pre-culture in production (P2) medium with the following composition (g/L): glucose (60), ammonium acetate (2.2), potassium dihydrogen phosphate (0.5), di-potassium hydrogen phosphate (0.5), magnesium sulfate (0.2), p-aminobenzoic acid (0.1), thiamin (0.1), sodium chloride (0.01), manganese sulfate (0.01), ferrous sulfate (0.01), and biotin (0.01). The medium pH was adjusted to 6.5 with 5 M potassium hydroxide and bubbled with nitrogen for 10 min before autoclaving at 120 °C for 20 min. Analytical grade chemicals were used in this study and were procured from Sigma Aldrich, Finland.




Fig. 1. The ABE production pathway in *Clostridium acetobutylicum* showing the formation of products and the role of reducing power NADH (AK: acetate kinase PAT: phosphate acetyltransferase, AAD: aldehyde alcohol dehydrogenase, AD: acetoacetate decarboxylase, BACoAT: butyrate-acetoacetate CoA-transferase, BADH: butyraldehyde dehydrogenase, BDH: butanol dehydrogenase, BK: butyryl kinase, PBT: phosphate butyryl transferase, Pi: inorganic phosphate) [7,13].

2.2. Batch study of the effect of reducing agents on ABE fermentation

Preliminary experiments were conducted to determine which reducing agents impact ABE fermentation the most and to find their optimum time of addition in the fermentation medium. It was performed in 125 mL airtight serum bottles having a 95 mL P2 medium. Serum bottles inoculated with a 5% v/v actively growing pre-culture of *Clostridium acetobutylicum* NRRL B-527. Seven reducing agents which are relatively economical than conventionally used viologens: glutathione (GLUT), sodium sulfide (SS), methylene blue (MB), phenosafranine (PS), ascorbic acid (AA), L-cysteine (CYS), and dithiothreitol (DTT) were selected. These reducing agents were dissolved in oxygen-free deionized water and were filter-sterilized (0.2-μm membrane filter). Reducing agents were added at the beginning of fermentation (0 h). The optimum concentration of reducing agents was identified by testing 0.1, 0.3, 0.5, and 0.7 mM concentrations. Furthermore, reducing agents were supplemented at 9, 12.5, and 17 h of fermentation to determine their optimal performance. These time intervals represent the initial, the mid, and the end of the exponential phase of fermentation. Batches were cultivated at 37 ± 2 °C and terminated after 96 h. Each experiment was carried at least in triplicate and the presented results report average \pm standard deviation.

2.3. Reducing agent-assisted fed-batch experiments

Fed-batches were conducted in 3 L bioreactors (BIOSTAT B plus, Sartorius, Germany). Experiments were started with a 1000 mL P2 medium. A pH-breakpoint is linked to the shift from acidogenesis to solventogenesis. Hence, pH control was not used [27]. The three best reducing agents from the preliminary experiments were selected and added at an optimized time of 12.5 h from the onset of fermentation. Glucose feeding was initiated when the residual glucose in the bioreactor decreased beneath 40 g/L. The required volume of each feed pulse was determined based on the amount of glucose necessary to replace the utilized glucose in the bioreactor. A total of 5 feedings were performed at 30 (36 for ascorbic acid), 48, 72, 96, and 120 h of fermentation. The fermentation was terminated after 168 h. In the end, the medium volume was 1144.29, 1454.83, and 1539.31 mL for ascorbic acid, DTT, and L-cysteine batch, respectively. The feed medium had a similar composition as the P2 medium except for the glucose concentration (120 g/L). Concentrated substrate feed helps to reduce the required fermentation volume and minimizes the waste disposal streams [27]. The control batch experiment was conducted in a bioreactor to compare its results with the fed-batches assisted by the reducing agents.

2.4. Analytical techniques

Solvents (acetone, butanol, and ethanol), acids (acetic, butyric, lactic), glucose, and acetoin were measured by high-performance liquid chromatography (HPLC) on Waters Alliance 2695 equipped with Rezex H (7.8 × 300 mm) column and Security Guard H⁺ pre-column. The refractive index detector was set at 30 °C, and 14 mM sulphuric acid was used as an eluent in isocratic mode with a 0.6 mL/min flow rate. *Clostridial* growth was estimated by measuring optical density at 600 nm (Eppendorf BioPhotometer plus, Germany).

ATP (Roche 11 699 709 001, Germany) and NADH (Sigma-Aldrich MAK037, USA) were determined by following the methods supplied with the kits. The ATP was assayed by measuring luminescence produced by the reaction between ATP, added D-luciferin, and luciferase. Luminescence was measured with BioTek Cytation 3 multimode reader (BioTek Instruments, USA). NADH was estimated after adding the NAD cycling enzyme mix by measuring absorbance at 450 nm.

NADH-BDH activity was analyzed by applying the method reported by Rajagopalan et al. [28]. The activity determined in the forward direction at 25 °C with 0.5 mM NADH in MES buffer (50 mM, pH 6) and 50 mM butyraldehyde. The reaction was started by adding 10% crude

enzyme extract and absorbance was determined at 340 nm by Eon microplate reader (BioTek Instruments, USA). One unit of activity (U) is equivalent to the amount of BDH needed for oxidation of 1 μmol of NADH/min.

Protein concentration was estimated by applying the Bradford assay and bovine serum albumin as a standard [29].

Amino acids in ABE fermentation broth were quantified by S433 automatic amino acid analyzer (Sykam Chromatography, Germany) with the following conditions: amino acids separated with cation separation column LCA K06/Na 4.6 × 150 mm; column temperature 80 °C; flow rate 0.45 mL/min; eluent was a mixture of buffer A (sodium citrate, pH 3.45) and buffer B (sodium citrate/borate, pH 10.85); colorimetric detection at 570 nm and 440 nm (proline) after ninhydrin reaction.

3. Results and discussion

3.1. Screening of reducing agents in the batch study

The reducing agents were supplemented at 0th h (beginning of the batch) of ABE fermentation. Their effect was studied on butanol and total solvent production. Fig. 2 summarizes the effect of reducing agents at four different concentrations in the batch operation. 0.3 mM DTT emerged as the best reducing agent producing 9.19 ± 0.23 g/L butanol against the control with 6.27 ± 0.02 g/L. 0.3 mM L-cysteine and 0.3 mM ascorbic acid generated 8.33 ± 0.23 and 8.19 ± 0.19 g/L butanol, respectively. Total solvent production followed a similar trend, a corresponding increase of 62.53, 59.56, and 44% when supplemented with 0.3 mM of DTT, L-cysteine, and ascorbic acid, respectively. Raising DTT concentration from 0.3 mM to 0.7 mM resulted in a steep reduction in butanol and ABE titer. DTT in high concentration (3 mM) has proven to be toxic to bacterial cells [30]. Furthermore, higher DTT concentration impacts disulfide bond formation and intracellular protein folding in *E. coli* [31,32]. The addition of 0.1–0.5 mM phenosafranine, sodium sulfide, glutathione, and methylene blue also showed an increase in ABE titer compared to the control.

The ABE pathway (Fig. 1) consists of a sequence of reduction reactions where NADH acts as a crucial coenzyme. Hence, the improvement in ABE titer evident in the preliminary study has probably stemmed from the elevation of both, the regeneration of NADH and the activity of key enzymes such as butanol dehydrogenase (BDH).

Although the present study was conducted in the anaerobic milieu, an improved oxygen-scavenging or aero-tolerance attributed by reducing agents (Ascorbic acid, DTT, and L-cysteine) could prove very beneficial since it is difficult to maintain strict anaerobic conditions on the industrial scale [33]. L-cysteine is reported to be beneficial for *Clostridial* growth in ABE fermentation under stress [34]. Other reducing agents such as glutathione also improved *Clostridial* aero-tolerance and enhanced butanol production [33].

We selected ascorbic acid, DTT, and L-cysteine at 0.3 mM concentration to further explore their effect on ABE production.

3.2. Adding reducing agents at different time intervals in batch mode

The physiological state of microbial cultures varies at different fermentation times. Therefore, we assumed that the supplementation of reducing agents at different fermentation times would affect butanol and total solvent titers to different extents. Ascorbic acid, DTT, and L-cysteine were added at 9, 12.5, and 17 h of fermentation, which represents the initial, the mid, and the end of the *Clostridial* exponential (high metabolic activity) phase.

All the reducing agents showed the best results when added at 12.5 h (Fig. 3). DTT produced the highest butanol and solvent titer of 10.48 ± 0.22 and 17.23 ± 0.23 g/L against 7.00 ± 0.26 and 10.56 ± 0.41 g/L of butanol and total solvents in the control (without reducing agent) experiment. L-cysteine and ascorbic acid generated 9.85 ± 0.11 and 9.64 ± 0.08 g/L butanol along with 15.53 ± 0.14 and 13.11 ± 0.01 g/L




Fig. 2. Effect of reducing agents on total solvent and butanol production by using *C. acetobutylicum* NRRL B-527 batch study after 96 h.




Fig. 3. Effect of time of addition of reducing agents on total solvent and butanol production using *C. acetobutylicum* NRRL B-527 batch study after 96 h.

of total solvents respectively. This translates into a 14–18% boost in the butanol titer compared to the addition of reducing agents at 0th h (Figs. 2 and 3).

In the mid-exponential phase (12.5 h), the NADH conversion circulation is high [35]. Thus, higher NADH generation possibly improved butanol production. The addition of reducing agents at the end of the exponential phase (17 h) declined solvent generation due to decreasing metabolic activity of pyruvate oxidation to acetyl-CoA. Reducing agent supplementation moderately improved the butanol and the total solvent titer in the batch study. Their positive effect was studied further in fed-batch operation.

3.3. Reducing agent-assisted fed-batch operation to enhance ABE production

3.3.1. Effect of reducing agents on ABE production

The threshold of glucose level necessary for solventogenesis can limit

the efficiency of ABE production. ABE generation is very low when the medium glucose concentration is below 20 g/L [22]. The current study followed a similar trend. Butanol titer in the batch fermentation stagnated in the late stationary phase, especially due to substrate limitation. Table 1 shows the residual glucose and ABE yields in batch fermentation experiments. Therefore, reducing agent addition after 12.5 h was coupled to fed-batch fermentation with controlled feeding to extend the solventogenic phase (thereby improving glucose consumption, ABE titer, and yield).

Preliminary fed-batches were conducted to determine the optimum glucose level necessary to achieve efficient ABE production in the solventogenic phase. Both continuous and pulse feeding strategies were tested to select a suitable substrate feeding mode. Among the tested medium glucose concentrations, 40 g/L glucose and pulse feeding strategy proved suitable for efficient ABE production (data not shown).

In ABE fed-batch operation, L-cysteine (0.3 mM) and DTT (0.3 mM) significantly improved solvent production. L-cysteine and DTT boosted the total ABE solvents up 98% (24.33 ± 0.30 g/L) and 87% (22.98 ± 0.09 g/L) compared to the control batch production (12.28 ± 0.1 g/L) (Fig. 4a). Ascorbic acid (0.3 mM) supplemented fed-batch showed a modest increase of 38% by producing 16.95 ± 0.21 g/L total solvents.

From 24 h onward, the DTT batch displayed relatively lower butanol and solvent production than the L-cysteine batch due to slower reassimilation of acids especially acetic acid (Figs. 4 and 5). Lower quantities of acids in the culture medium reflect their higher reassimilation in *Clostridia* and their effective conversion in solvents [36]. A study

Table 1

Residual glucose and ABE solvent yield after 96 h of batch fermentation using *C. acetobutylicum* NRRL B-527 in serum bottles.

Sample	Residual glucose (g/L)	% Residual glucose	ABE yield (g/g)
Control	14.64 ± 0.51	24.40	0.233
0.3 mM DTT	7.50 ± 0.73	12.50	0.328
0.3 mM L-Cysteine	9.77 ± 0.11	16.28	0.309
0.3 mM Ascorbic acid	9.93 ± 0.09	16.55	0.262




Fig. 4. Effect of reducing agents on ABE production using *C. acetobutylicum* NRRL B-527 in fed-batch operation (a) total ABE solvents, (b) butanol production, (c) acetone production, (d) ethanol production. F1, F2, F3, F4, and F5 represent feeding points at 30 (36 for ascorbic acid), 48, 72, 96, and 120 h of fermentation.




Fig. 5. Effect of reducing agents on (a) acetic acid and (b) butyric acid production in the fed-batch study using *C. acetobutylicum* NRRL B-527.

conducted with C^{14} labeled acids showed that a major quantity of reassimilated butyrate and more than half of the acetate could be both reduced to butanol [7]. Likely, the slower reassimilation of acids into *Clostridia* can be due to the modulation of the membrane proteins and the transport systems by DTT during acid reuptake. In fact, DTT is a membrane permeating reducing agent. It affects the thiol group of proteins in the bacterial membrane, which in turn influence the affinity and the activity of both transport systems and membrane-associated enzymes [37,38]. Despite the slower reassimilation at the beginning, DTT showed a significant gain in solvent production during late solventogenesis.

The addition of L-cysteine demonstrated a significant effect on solvent production and ABE total solvent yield in fed-batch operation. The L-cysteine batch manifested higher acid reuptake than DTT and ascorbic acid batches. Butyric acid reassimilation was faster in the L-cysteine batch than other batches including the control batch. Fed-batch operation re-assimilated even lower quantities of organic acids in bacteria from the fermentation medium (Fig. 5). This is also evident in Fig. 6, where L-cysteine showed the highest solvents to acids ratio of 27.97 followed by DTT with 25.25, and ascorbic acid with 14.49 against 8.90




Fig. 6. Solvents to acids ratio in reducing agent-assisted fed-batches.

in the control.

Ascorbic acid showed only a moderate impact on ABE production compared to L-cysteine and DTT in fed-batch experiments. It is reported that ascorbic acid forms a chelate complex with transition metal ions [39]. Hence, ascorbic acid might have formed a chelate complex with transition metal ions, iron, and manganese present in the production medium and feeding medium resulted in delayed *Clostridial* growth (Supplementary figure). The acid production profile reflected this delay where it reached the maximum level at 21st h contrary to 14th h in presence of other reducing agents (Fig. 5). Consequently, the ascorbic acid batch indicated slower acid re-assimilation and thereby lower ABE production (Fig. 4).

In fed-batch operation, the butanol level passed from 8.17 ± 0.12 g/L in the control to a maximum of 16.26 ± 0.51 g/L in the L-cysteine batch. DTT and ascorbic acid batches measured 15.35 ± 0.03 and 10.58 ± 0.09 g/L butanol, respectively. Addition of DTT and L-cysteine decrease the redox potential of fermentation medium [38,40,41]. Negative redox potential is linked to the amplification of reduction processes which is a peculiar feature of anaerobic fermentation [42], and it enhances intracellular NADH [42–44]. Reducing agents reduce NAD^+ back to NADH by donating electrons. This occurs if their redox potential is below -316 mV (NAD^+ redox potential) [45] and thus assists in NADH regeneration. DTT and L-cysteine have redox potential of -332 mV [46] and -340 mV [47] respectively. Therefore, DTT and L-cysteine increased NADH regeneration and consequently improved butanol production. Furthermore, elevated NADH levels might have contributed in enhancing activities of crucial enzymes of the butanol production pathway, such as butanol dehydrogenase (BDH) and butyraldehyde dehydrogenase (BADH) which are NADH dependent [48,49].

Wild-type *C. acetobutylicum* typically produces maximal butanol titer of only 13 g/L [7,50]. Butanol accumulation deranges physiological functions of bacterial cell membrane including membrane ATPase, pH gradient, membrane potential, and transportation [51–53]. The disruption of the membrane functions may impede the proteins involved in oxidation–reduction reactions, promoting the accumulation of reactive oxygen species (ROS) [54]. *C. acetobutylicum* is armed with several pathways for detoxification of ROS including glutathione peroxidase-like proteins [55] and superoxide reductase (SOR) such as desulfo-ferrodoxin [56].

The enhanced fermentation performance achieved in the present study shows improved survival of *Clostridia* to solvent, acid, and osmotic stress reflected in conventional ABE solvent production (Fig. 4), glucose consumption (Fig. 7), and growth profile (Supplementary figure). Experimental data validate that the fermentation did not falter even in the presence of 24 g/L total solvents and 16 g/L butanol. These results are in alignment with Burhan et al. [57] study where cysteine derivative N-acetyl-L-cysteine (NAC) mitigated the oxidative stress in yeast




Fig. 7. Effect of reducing agents on *C. acetobutylicum* NRRL B-527 glucose consumption in fed-batch mode.

and thereby enhanced the fermentation performance. NAC also enhanced the activity of cytosolic Cu-Zn supraoxide dismutase (Cu-Zn SOD), which is important in protection from ROS in the yeast [57]. Desulfoferrodoxin enzyme in *Clostridium acetobutylicum* functions as a superoxide reductase in ROS detoxification, and its active center consists of $[\text{Fe}(\text{NH}_3)_4(\text{SCys})]$ cysteine residue [56]. Considering this, the present study raises the possibility that L-cysteine may have exerted a positive impact on reductases and peroxidases such as desulfoferridoxin involved in ROS scavenging in ABE fermentation.

Interestingly DTT is also reported to decrease the level of ROS by 65.7% in *Pseudomonas chlororaphis* GP72AN and enhanced generation of biocontrol antibiotic phenazine-1-carboxylic acid (PCA) by 1.8- fold [58]. Therefore, DTT as well might have protected the *Clostridia* against ROS in ABE fermentation.

The protective effect of ascorbic acid is well documented in aerobic fermentation. However, the literature on its impact on anaerobic fermentation is scarce. Ascorbic acid exerted a positive effect on bacterial and yeast cells through modulating medium redox potential, prolonging the mean and maximal life span in Cu-Zn superoxide dismutase mutant yeast [59,60]. Nonetheless, in the present study ascorbic acid supplemented experiments could not cross the lethal butanol level. Hence, the effect of ascorbic acid in mitigating solvent stress is not clear from this study.

Other reducing agents, neutral red and methyl viologen have been frequently used in ABE fermentation to divert metabolic flux towards butanol production [20,21]. Interestingly, the mechanism to improve butanol titer seems dissimilar in the current study, especially because of the prominent acetone production (Fig. 4 and Table 2) unlike in production processes supplemented with methyl viologen or neutral red. In fact, all reducing agents improved acetone yield compared to the control (Table 2). The improvement in acetone yield can be attributed to improved *Clostridial* survival and intermittent glucose feeding. Reducing agents also exerted a beneficial effect on ethanol production and yield (Fig. 4 and Table 2). Since both ethanol and butanol production pathways require NADH (Fig. 1), their order of effect on production and yield is also similar (L-cysteine > DTT > Ascorbic acid). Butanol yield increased noticeably than ethanol in the present study (Fig. 4 and Table 2). The huge difference in ethanol and butanol yield is explained by the fact that, when *Clostridia* in ABE fermentation transit from the acidogenic phase to the solventogenic phase amidst the adverse environment of low pH, butanol is the primary product, while ethanol is only

Table 2
Effect of reducing agents on ABE fermentation after 168 h.

Parameter	Control	L-cysteine	DTT	Ascorbic acid
Butanol (g/L)	8.17 ± 0.12	16.26 ± 0.51	15.35 ± 0.03	10.58 ± 0.09
Acetone (g/L)	3.21 ± 0.10	6.38 ± 0.10	6.25 ± 0.09	5.30 ± 0.20
Ethanol (g/L)	0.90 ± 0.01	1.69 ± 0.04	1.38 ± 0.01	1.07 ± 0.03
Acetic acid (g/L)	0.95 ± 0.03	0.60 ± 0.03	0.66 ± 0.01	0.76 ± 0.02
Butyric acid (g/L)	0.43 ± 0.04	0.27 ± 0.01	0.25 ± 0.03	0.41 ± 0.06
Total ABE (g/L)	12.28 ± 0.10	24.33 ± 0.30	22.98 ± 0.09	16.95 ± 0.21
Total acids (g/L)	1.38 ± 0.04	0.87 ± 0.03	0.91 ± 0.03	1.17 ± 0.06
ABE productivity (g/L.h) ^a	0.11 ± 0.02	0.26 ± 0.05	0.19 ± 0.02	0.16 ± 0.01
Butanol yield (g/g)	0.166	0.250	0.245	0.199
Acetone yield (g/g)	0.065	0.100	0.099	0.099
Ethanol yield (g/g)	0.018	0.026	0.022	0.020
ABE yield (g/g)	0.250	0.382	0.367	0.318
Solvents:acids ratio (g/g)	8.90	27.97	25.25	14.49

^a Value measured at 48 h of fermentation.

the marginal one.

The butanol level and the solvent yield benefitted from the application of both controlled glucose feeding and reducing agents. Although experiments with reducing agents consumed more glucose (Fig. 7), they produced less biomass as evident from the OD profile (Supplementary figure). Thus, reducing agents apparently diverted glucose more towards solvent production. Therefore, at the end of the fermentation (168 h), the L-cysteine batch exhibited a 51% increase in butanol yield followed by DTT with 48%, and ascorbic acid with a 20% increase compared to the control. The total ABE solvent yield followed a similar trend since butanol is the major product (Fig. 8 and Table 2). From the very beginning, the L-cysteine batch exhibited better acid reassimilation and its effective conversion in solvents, thereby displayed the highest solvents to acids ratio (Fig. 6) throughout the fermentation period. Hence, L-cysteine was found to be the most active in terms of total ABE solvents yield. This system also enhanced solvent productivity compared to the control. The maximum productivity of 0.26 g/L h in L-cysteine, 0.19 g/L h in DTT, and 0.16 g/L h in ascorbic acid versus 0.11 g/L h in the control batch were observed at 48 h of fermentation.

Table 3 shows a comparison of ABE fermentation performance with commonly used reducing agents and the current study. The present study (with L-cysteine and DTT) is competitive with the other studies in terms of butanol and ABE titer. However, ABE productivity still needs to be improved for economic large-scale production. The productivity can be improved by implementing advanced bioprocess technologies such as two-stage packed bed column reactor operation with simultaneous product recovery module system [26]. Likewise, the employment of engineered *Clostridia* [61,73] can further enhance both butanol yields and productivities. Hence, initial findings in this study indicate the applicability of reducing agents in fed-batch combination to further improve ABE fermentation production. The use of ascorbic acid is not as promising as L-cysteine and DTT.

3.3.2. Effect of reducing agents on NADH

After verifying the effect of reducing agents on the butanol production, the effect of the same agents was evaluated on the NADH levels in *Clostridia* during the solventogenic phase. NADH accumulates during acidogenesis and then decreases to the basic level during the solvent production phase [64]. As visible in Table 4, L-cysteine, DTT, and ascorbic acid increased NADH concentration by 36%, 29%, and 7% compared to the control.

Enhanced solvent production and yield reflected the beneficial effect of elevated NADH. Nevertheless, it is conceivable that increased NADH




Fig. 8. Effect of reducing agents on butanol yield and solvent yield in *C. acetobutylicum* NRRL B-527 fed-batch study.

levels achieved after the addition of reducing agents is not the sole factor responsible for higher butanol production. Presumably, the reducing agents (especially L-cysteine and DTT) must have exerted their favorable effect on the ABE pathway through the multiple mechanisms discussed in Section 3.3.1, culminating in the high butanol yield and titer.

3.3.3. Effect of reducing agents on NADH-dependent butanol dehydrogenase activity

Clostridia has multiple aldehyde/alcohol dehydrogenases. NADH-dependent butanol dehydrogenase (NADH-BDH) is the major enzyme accountable for butanol synthesis in ABE fermentation as it exhibits the highest activity during solventogenesis [28].

Since a reduced environment is conducive for catalytical activity and stability of alcohol dehydrogenases including BDH [28], the reducing agent supplemented fed-batches displayed higher NADH-BDH activities (Table 5). L-cysteine batch showed the highest NADH-BDH activity (3-fold higher than control), followed by DTT (2-fold higher than control) and ascorbic acid (a slight increase over control). The butanol levels achieved in the present study highlight the quintessential role of NADH-BDH in ABE fermentation. Besides, enhanced NADH-BDH activities led to elevated solvents to acids ratio of 27.97 and 25.25 in L-cysteine and DTT batches, relative to the ratio of 8.9 in the control (Fig. 6).

3.3.4. Effect of reducing agents on ATP

The promotion of solvent formation reduces ATP demand [64]. Here, solvent production improved in the presence of reducing agents. Hence ATP analysis was performed to check their impact on ATP levels during fermentation (Fig. 9). Before the addition of reducing agents (12 h), ATP levels were high due to the exponential phase conversion of glucose into acetic and butyric acid (acid formation generates ATP) [65]. After 24 h, ATP concentration decreased in all the batches, especially in the ascorbic acid batch. This decrease can be explained by higher ATP demand (utilization) to enhance membrane-active (ATPase) permeability to maintain physiological intracellular pH in the presence of high acid concentrations [66]. Interestingly, the steep ATP decrease in the ascorbic acid batch coincides with the delayed acid peak and the prolonged acidogenic phase (Figs. 5 and 9). ATP increased till 48 h accompanied by acid reuptake and biomass increase (Supplementary figure). A decrease in biomass from 48 to 120 h led to a decrease in the ATP level in *Clostridia*. In the late solventogenic phase (72–120 h), the ATP levels in the DTT and L-cysteine batches were still high compared to the control.

As discussed in Section 3.3.1, DTT and L-cysteine might have protected *Clostridia*, from fermentation stress while the same cannot be stated for ascorbic acid. ATP is crucial in cell maintenance and repair. Hence high ATP levels detected in DTT and L-cysteine batches can be attributed to lower ATP demand due to their protective actions. On the other hand, the protective effect was not evident in the ascorbic acid batch. Thus, it might have led to higher ATP consumption for cell repair and maintenance and therefore lower levels of ATP detected in the ascorbic acid batch. The high ATP levels obtained in DTT and L-cysteine batches reflected in increased butanol and acetone production as decreased ATP demand promotes solventogenesis [64].

3.3.5. Effect of reducing agents on protective amino acids secretion in ABE culture

During ABE fermentation, *Clostridia* are subjected to multiple stresses, including acid and solvent stress. Cells respond, adapt, and develop tolerance to multiple stressors during cultivation. However, the mechanisms involved in encountering the overlapped stresses are only partially investigated and remain poorly understood [67]. Metabolomic analysis of *Clostridia* has revealed that increased amino acid levels, such as isoleucine and valine, confer butanol tolerance to bacteria [68]. Several studies found that to better survive under the stress caused by ABE fermentation, *Clostridia* secrete amino acids such as phenylalanine, methionine, leucine, valine, and tyrosine in culture medium [15,69].

Table 3

Comparison of performance indicators of ABE fermentation with different reducing agents.

Reducing agent	Strain	Operation mode, carbon source	Butanol (g/L)	Improvement to control (%) (Butanol titer)	Butanol yield (g/g)	ABE (g/L)	ABE yield (g/g)	ABE productivity (g/L h)	Reference
0.1 mM Neutral red	<i>Clostridium beijerinckii</i> IB4	Batch in micro-electrolysis cell (MEC), glucose	10.5	12	0.250	14.68	0.349	0.52	[20]
5.1 mM Sodium sulfide	<i>Clostridium acetobutylicum</i> CICC8012	Batch, wheat straw hydrolysate	7.63	48	0.192	13.85	0.350	0.19	[62]
1.0 mM Methyl viologen	<i>Clostridium acetobutylicum</i> ATCC 824	Batch, glucose	11.93	23	–	13.98	–	–	[63]
0.5 mM Methyl viologen	<i>Clostridium tyrobutyricum</i> <i>CtAck-adhE2</i>	Batch, glucose	13.00	41	0.340	–	–	–	[61]
0.025 mM Benzyl viologen	<i>Clostridium tyrobutyricum</i> <i>CtAck-adhE2</i>	Batch, glucose	13.70	49	0.310	–	–	–	[61]
0.5 mM Methyl viologen and 0.025 mM Benzyl viologen	<i>Clostridium tyrobutyricum</i> <i>CtAck-adhE2</i>	Repeated batch with immobilized cells in a fi-brous-bed bioreactor (FBB), first batch with glucose, subsequent batches with cassava bagasse hydrolysate	15.00	63	0.300	–	–	–	[61]
0.3 mM L-cysteine	<i>Clostridium acetobutylicum</i> NRRL B-527	Fed-batch, glucose	16.26	99	0.250	24.33	0.382	0.26	This study
0.3 mM DTT	<i>Clostridium acetobutylicum</i> NRRL B-527	Fed-batch, glucose	15.35	88	0.245	22.98	0.367	0.19	This study
0.3 mM Ascorbic acid	<i>Clostridium acetobutylicum</i> NRRL B-527	Fed-batch, glucose	10.58	29	0.199	16.95	0.318	0.16	This study

Table 4Effect of reducing agents on NADH after 48 h *C. acetobutylicum* NRRL B-527 fed-batch operation.

Fermentation sample	NADH (μmole/g dry cell weight)
Control	0.809 (±0.01)
Ascorbic acid	0.862 (±0.03)
DTT	1.040 (±0.01)
L-cysteine	1.101 (±0.02)


Table 5Effect of reducing agents on NADH-BDH activity after 168 h *C. acetobutylicum* NRRL B-527 fed-batch cultivation.

Fermentation sample	NADH-BDH (U/mg)
Control	0.021 (±0.002)
Ascorbic acid	0.025 (±0.001)
DTT	0.047 (±0.002)
L-cysteine	0.070 (±0.003)

Given this, we verified whether elevated butanol levels achieved in the current study increased the secretion of protective amino acids in the *Clostridia*. Thus, after the supplementation of reducing agents, we analyzed the secretion of stress-responsive amino acids in the ABE culture medium.

The majority of amino acids were quantified after 30 h of fermentation. Interestingly, an increase in proline, methionine, isoleucine, alanine, and histidine levels were detected (Fig. 10). DTT, L-cysteine, and ascorbic acid batch demonstrated an approximately 2 to 3-fold increase in amino acid release in the medium. Branched-chain amino acids (BCAAs) including isoleucine help in the adaptation of membrane fluidity in response to butanol stress in *Clostridia* [68,70]. Proline protects the cells from alcohol stress [71].

Although ascorbic acid batch only moderately improved butanol production, histidine level manifested a 64% increase over the control (Figs. 4b and 10). This led us to believe that, besides the butanol stress,

Fig. 9. Effect of reducing agents on ATP level in *C. acetobutylicum* NRRL B-527 fed-batch cultivation.

some other stress might have promoted histidine secretion in the ascorbic acid culture broth. Some protective amino acids are not only involved in countering solvent stress but also carboxylic acid stress in bacteria [67]. Broadbent et al. [72] proved that the induction of histidine biosynthesis genes improved acid tolerance in *Lactobacillus casei*. In the present study, ascorbic acid experiments showed decreased reuptake of acids. It led to their high residual levels in the fermentation medium (Fig. 5). Therefore, higher secretion of histidine in ascorbic acid batch, confirms its direct protective role in acid tolerance in *C. acetobutylicum*. This finding is interesting since Wang et al. [70] found that histidine biosynthesis genes (CAC0935-43) were upregulated under both butyrate and butanol stress in *C. acetobutylicum*. Hence, the current study proposes the protection of *Clostridia* by histidine from butyrate as well as acetate stress. The experimental results also validate the importance of protective amino acids in *Clostridial* stress. These amino acids may influence other aspects of *Clostridial* physiology in addition to those discussed here. Therefore, to gain insights, we recommend further study such as transcriptional analysis of amino acid synthesis in the presence




Fig. 10. Effect of reducing agents on protective amino acid secretion in *C. acetobutylicum* NRRL B-527 culture broth after 30 h.

of these reducing agents.

4. Cost comparison of reducing agents

The current study demonstrates the positive impact of L-cysteine, DTT, and ascorbic acid on solvent yield and titer in ABE fermentation. However, to apply them on industrial-scale, cost-effectiveness is crucial. The reducing agents used in this study are noticeably more affordable than commonly used reducing agents in ABE fermentation, neutral red, and methyl viologen (Figs. 11 and 12). Moreover, the use of smaller amounts (0.3 mM) would further facilitate their application. Although both L-cysteine and DTT significantly improved butanol and ABE titer, in terms of reducing agent cost per unit butanol and ABE production, L-cysteine is a more attractive option compared to DTT. Hence, more apt for large scale application. The cost can be further cut down by employing food wastes rich in these reductants, for instance, ascorbic acid from citrus fruits juice and L-cysteine from chicken broth. We recommend detailed techno-economic analysis to establish the economic feasibility of the current process.

5. Conclusions

An efficient ABE production system was developed by employing cost-competitive reducing agents with controlled glucose feeding regimen in fed-batch operation. ABE titer, yield, and consequently the efficacy of substrate utilization were significantly increased in this study. Substantial increase in NADH, BDH, and ATP levels were observed in presence of L-cysteine and DTT, which reflected in improved ABE production. Reducing agents used in the present study did not impact acetone production. Hence, it is proposed that L-cysteine, ascorbic acid, and DTT have a different mechanism of action than conventional reducing agents such as viologens and neutral red. Furthermore, histidine secretion emerged as an important factor in *Clostridial* acid stress tolerance in this study. This factor will be useful in developing acid-tolerant *Clostridial* strain for industrial applications. ABE productivity of the current system can further be enhanced by integrating it with *in-situ* ABE solvent recovery module. To gain a deeper understanding of stress-responsive amino acid secretion in ABE fermentation, further studies such as transcriptional analysis of amino acid synthesis in the presence of reducing agents is recommended. The results from current study will be useful in developing an advanced cultivation system and separation technology by carefully considering the feedstock applicability in large-scale operation.




Fig. 11. Cost comparison of reducing agents used in the present study versus commonly used reducing agents in ABE fermentation. Costs are based on data from Molekula Group and Acros Organics (Methyl viologen) on 22/04/2020.




Fig. 12. Comparison of reducing agents cost per unit butanol and ABE production. Ascorbic acid, DTT, and L-cysteine: present study, neutral red: [73], methyl viologen: [63].

CRediT authorship contribution statement

Vijaya Chandgude: Conceptualization, Investigation, Methodology, Formal analysis, Writing - original draft, Visualization, Writing - review & editing. **Teemu Välijämsi:** Formal analysis. **Juha Linnekoski:** Formal analysis. **Tom Granström:** Conceptualization. **Bruna Pratto:** Visualization, Writing - review & editing. **Tero Eerikäinen:** Conceptualization. **German Jürgens:** Conceptualization. **Sandip Bankar:** Funding acquisition, Supervision, Writing - review & editing, Conceptualization.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are grateful to the Aalto School of Chemical Engineering and the Finnish Cultural Foundation for the financial support of this study. We acknowledge experimental help from Lauri Honkanen and analysis help from Esa Uosukainen. We thank Riikka Särkälä, Ville Aumala, Sami Havukainen, and He Li for helpful suggestions. We also acknowledge Nahla Osmanbegovic for useful comments on the manuscript. Aalto University Bioeconomy infrastructure is acknowledged.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.enconman.2020.113627>.

References

- Pachauri RK, Reisinger A, editors. Climate change 2007: synthesis report. Contribution of working groups I, II and III to the fourth assessment report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland; 2007. <https://www.ipcc.ch/report/ar4/syr/> (downloaded on May 5, 2020).
- Liao JC, Mi L, Pontrelli S, Luo S. Fuelling the future: Microbial engineering for the production of sustainable biofuels. *Nat Rev Microbiol* 2016;14:288–304. <https://doi.org/10.1038/nrmicro.2016.32>.
- Galloni E, Fontana G, Staccone S, Scala F. Performance analyses of a spark-ignition engine firing with gasoline-butanol blends at partial load operation. *Energy Convers Manag* 2016;110:319–26. <https://doi.org/10.1016/j.enconman.2015.12.038>.
- Mendez CJ, Parthasarathy RN, Gollahalli SR. Performance and emission characteristics of butanol/Jet A blends in a gas turbine engine. *Appl Energy* 2014; 118:135–40. <https://doi.org/10.1016/j.apenergy.2013.12.011>.
- Yun H, Choi K, Lee CS. Effects of biobutanol and biobutanol-diesel blends on combustion and emission characteristics in a passenger car diesel engine with pilot injection strategies. *Energy Convers Manag* 2016;111:79–88. <https://doi.org/10.1016/j.enconman.2015.12.017>.
- Nilsson A, Shabestary K, Brandão M, Hudson EP. Environmental impacts and limitations of third-generation biobutanol life cycle assessment of n-butanol produced by genetically engineered cyanobacteria. *J Ind Ecol* 2020;24:205–16. <https://doi.org/10.1111/jiec.12843>.
- Jones DT, Woods DR. Acetone-butanol fermentation revisited. *Microbiol Rev* 1986; 50:484–524. <https://doi.org/10.1128/mmbr.50.4.484-524.1986>.
- Jang YS, Malaviya A, Cho C, Lee J, Lee SY. Butanol production from renewable biomass by *Clostridia*. *Bioresour Technol* 2012;123:653–63. <https://doi.org/10.1016/j.biortech.2012.07.104>.
- Blaschek H, Bassam A, Formanek J, Chen C, Wales N. Inventors; The Board of Trustees of the University of Illinois, Urbana, IL (US), assignee. Method of producing butanol using a mutant strain of *Clostridium beijerinckii*. US patent 6,358,717B1; 2002 Mar 19.
- Pratto B, Chandgude V, de Sousa R, Cruz AJG, Bankar S. Biobutanol production from sugarcane straw: defining optimal biomass loading for improved ABE fermentation. *Ind Crops Prod* 2020;148:1–9. <https://doi.org/10.1016/j.indcrop.2020.112265>.
- Farmanbordar S, Karimi K, Amiri H. Municipal solid waste as a suitable substrate for butanol production as an advanced biofuel. *Energy Convers Manag* 2018;157: 396–408. <https://doi.org/10.1016/j.enconman.2017.12.020>.
- Yu M, Zhang Y, Tang IC, Yang ST. Metabolic engineering of *Clostridium tyrobutyricum* for n-butanol production. *Metab Eng* 2011;13:373–82. <https://doi.org/10.1016/j.ymben.2011.04.002>.
- Buehler EA, Mesbah A. Kinetic study of acetone-butanol-ethanol fermentation in continuous culture. *PLoS One* 2016;11:1–21. <https://doi.org/10.1371/journal.pone.0158243>.
- Liu X, Yu X. Enhancement of butanol production: From biocatalysis to biocatalysis. *ACS Energy Lett* 2020;5:867–78. <https://doi.org/10.1021/acsenergylett.9b02596>.
- Ding J, Luo H, Xie F, Wang H, Xu M, Shi Z. Electron receptor addition enhances butanol synthesis in ABE fermentation by *Clostridium acetobutylicum*. *Bioresour Technol* 2018;247:1201–5. <https://doi.org/10.1016/j.biortech.2017.09.010>.
- Khosravaniour Mostafazadeh A, Drogui P, Brar SK, Tyagi RD, Le Bihan Y, Buelna G, et al. Enhancement of biobutanol production by electromicrobial glucose conversion in a dual chamber fermentation cell using *C. pasteurianum*. *Energy Convers Manag* 2016;130:165–75. <https://doi.org/10.1016/j.enconman.2016.10.050>.
- Park J, Choi Y. Cofactor engineering in cyanobacteria to overcome imbalance between NADPH and NADH: a mini review. *Front Chem Sci Eng* 2017;11:66–71. <https://doi.org/10.1007/s11705-016-1591-1>.
- Meng H, Liu P, Sun H, Cai Z, Zhou J, Lin J, et al. Engineering a d-lactate dehydrogenase that can super-efficiently utilize NADPH and NADH as cofactors. *Sci Rep* 2016;6:1–8. <https://doi.org/10.1038/srep24887>.
- Liu C-G, Qin J-C, Lin Y-H. Fermentation and redox potential. *Ferment Process* 2017;13.
- He AY, Yin CY, Xu H, Kong XP, Xue JW, Zhu J, et al. Enhanced butanol production in a microbial electrolysis cell by *Clostridium beijerinckii* IB4. *Bioprocess Biosyst Eng* 2016;39:245–54. <https://doi.org/10.1007/s00449-015-1508-2>.
- Du Y, Jiang W, Yu M, Tang I, Yang S. Metabolic Process Engineering of *Clostridium tyrobutyricum* Δack – adhE2 for Enhanced n-butanol Production From Glucose : Effects of Methyl Viologen on NADH Availability, Flux Distribution, and Fermentation Kinetics n.d. <https://doi.org/10.1002/bit.25489>.
- Long S, Jones DT, Woods DR. Initiation of solvent production, clostridial stage and endospore formation in *Clostridium acetobutylicum* P262. *Appl Microbiol Biotechnol* 1984;20:256–61. <https://doi.org/10.1007/BF00250635>.
- Ibrahim MF, Abd-Aziz S, Yusoff MEM, Phang LY, Hassan MA. Simultaneous enzymatic saccharification and ABE fermentation using pretreated oil palm empty fruit bunch as substrate to produce butanol and hydrogen as biofuel. *Renew Energy* 2015;77:447–55. <https://doi.org/10.1016/j.renene.2014.12.047>.
- Dolejš I, Krasňan V, Stloukal R, Rosenberg M, Rebroš M. Butanol production by immobilised *Clostridium acetobutylicum* in repeated batch, fed-batch, and continuous modes of fermentation. *Bioresour Technol* 2014;169:723–30. <https://doi.org/10.1016/j.biortech.2014.07.039>.
- Maiti S, Gallastegui G, Sarma SJ, Brar SK, Le Bihan Y, Drogui P, et al. A re-look at the biochemical strategies to enhance butanol production. *Biomass Bioenergy* 2016;94:187–200. <https://doi.org/10.1016/j.biombioe.2016.09.001>.
- Bankar SB, Survase SA, Singh RS, Granström T. Continuous two stage acetone-butanol-ethanol fermentation with integrated solvent removal using *Clostridium acetobutylicum* B 5313. *Bioresour Technol* 2012;106:110–6. <https://doi.org/10.1016/j.biortech.2011.12.005>.
- Ezeji T, Qureshi N, Blaschek H. Acetone butanol ethanol (ABE) production from concentrated substrate: reduction in substrate inhibition by fed-batch technique and product inhibition by gas stripping. *Appl Microbiol Biotechnol* 2004;63:653–8. <https://doi.org/10.1007/s00253-003-1400-x>.
- Rajagopalan G, He J, Yang KL. A highly efficient NADH-dependent butanol dehydrogenase from high-butanol-producing *Clostridium* sp. BOH3. *Bioenergy Res* 2013;6:240–51. <https://doi.org/10.1007/s12155-012-9253-8>.
- Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–54. <https://doi.org/10.2135/cropsci1994.0011183X003400020031x>.
- Wang XD, Mauvais G, Cachon R, Diviès C, Feron G. Addition of reducing agent dithiothreitol improves 4-decanolide synthesis by the genus *Sporidiobolus*. *J Biosci Bioeng* 2000;90:338–40. [https://doi.org/10.1016/S1389-1723\(00\)80092-4](https://doi.org/10.1016/S1389-1723(00)80092-4).
- Missakas D, Georgopoulos C, Raina S. Identification and characterization of the *Escherichia coli* gene dsbB, whose product is involved in the formation of disulfide bonds in vivo. *Proc Natl Acad Sci USA* 1993;90:7084–8. <https://doi.org/10.1073/pnas.90.15.7084>.
- Gill R, Cha H, Jain A, Rao G, Bentley W. Generating controlled reducing environments in aerobic recombinant *Escherichia coli* fermentations: effects on cell growth, oxygen uptake, heat shock protein expression, and in vivo CAT activity. *Biotechnol Bioeng* 1998;59:248–59. [https://doi.org/10.1002/\(SICI\)1097-0290\(19980720\)59:2<248::AID-BIT12>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1097-0290(19980720)59:2<248::AID-BIT12>3.0.CO;2-A).
- Zhu L, Dong H, Zhang Y, Li Y. Engineering the robustness of *Clostridium acetobutylicum* by introducing glutathione biosynthetic capability. *Metab Eng* 2011; 13:426–34. <https://doi.org/10.1016/j.ymben.2011.01.009>.
- Heluane H, Evans MR, Dagher SF. Meta-analysis and functional validation of nutritional requirements of solventogenic clostridia growing under butanol stress conditions and coutilization of D-Glucose and D-Xylose. *Appl Environ Microbiol* 2011;77:4473–85. <https://doi.org/10.1128/AEM.00116-11>.
- Chauvatcharin S, Siripatana C, Seki T, Takagi M, Yoshida T. Metabolism analysis and on-line physiological state diagnosis of acetone-butanol fermentation. *Biotechnol Bioeng* 1998;58:561–71. [https://doi.org/10.1002/\(SICI\)1097-0290\(19980620\)58:6<561::AID-BIT1>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1097-0290(19980620)58:6<561::AID-BIT1>3.0.CO;2-G).
- Pang ZW, Lu W, Zhang H, Liang ZW, Liang JJ, Du LW, et al. Butanol production employing fed-batch fermentation by *Clostridium acetobutylicum* GX01 using alkali-pretreated sugarcane bagasse hydrolysed by enzymes from *Thermoascus aurantiacus* QS 7–2–4. *Bioresour Technol* 2016;212:82–91. <https://doi.org/10.1016/j.biortech.2016.04.013>.
- Kirakosyan G, Bagramyan K, Trchounian A. Redox sensing by *Escherichia coli*: effects of dithiothreitol, a redox reagent reducing disulphides, on bacterial growth. *Biochem Biophys Res Commun* 2004;325:803–6. <https://doi.org/10.1016/j.bbrc.2004.10.119>.
- Soghomanian D, Akopyan K, Trchounian A. pH and oxidation-reduction potential change of environment during growth of lactic acid bacteria: effects of oxidizers and reducers. *Appl Biochem Microbiol* 2011;47:27–31. <https://doi.org/10.1134/S0003683811010157>.
- Davies MB. Reactions of L-ascorbic-acid with transition-metal complexes. *Polyhedron* 1992;11:285–321.
- Zhao X, Ye S, Qi N, Li X, Bao N, Xing D, et al. Mechanisms of enhanced bio-H2 production in *Ethanoligenes harbinense* by L-cysteine supplementation: analyses at growth and gene transcription levels. *Fuel* 2019;252:143–7. <https://doi.org/10.1016/j.fuel.2019.04.037>.
- Gabrielyan L, Sargsyan H, Hakobyan L, Trchounian A. Regulation of hydrogen photoproduction in *Rhodobacter sphaeroides* batch culture by external oxidizers and reducers. *Appl Energy* 2014;131:20–5. <https://doi.org/10.1016/j.apenergy.2014.06.019>.
- Vassilian A, Trchounian A. Environment oxidation-reduction potential and redox sensing by bacteria. *Bact Membr* 2009;163–95.
- Snoep JL, De Graef MR, Teixeira De Mattos MJ, Neijssel OM. Pyruvate catabolism during transient state conditions in chemostat cultures of *Enterococcus faecalis* NCTC 775: importance of internal pyruvate concentrations and NADH/NAD+ ratios. *J Gen Microbiol* 1992;138:2015–20. <https://doi.org/10.1099/00221287-138-10-2015>.
- Sridhar J, Eiteman M. Metabolic flux analysis of *Clostridium thermosuccinogenes*. *Appl Biochem Biotechnol* 2001;94:51–69.
- Schafer FQ, Buettner GR. Redox state and redox environment in biology. In: Forman HJ, Fukuto J, TM, editors. *Signal Transduct*. by React. Oxyg. Nitrogen Species Pathways. Chem. Princ., Springer, Dordrecht; 2003. p. 1–14. https://doi.org/10.1007/0-306-48412-9_1.
- Cleland WW. Dithiothreitol, a new protective reagent for SH groups. *Biochemistry* 1964;3:480–2. <https://doi.org/10.1021/bi00892a002>.
- Loach PA. *Handbook of biochemistry and molecular biology*. Fourth: CRC Press; 2010.

[48] Welch RW, Rudolph FB, Papoutsakis ET. Purification and characterization of the nadh-dependent butanol dehydrogenase from *Clostridium acetobutylicum* (ATCC 824). *Arch Biochem Biophys* 1989;273:309–18.

[49] Dürre P, Kuhn A, Gottwald M, Gottschalk G. Enzymatic investigations on butanol dehydrogenase and butyraldehyde dehydrogenase in extracts of *Clostridium acetobutylicum*. *Appl Microbiol Biotechnol* 1987;26:268–72. <https://doi.org/10.1007/BF00286322>.

[50] Mariano AP, Keshtkar MJ, Atala DIP, Filho FM, Macle MRW, Filho RMI, et al. Energy requirements for butanol recovery using the flash fermentation technology. *Energy Fuels* 2011;25:2347–55. <https://doi.org/10.1021/ef200279w>.

[51] Bowles LK, Ellefson WL. Effects of butanol on *Clostridium acetobutylicum*. *Appl Environ Microbiol* 1985;50:1165–70. <https://doi.org/10.1128/aem.50.5.1165-1170.1985>.

[52] Huang L, Gibbins LN, Forsberg CW. Transmembrane pH gradient and membrane potential in *Clostridium acetobutylicum* during growth under acetogenic and solventogenic conditions. *Appl Environ Microbiol* 1985;50:1043–7. <https://doi.org/10.1128/aem.50.4.1043-1047.1985>.

[53] Terracciano JS, Kashket ER. Intracellular conditions required for initiation of solvent production by *Clostridium acetobutylicum*. *Appl Environ Microbiol* 1986;52: 86–91. <https://doi.org/10.1128/aem.52.1.86-91.1986>.

[54] Rutherford BJ, Dahl RH, Price RE, Szmidt HL, Benke PI, Mukhopadhyay A, et al. Functional genomic study of exogenous n-butanol stress in *Escherichia coli*. *Appl Environ Microbiol* 2010;76:1935–45. <https://doi.org/10.1128/AEM.02323-09>.

[55] Kawasaki S, Watamura Y, Ono M, Watanabe T, Takeda K, Niimura Y. Adaptive responses to oxygen stress in obligatory anaerobes *Clostridium acetobutylicum* and *Clostridium aminovalericum*. *Appl Environ Microbiol* 2005;71:8442–50. <https://doi.org/10.1128/AEM.71.12.8442-8450.2005>.

[56] Riebe O, Fischer RJ, Bahl H. Desulfovirodotoxin of *Clostridium acetobutylicum* functions as a superoxide reductase. *FEBS Lett* 2007;581:5605–10. <https://doi.org/10.1016/j.febslet.2007.11.008>.

[57] Burphan T, Tatip S, Limcharoensuk T, Kangboonruang K, Boonchird C, Auesukaree C. Enhancement of ethanol production in very high gravity fermentation by reducing fermentation-induced oxidative stress in *Saccharomyces cerevisiae*. *Sci Rep* 2018;8:1–11. <https://doi.org/10.1038/s41598-018-31558-4>.

[58] Yue SJ, Huang P, Li S, Jan M, Hu HB, Wang W, et al. Enhanced production of 2-hydroxyphenazine from glycerol by a two-stage fermentation strategy in *Pseudomonas chlororaphis* GP72AN. *J Agric Food Chem* 2020;68:561–6. <https://doi.org/10.1021/acs.jafc.9b05033>.

[59] Dai J, Cheng J, Liang Y, Jiang T, Yuan Y. Regulation of extracellular oxidoreduction potential enhanced (R, R)-2,3-butanediol production by *Paenibacillus polymyxa* CJX518. *Bioresour Technol* 2014;167:433–40. <https://doi.org/10.1016/j.biortech.2014.06.044>.

[60] Krzepilko A, Świecić A, Wawryń J, Zadrag R, Koziot S, Bartosz G, et al. Ascorbate restores lifespan of superoxide-dismutase deficient yeast. *Free Radic Res* 2004;38: 1019–24. <https://doi.org/10.1080/10715760410001717327>.

[61] Huang J, Du Y, Bao T, Lin M, Wang J, Yang S. Bioresource Technology Production of n-butanol from cassava bagasse hydrolysate by engineered *Clostridium* tyrobutyricum overexpressing adh E2: kinetics and cost analysis. *Bioresour Technol* 2019;292:121969. <https://doi.org/10.1016/j.biortech.2019.121969>.

[62] Jin Y, Fang Y, Huang M, Sun J, Huang Y, Gao X, et al. Combination of RNA sequencing and metabolite data to elucidate improved toxic compound tolerance and butanol fermentation of *Clostridium acetobutylicum* from wheat straw hydrolysate by supplying sodium sulfide. *Bioresour Technol* 2015;198:77–86. <https://doi.org/10.1016/j.biortech.2015.08.139>.

[63] Hönicke D, Janssen H, Grimmer C, Ehrenreich A, Lütke-Eversloh T. Global transcriptional changes of *Clostridium acetobutylicum* cultures with increased butanol: acetone ratios. *N Biotechnol* 2012;29:485–93. <https://doi.org/10.1016/j.nbt.2012.01.001>.

[64] Zhao X, Condruz S, Chen J, Jolicoeur M. A quantitative metabolomics study of high sodium response in *Clostridium acetobutylicum* ATCC 824 acetone-butanol-ethanol (ABE) fermentation. *Sci Rep* 2016;6:1–13. <https://doi.org/10.1038/srep28307>.

[65] Lütke-Eversloh T. Application of new metabolic engineering tools for *Clostridium acetobutylicum*. *Appl Microbiol Biotechnol* 2014;98:5823–37. <https://doi.org/10.1007/s00253-014-5785-5>.

[66] Gottwald M, Gottschalk G. The internal pH of *Clostridium acetobutylicum* and its effect on the shift from acid to solvent formation. *Arch Microbiol* 1985;143:42–6. <https://doi.org/10.1007/BF00414766>.

[67] Nicolaou SA, Gaida SM, Papoutsakis ET. A comparative view of metabolite and substrate stress and tolerance in microbial bioprocessing: from biofuels and chemicals, to biocatalysis and bioremediation. *Metab Eng* 2010;12:307–31. <https://doi.org/10.1016/j.ymben.2010.03.004>.

[68] Wang YF, Tian J, Ji ZH, Song MY, Li H. Intracellular metabolic changes of *Clostridium acetobutylicum* and promotion to butanol tolerance during biobutanol fermentation. *Int J Biochem Cell Biol* 2016;78:297–306. <https://doi.org/10.1016/j.biocel.2016.07.031>.

[69] Luo H, Ge L, Zhang J, Zhao Y, Ding J, Li Z, et al. Enhancing butanol production under the stress environments of co-culturing *Clostridium acetobutylicum*/ *Saccharomyces cerevisiae* integrated with exogenous butyrate addition. *PLoS One* 2015;10:1–25. <https://doi.org/10.1371/journal.pone.0141160>.

[70] Wang Q, Venkataramanan KP, Huang H, Papoutsakis ET, Wu CH. Transcription factors and genetic circuits orchestrating the complex, multilayered response of *Clostridium acetobutylicum* to butanol and butyrate stress. *BMC Syst Biol* 2013;7. <https://doi.org/10.1186/1752-0509-7-120>.

[71] Kaino T, Takagi H. Gene expression profiles and intracellular contents of stress protectants in *Saccharomyces cerevisiae* under ethanol and sorbitol stresses. *Appl Microbiol Biotechnol* 2008;79:273–83. <https://doi.org/10.1007/s00253-008-1431-4>.

[72] Broadbent JR, Larsen RL, Deibel V, Steele JL. Physiological and transcriptional response of *Lactobacillus casei* ATCC 334 to acid stress. *J Bacteriol* 2010;192: 2445–58. <https://doi.org/10.1128/JB.01618-09>.

[73] Wu Y, Wang Z, Xin X, Bai F, Xue C. Synergetic engineering of central carbon, energy, and redox metabolisms for high butanol production and productivity by *Clostridium acetobutylicum*. *Ind Eng Chem Res* 2020. <https://doi.org/10.1021/acs.iecr.0c01187>.