

MATH 303 — Measures and Integration
Lecture Notes, Fall 2024

Ethan Ackelsberg

Contents

Part 1. Motivation and Basics of Abstract Measure Theory	1
Chapter 1. Motivating Problems of Measure Theory	3
1. The Problem of Measurement	3
2. Riemann Integration and Jordan Content	3
3. Limits of Integrable Functions	8
4. The Solution of Lebesgue	8
5. Applications of Abstract Measure Theory	9
Chapter Notes	10
Exercises	10
Chapter 2. Measure Spaces	13
1. σ -Algebras	13
2. Measurable Functions	14
3. The Extended Real Numbers and Extended Real-Valued Functions	15
4. Measures	18
Chapter Notes	19
Exercises	19
Chapter 3. Integration Against a Measure	21
1. Integration of Simple Functions	21
2. Integration of Nonnegative Measurable Functions	23
3. Integration of Real and Complex-Valued Functions	26
4. Integral Identities and Inequalities	27
5. Sets of Measure Zero	30
6. The Dominated Convergence Theorem	33
Chapter Notes	34
Exercises	34
Part 2. Constructions of Measures	37
Chapter 4. Taxonomy of Measures	39
1. Properties of Measures	40
2. Finiteness Properties	40
3. Decompositions of Measures	41
Exercises	43
Chapter 5. Lebesgue–Stieltjes Measures	45
1. The π – λ Theorem and Uniqueness of Lebesgue–Stieltjes Measures	46
2. Half-Open Intervals	50
3. Premeasures and Outer Measures	50
4. Lebesgue Measure	56
5. Cantor Measure	57

6. Regularity of Lebesgue–Stieltjes Measures	58
Chapter Notes	60
Exercises	60
Chapter 6. Borel Measures on Locally Compact Hausdorff Spaces	63
1. Locally Compact Hausdorff Spaces	63
2. Radon Measures and the Riesz Representation Theorem	64
3. Topological Lemmas	64
4. Proof of the Riesz Representation Theorem	67
Exercises	69
Chapter 7. Products of Measure Spaces	71
1. Cross-Sectional Product Measures and the Fubini–Tonelli Theorem	72
2. The Maximal Product Measure	76
Exercises	78
Part 3. Additional Topics in Measure Theory	79
Chapter 8. L^p Spaces	81
1. Topological Vector Spaces	81
2. L^p Norms	82
3. Convexity and the Inequalities of Jensen and Minkowski	83
4. Riesz–Fischer Theorem	85
5. Hölder’s Inequality	87
6. Convolutions and Young’s Inequality	89
Exercises	89
Chapter 9. Regularity and Littlewood’s Principles	91
1. The First Principle: Approximation of Measurable Sets	91
2. The Second Principle: L^p Functions are Nearly Continuous	94
3. The Third Principle: Convergent Sequences of Functions are Nearly Uniformly Convergent	96
Chapter Notes	97
Exercises	98
Chapter 10. Differentiation of Measures	99
1. Signed and Complex Measures	99
2. Radon–Nikodym Derivatives	102
3. Complex Borel Measures on \mathbb{R} and their Derivatives	106
4. The Dual of L^p	108
5. The Riesz Representation Theorem, Revisited	110
Chapter Notes	114
Exercises	114
Bibliography	117

Part 1

Motivation and Basics of Abstract Measure Theory

CHAPTER 1

Motivating Problems of Measure Theory

1. The Problem of Measurement

A basic (and very old) problem in mathematics is to compute the size (length, area, volume) of geometric objects. Areas of polygons and circles can be computed by elementary methods. More complicated regions bounded by continuous curves can be attacked with methods from calculus. But what about more general subsets of Euclidean space? Does it always make sense to talk about the (hyper-)volume of a subset of \mathbb{R}^d ? What properties does volume have, and how do we compute it?

We will consider these general questions as the “problem of measurement” in Euclidean space and discuss some approaches to a solution.

2. Riemann Integration and Jordan Content

A good first attempt at solving the problem of measurement comes from the Riemann theory of integration. The basic strategy is to approximate general regions by finite collections of boxes (sets of the form $B = \prod_{i=1}^d [a_i, b_i]$). For such a box B , we declare the volume to be $\text{Vol}(B) = \prod_{i=1}^d (b_i - a_i)$ and use this to define the volume of more general regions. We will now make this idea rigorous.

DEFINITION 1.1: DARBOUX INTEGRATION

Let $B = \prod_{i=1}^d [a_i, b_i]$ be a box in \mathbb{R}^d , and let $f : B \rightarrow \mathbb{R}$ be a bounded function.

- A *Darboux partition* of B is a family of finite sequences $(x_{i,j})_{1 \leq i \leq d, 0 \leq j \leq n_i}$ such that $a_i = x_{i,0} < x_{i,1} < \dots < x_{i,n_i} = b_i$ for each $i \in \{1, \dots, d\}$.




FIGURE 1.1. A Darboux partition in dimension $d = 2$ with $n_1 = 4$ and $n_2 = 6$.

- Given a Darboux partition $P = (x_{i,j})_{1 \leq i \leq d, 0 \leq j \leq n_i}$ of B , the *upper* and *lower Darboux sums of f over B* are given by

$$U_B(f, P) = \sum_{\mathbf{j} \in \prod_{i=1}^d \{1, \dots, n_i\}} \sup_{\mathbf{x} \in B_{\mathbf{j}}} f(\mathbf{x}) \cdot \text{Vol}(B_{\mathbf{j}})$$

and

$$L_B(f, P) = \sum_{\mathbf{j} \in \prod_{i=1}^d \{1, \dots, n_i\}} \inf_{\mathbf{x} \in B_{\mathbf{j}}} f(\mathbf{x}) \cdot \text{Vol}(B_{\mathbf{j}}),$$

where $B_{\mathbf{j}}$ is the box $\prod_{i=1}^d [x_{i,j_i-1}, x_{i,j_i}]$, and $\text{Vol}(B_{\mathbf{j}}) = \prod_{i=1}^d (x_{i,j_i} - x_{i,j_i-1})$ is the volume of $B_{\mathbf{j}}$.




FIGURE 1.2. Upper (red) and lower (blue) Darboux sums of a function f over an interval ($d = 1$).

- The *upper* and *lower Darboux integral of f over B* are

$$U_B(f) = \inf\{U_B(f, P) : P \text{ is a Darboux partition of } B\}$$

and

$$L_B(f) = \sup\{L_B(f, P) : P \text{ is a Darboux partition of } B\}.$$

- The function f is *Darboux integrable over B* if $U_B(f) = L_B(f)$, and their common value is called the *Darboux integral of f over B* and is denoted by $\int_B f(\mathbf{x}) d\mathbf{x}$.

PROPOSITION 1.2

A function f is Darboux integrable if and only if it is Riemann integrable. Moreover, the value of the Darboux integral and the Riemann integral (for a Riemann–Darboux integrable function) are the same.

DEFINITION 1.3

A bounded set $E \subseteq \mathbb{R}^d$ is a *Jordan measurable set* if $\mathbb{1}_E$ is Riemann–Darboux integrable over a box containing E . The *Jordan content* of a Jordan measurable set E is the value $J(E) = \int_B \mathbb{1}_E(\mathbf{x}) d\mathbf{x}$, where B is any box containing E .

Jordan measurable sets include basic geometric objects such as polyhedra, conic sections, regions bounded by finitely many smooth curves/surfaces, etc.

DEFINITION 1.4

A set $S \subseteq \mathbb{R}^d$ is a *simple set* if it is a finite union of boxes $S = \bigcup_{j=1}^k B_j$.

If the boxes B_1, \dots, B_k are disjoint, then the volume of the simple set $S = \bigcup_{j=1}^k B_j$ is $\text{Vol}(S) = \sum_{j=1}^k \text{Vol}(B_j)$. If some of the boxes intersect, then the volume of $S = \bigcup_{j=1}^k B_j$ can be computed using inclusion-exclusion:

$$\text{Vol}(S) = \sum_{j=1}^k \text{Vol}(B_j) - \sum_{1 \leq j_1 < j_2 \leq k} \text{Vol}(B_{j_1} \cap B_{j_2}) + \sum_{1 \leq j_1 < j_2 < j_3 \leq k} \text{Vol}(B_{j_1} \cap B_{j_2} \cap B_{j_3}) - \dots$$

This expression is well-defined, since the intersection of two boxes is again a box. A Jordan measurable set is a set that is “well-approximated” by simple sets, as we will make precise now.

DEFINITION 1.5

For a bounded set $E \subseteq \mathbb{R}^d$, define the *inner* and *outer Jordan content* by

$$J_*(E) = \sup \{ \text{Vol}(S) : S \subseteq E \text{ is a simple set} \}.$$

and

$$J^*(E) = \inf \{ \text{Vol}(S) : S \supseteq E \text{ is a simple set} \}.$$




FIGURE 1.3. Simple sets approximating the inner (red) and outer Jordan content (blue) of a region in dimension $d = 2$. With the red boxes removed from the blue, we get a simple set covering the boundary (in green).

THEOREM 1.6

Let $E \subseteq \mathbb{R}^d$ be a bounded set. The following are equivalent:

- (i) E is Jordan measurable;
- (ii) $J_*(E) = J^*(E)$ (in which case $J(E)$ is equal to this same value);
- (iii) $J^*(\partial E) = 0$.

PROOF. We will prove the $d = 1$ case. The multidimensional case is similar but more notationally cumbersome, so we omit it to avoid additional technical details that would largely obscure the main ideas.

(i) \iff (ii). To establish this equivalence, it suffices to show

$$U_B(\mathbb{1}_E) = J^*(E) \quad \text{and} \quad L_B(\mathbb{1}_E) = J_*(E)$$

for any box (interval) $B \supseteq E$. Let us prove $U_B(\mathbb{1}_E) = J^*(E)$.

CLAIM 1. $U_B(\mathbb{1}_E) \leq J^*(E)$.

Let $\varepsilon > 0$. Then from the definition of the outer Jordan content, there exists a simple set $S \subseteq \mathbb{R}$ such that $E \subseteq S$ and $\text{Vol}(S) < J^*(E) + \varepsilon$. By assumption, B is an interval containing E , so $S \cap B$ is also a simple set containing E , and $\text{Vol}(S \cap B) \leq \text{Vol}(S) < J^*(E) + \varepsilon$. We may therefore assume without loss of generality that $S \subseteq B$. Write $B = [a, b]$ and $S = [a_1, b_1] \sqcup [a_2, b_2] \sqcup \cdots \sqcup [a_n, b_n]$ with $a \leq a_1 \leq b_1 < a_2 \leq b_2 < \cdots < a_n \leq b_n \leq b$. We define a Darboux partition^a P of $[a, b]$ by $P = (x_i)_{i=0}^{2n+1}$ with $x_0 = a$, $x_1 = a_1$, $x_2 = b_1$, \dots , $x_{2n-1} = a_n$, $x_{2n} = b_n$, $x_{2n+1} = b$. Then since $E \subseteq S$, we have

$$\begin{aligned} U_B(\mathbb{1}_E, P) &= \sum_{i=1}^{2n+1} \sup_{x_{i-1} \leq x \leq x_i} \mathbb{1}_E(x) \cdot (x_i - x_{i-1}) \\ &\leq 0 \cdot (a_1 - a) + 1 \cdot (b_1 - a_1) + 0 \cdot (a_2 - b_1) + \cdots + 1 \cdot (b_n - a_n) + 0 \cdot (b - b_n) \\ &= \text{Vol}(S). \end{aligned}$$

Hence, $U_B(\mathbb{1}_E) \leq U_B(\mathbb{1}_E, P) \leq \text{Vol}(S) < J^*(E) + \varepsilon$. This proves the claim.

^aStrictly speaking, this may fail to be a Darboux partition, since some of the points are allowed to coincide. However, the value we compute for $U_B(\mathbb{1}_E, P)$ will be the correct value for the partition where we remove repetitions of the same point.

CLAIM 2. $J^*(E) \leq U_B(\mathbb{1}_E)$.

Let $\varepsilon > 0$. Write $B = [a, b]$. Then there exists a Darboux partition $a = x_0 < x_1 < \cdots < x_n = b$ such that $U_B(\mathbb{1}_E, P) < U_B(\mathbb{1}_E) + \varepsilon$. Let $M_i = \sup_{x_{i-1} \leq x \leq x_i} \mathbb{1}_E(x) \in \{0, 1\}$, and note that, by definition, $U_B(\mathbb{1}_E, P) = \sum_{i=1}^n M_i(x_i - x_{i-1})$. Let $I \subseteq \{1, \dots, n\}$ be the set $I = \{1 \leq i \leq n : M_i = 1\}$, and let $S = \bigcup_{i \in I} [x_{i-1}, x_i]$. Then S is a simple set with length $\text{Vol}(S) = \sum_{i \in I} (x_i - x_{i-1}) = U_B(\mathbb{1}_E, P)$. Moreover, $E \subseteq S$, since S is the union of all intervals that have nonempty intersection with E . Thus, $J^*(E) \leq \text{Vol}(S) = U_B(\mathbb{1}_E, P) < U_B(\mathbb{1}_E) + \varepsilon$.

The identity $L_B(\mathbb{1}_E) = J_*(E)$ is proved similarly.

(ii) \iff (iii). It suffices to prove $J^*(\partial E) = J^*(E) - J_*(E)$. (See Figure 1.3.)

CLAIM 3. $J^*(\partial E) \leq J^*(E) - J_*(E)$.

Let $\varepsilon > 0$. Let S_1 be a simple set such that $E \subseteq S_1$ and $\text{Vol}(S_1) < J^*(E) + \frac{\varepsilon}{2}$. Since S_1 is closed, we have $\overline{E} \subseteq S_1$. Let S_2 be a simple set with $S_2 \subseteq E$ such that $\text{Vol}(S_2) > J_*(E) - \frac{\varepsilon}{2}$. Note that $\text{int}(S_2) \subseteq \text{int}(E)$. Therefore, $S = S_1 \setminus \text{int}(S_2)$ is a simple set and $\partial E = \overline{E} \setminus \text{int}(E) \subseteq S$, so $J^*(\partial E) \leq \text{Vol}(S) = \text{Vol}(S_2) - \text{Vol}(S_1) < J^*(E) - J_*(E) + \varepsilon$. But ε was arbitrary, so we conclude $J^*(\partial E) \leq J^*(E) - J_*(E)$.

CLAIM 4. $J^*(E) - J_*(E) \leq J^*(\partial E)$.

Let $\varepsilon > 0$, and let $S \supseteq \partial E$ be a simple set with $\text{Vol}(S) < J^*(\partial E) + \frac{\varepsilon}{2}$. Write $S = \bigcup_{i=1}^n [a_i, b_i]$ with $a_1 \leq b_1 < a_2 \leq b_2 < \dots < a_n \leq b_n$. Let $[a, b] \subseteq \mathbb{R}$ such that $E \subseteq [a, b]$ and $a < a_1$ and $b < b_n$. For notational convenience, let $b_0 = a$ and $a_{n+1} = b$. Let $I \subseteq \{0, \dots, n\}$ be the collection of indices i such that $(b_i, a_{i+1}) \cap E \neq \emptyset$. For each $i \in I$, we claim that $(b_i, a_{i+1}) \subseteq E$. If not, then (b_i, a_{i+1}) contains a boundary point of E , but $\partial E \subseteq S$, so this is a contradiction. Thus, $S' = \bigcup_{i \in I} [b_i, a_{i+1}]$ is a simple set with $\text{int}(S') \subseteq E$. Shrinking slightly each interval in S' , we obtain a simple set

$$S'' = \bigcup_{i \in I} \left[b_i + \frac{\varepsilon}{4(n+1)}, a_{i+1} - \frac{\varepsilon}{4(n+1)} \right]$$

such that $S'' \subseteq E$. Moreover, $\text{Vol}(S'') \geq \text{Vol}(S') - \frac{\varepsilon}{2(n+1)}|I| \geq \text{Vol}(S') - \frac{\varepsilon}{2}$. Noting that $S \cup S'$ is a simple set containing E , we arrive at the inequality

$$J^*(E) - J_*(E) \leq \text{Vol}(S \cup S') - \text{Vol}(S'') = \text{Vol}(S) + \text{Vol}(S') - \text{Vol}(S'') < J^*(\partial E) + \varepsilon.$$

This completes the proof of Theorem 1.6. □

EXAMPLE 1.7

The sets $\mathbb{Q} \cap [0, 1]$ and $[0, 1] \setminus \mathbb{Q}$ are not Jordan measurable (see Exercise 1.4).

In addition to the above example, there are many other “nice” sets that are not Jordan measurable. There are, for instance, bounded open sets in \mathbb{R} that are not Jordan measurable. We will work out one such example in detail.

EXAMPLE 1.8

The complement U of the fat Cantor set (also known as the Smith–Volterra–Cantor set) $K \subseteq [0, 1]$ is Jordan non-measurable. We construct K iteratively, starting from $[0, 1]$, by removing intervals of length 4^{-n} at step n . In other words, at step n , we remove an interval of length 4^{-n} around each rational point with denominator 2^n .



FIGURE 1.4. Iterative construction of the fat Cantor set.

Let

$$U = \bigcup_{n=0}^{\infty} \bigcup_{j=1}^{2^n} \left(\frac{2j+1}{2^{n+1}} - \frac{1}{2 \cdot 4^{n+1}}, \frac{2j+1}{2^{n+1}} + \frac{1}{2 \cdot 4^{n+1}} \right).$$

Then $K = [0, 1] \setminus U$. The inner Jordan content of U is

$$J_*(U) = \sum_{n=0}^{\infty} \sum_{j=1}^{2^n} \text{Len} \left(\frac{2j+1}{2^{n+1}} - \frac{1}{2 \cdot 4^{n+1}}, \frac{2j+1}{2^{n+1}} + \frac{1}{2 \cdot 4^{n+1}} \right) = \sum_{n=0}^{\infty} 2^n \cdot \frac{1}{4^{n+1}} = \frac{1}{4} \sum_{n=0}^{\infty} 2^{-n} = \frac{1}{2}.$$

However, $\overline{U} = [0, 1]$ (since U contains every rational number whose denominator is a power of 2), so the outer Jordan content of U is $J^*(U) = J^*([0, 1]) = 1$.

3. Limits of Integrable Functions

You may recall from the theory of Riemann integration that *uniform* limits of Riemann integrable functions are Riemann integrable, and one may in this case interchange the order of taking limits and computing the integral. More precisely:

THEOREM 1.9

Let B be a box in \mathbb{R}^d . Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of Riemann integrable functions on B , and suppose f_n converges uniformly to a function $f : B \rightarrow \mathbb{R}$. Then f is Riemann integrable, and

$$\int_B f(\mathbf{x}) \, d\mathbf{x} = \lim_{n \rightarrow \infty} \int_B f_n(\mathbf{x}) \, d\mathbf{x}.$$

One of the deficiencies of the Riemann–Darboux–Jordan approach to integration and measurement is that pointwise (non-uniform) limits do not share this property.

EXAMPLE 1.10

Enumerate the set $\mathbb{Q} \cap [0, 1] = \{q_1, q_2, \dots\}$. Let $f_n : [0, 1] \rightarrow [0, 1]$ be the function

$$f_n(x) = \begin{cases} 1, & \text{if } x \in \{q_1, \dots, q_n\} \\ 0, & \text{otherwise.} \end{cases}$$

Then f_n is Riemann integrable and $f_n \rightarrow \mathbb{1}_{\mathbb{Q} \cap [0, 1]}$ pointwise, but $\mathbb{1}_{\mathbb{Q} \cap [0, 1]}$ is not Riemann integrable.

Since analysis so often deals with limits, it is desirable to develop a theory of integration that accommodates pointwise limits. The Lebesgue measure and Lebesgue integral resolve this shortcoming.

4. The Solution of Lebesgue

The Jordan non-measurable set in Example 1.8 appears to have a sensible notion of “length.” Indeed, the complement U , being a disjoint union of intervals, could be reasonably assigned as a “length” the sum of the lengths of the (countably many) intervals of which it is made. This produces a value of $\frac{1}{2}$ for the length of U , and so we should take K to also have length $\frac{1}{2}$, since $K \sqcup U = [0, 1]$ is an interval of length 1. The feature that U is a disjoint union of intervals turns out to not be any special feature of U at all but instead a general feature of open sets in \mathbb{R} .

PROPOSITION 1.11

Let $U \subseteq \mathbb{R}$ be an open set. Then U can be expressed as a countable disjoint union of open intervals.

PROOF. Exercise 1.1. □

By Proposition 1.11, it seems reasonable to define the length of an open set $U \subseteq \mathbb{R}$ as follows. Write $U = (a_1, b_1) \sqcup (a_2, b_2) \sqcup \dots$ as a disjoint union of open intervals, and define its length as $(b_1 - a_1) + (b_2 - a_2) + \dots$. Then open sets may play the role that simple sets played in the definition of the Jordan content, and this leads to the Lebesgue measure.

REMARK. In higher dimensions, Proposition 1.11 needs to be modified, but one can still reasonably talk about the d -dimensional volume of open sets in \mathbb{R}^d . See Exercises 1.2 and 1.3.

DEFINITION 1.12

Let $E \subseteq \mathbb{R}^d$.

- The *outer Lebesgue measure of E* is the quantity

$$\lambda^*(E) = \inf \{\text{Vol}(U) : U \supseteq E \text{ is open}\}$$

$$= \inf \left\{ \sum_{j=1}^{\infty} \text{Vol}(B_j) : B_1, B_2, \dots \text{ are boxes, and } E \subseteq \bigcup_{j=1}^{\infty} B_j \right\}.$$

- The set E is *Lebesgue measurable* (with *Lebesgue measure* $\lambda(E) = \lambda^*(E)$) if for every $\varepsilon > 0$, there exists an open set $U \subseteq \mathbb{R}^d$ such that $E \subseteq U$ and $\lambda^*(U \setminus E) < \varepsilon$.

PROPOSITION 1.13

If $E \subseteq \mathbb{R}^d$ is Jordan measurable, then E is Lebesgue measurable and $J(E) = \lambda(E)$.

The family of Lebesgue measurable sets is much larger than the family of Jordan measurable sets. Among the several nice properties of the Lebesgue measure (and abstract measures) that we will see later in the course are:

PROPOSITION 1.14

- (1) If $(E_n)_{n \in \mathbb{N}}$ are Lebesgue measurable sets, then $\bigcup_{n=1}^{\infty} E_n$ and $\bigcap_{n=1}^{\infty} E_n$ are Lebesgue measurable.
- (2) If $(E_n)_{n \in \mathbb{N}}$ are pairwise disjoint and Lebesgue measurable, then $\lambda(\bigcup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \lambda(E_n)$.
- (3) If $E_1 \subseteq E_2 \subseteq \dots \subseteq \mathbb{R}^d$ are Lebesgue measurable sets, then $\lambda(\bigcup_{n=1}^{\infty} E_n) = \lim_{n \rightarrow \infty} \lambda(E_n)$.
- (4) If $E_1 \supseteq E_2 \supseteq \dots$ are Lebesgue measurable subsets of \mathbb{R}^d and $\lambda(E_1) < \infty$, then $\lambda(\bigcap_{n=1}^{\infty} E_n) = \lim_{n \rightarrow \infty} \lambda(E_n)$.

5. Applications of Abstract Measure Theory

The mathematical language and tools encompassed in measure theory play a foundational role in many other areas of mathematics. A highly abbreviated sampling follows.

PROBABILITY THEORY. Measure theory provides the axiomatic foundations of probability theory, providing rigorous notions of *random variables* and *probabilities of events*. Important limit laws (the law of large numbers and central limit theorem, for example) are phrased mathematically using measure-theoretic notions of convergence.


FOURIER ANALYSIS. Periodic (say, continuous or Riemann-integrable) functions on the real line have corresponding Fourier series representations $f(x) \sim \sum_{n \in \mathbb{Z}} \widehat{f}(n) e^{2\pi i n x}$. The functions $e^{2\pi i n x}$ are orthonormal, and Parseval's identity gives $\sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2 = \int_0^1 |f(x)|^2 dx$. Given a sequence $(a_n)_{n \in \mathbb{N}}$, one may ask whether $\sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ is the Fourier expansion of some function f , and if so, what properties does f have? Another natural question is whether the series $\sum_{n \in \mathbb{Z}} \widehat{f}(n) e^{2\pi i n x}$ actually converges to the function f , and if so, in which sense? Both of these questions are properly answered in a measure-theoretic framework. If one is interested in decomposing functions defined on other groups (for instance, on compact abelian groups) into their Fourier series, then one also

needs to develop a method of integrating functions on groups in order to compute Fourier coefficients and make sense of Parseval's identity.

FUNCTIONAL ANALYSIS AND OPERATOR THEORY. When one studies familiar concepts from linear algebra in infinite-dimensional spaces, measures become unavoidable for many tasks. For example, versions of the spectral theorem (generalizing the representation of suitable matrices in terms of their eigenvalues and eigenvectors) for operators on infinite-dimensional spaces require the abstract notion of a measure.

ERGODIC THEORY. Ergodic theory was developed to study the long-term statistical behavior of dynamical (time-dependent) systems, providing a framework to resolve important problems in physics related to the “ergodic hypothesis” in thermodynamics and the “stability” of the solar system. It turns out that the appropriate mathematical formalism for understanding these problems comes from abstract measure theory.

FRACTAL GEOMETRY. Self-similar geometric objects such as the Koch snowflake, Sierpiński carpet, and the middle-thirds Cantor set (see Figure 1.5) can be meaningfully assigned a notion of “dimension” that can take a non-integer value. How does one determine the dimension of a fractal object? There are several different approaches to dimension, but one of the most popular is the *Hausdorff dimension*, which relies on a family of measures that interpolate between the integer-dimensional Lebesgue measures.



Chapter Notes

This introductory chapter is heavily influenced by the book of Tao [10] on measure theory. Many of the results in this chapter are discussed in greater detail in [10, Section 1.1].

Exercises

1.1 Let $U \subseteq \mathbb{R}$ be an open set. Show that U can be written as a disjoint union of countably many open intervals.

1.2 Let $U = \{(x, y) : x^2 + y^2 < 1\} \subseteq \mathbb{R}^2$ be the open unit disk. Show that U cannot be expressed as a disjoint union of countably many open boxes.

1.3 Let $U \subseteq \mathbb{R}^d$ be an open set. Show that U can be written as a disjoint union of countably many half-open boxes (i.e., sets of the form $B = \prod_{i=1}^d [a_i, b_i)$).

1.4 Show that $J^*(\mathbb{Q} \cap [0, 1]) = J^*([0, 1] \setminus \mathbb{Q}) = 1$, and $J_*(\mathbb{Q} \cap [0, 1]) = J_*([0, 1] \setminus \mathbb{Q}) = 0$.

1.5 (Area Interpretation of the Riemann Integral) Let $[a, b]$ be an interval and $f : [a, b] \rightarrow [0, \infty)$ a bounded function. Show that f is Riemann integrable if and only if the set

$$E_f = \{(x, t) : a \leq x \leq b, 0 \leq t \leq f(x)\}$$

is a Jordan measurable set in \mathbb{R}^2 , in which case

$$\int_a^b f(x) \, dx = J(E_f).$$

1.6 Give an example to show that the statement

$$\lambda^*(E) = \sup_{U \subseteq E, U \text{ open}} \lambda^*(U).$$

is false.

CHAPTER 2

Measure Spaces

1. σ -Algebras

Before defining measures, we must determine which subsets of a given set X we would like to be able to measure. The full set X should be measurable, and we should allow ourselves to perform the basic set-theoretic operations (complements, unions, and intersections). Allowing *finite* unions and intersections produces an *algebra* of sets. Algebras are a very useful notion, but (as with the Jordan content discussed in the previous chapter) they are insufficient for appropriately handling limits. We will therefore upgrade from algebras to σ -*algebras*:

DEFINITION 2.1

Let X be a set. A σ -*algebra* on X is a family $\mathcal{B} \subseteq \mathcal{P}(X)$ of subsets of X with the following properties:

- $X \in \mathcal{B}$;
- If $B \in \mathcal{B}$, then $X \setminus B \in \mathcal{B}$;
- If $(B_n)_{n \in \mathbb{N}}$ is a countable family of elements of \mathcal{B} , then $\bigcup_{n \in \mathbb{N}} B_n \in \mathcal{B}$.

REMARK. In the definition of a σ -algebra, we have made no explicit mention of intersections. However, by De Morgan's laws, we can also generate the countable intersection of sets: $\bigcap_{n \in \mathbb{N}} B_n = X \setminus (\bigcup_{n \in \mathbb{N}} (X \setminus B_n))$.

EXAMPLE 2.2

Some examples of σ -algebras include the following:

- For any set X , the power set $\mathcal{P}(X)$ is a σ -algebra, as is the pair $\{\emptyset, X\}$.
- The family $\mathcal{B} = \{B \subseteq \mathbb{R} : \text{either } B \text{ or } \mathbb{R} \setminus B \text{ is countable}\}$ of countable and co-countable subsets of \mathbb{R} is a σ -algebra.
- Unions of unit-length intervals in \mathbb{R} form a σ -algebra $\mathcal{B} = \{\bigcup_{n \in S} [n, n+1) : S \subseteq \mathbb{Z}\}$.

PROPOSITION 2.3

Suppose $(\mathcal{B}_i)_{i \in I}$ is a family of σ -algebras on X . Then $\bigcap_{i \in I} \mathcal{B}_i$ is a σ -algebra.

PROOF. Let $\mathcal{B} = \bigcap_{i \in I} \mathcal{B}_i$.

For every $i \in I$, we have $X \in \mathcal{B}_i$, so $X \in \mathcal{B}$.

Suppose $B \in \mathcal{B}$. Then $B \in \mathcal{B}_i$ for every $i \in I$, so $X \setminus B \in \mathcal{B}_i$ for every $i \in I$. Hence, $X \setminus B \in \mathcal{B}$.

Let $(B_n)_{n \in \mathbb{N}}$ be a countable family of sets in \mathcal{B} . For each $i \in I$, the sets $(B_n)_{n \in \mathbb{N}}$ belong to \mathcal{B}_i , so $\bigcup_{n \in \mathbb{N}} B_n \in \mathcal{B}_i$. Therefore, $\bigcup_{n \in \mathbb{N}} B_n \in \mathcal{B}$. \square

DEFINITION 2.4

The *σ -algebra generated by a family* $\mathcal{S} \subseteq \mathcal{P}(X)$ is the smallest σ -algebra containing \mathcal{S} , denoted by $\sigma(\mathcal{S})$.

REMARK. Note that $\sigma(\mathcal{S})$ is well-defined by Proposition 2.3:

$$\sigma(\mathcal{S}) = \bigcap \{\mathcal{B} : \mathcal{B} \text{ is a } \sigma\text{-algebra on } X, \mathcal{S} \subseteq \mathcal{B}\}.$$

In topological spaces (such as the real line), we will often consider the σ -algebra generated by the topology.

DEFINITION 2.5

Let (X, τ) be a topological space. The *Borel σ -algebra* is the σ -algebra generated by the open subsets of X , i.e. $\text{Borel}(X) = \sigma(\tau)$.

Borel sets can be placed in a hierarchy in terms of their level of complexity. At the simplest level are the open (G) and closed (F) sets. Next come countable intersections of open sets (G_δ sets) and countable unions of closed sets (F_σ sets) and so on.

$$\begin{array}{ccccccccccc} \Sigma_1^0 = G & \xrightarrow{\cap} & \Pi_2^0 = G_\delta & \xrightarrow{\cup} & \Sigma_3^0 = G_{\delta\sigma} & \xrightarrow{\cap} & \Pi_4^0 = G_{\delta\sigma\delta} & \xrightarrow{\cup} & \dots & \xrightarrow{\cup} & \Sigma_\omega^0 & \xrightarrow{\cap} & \dots \\ \downarrow c & & \downarrow c & & \downarrow c & & \downarrow c & & & & \downarrow c & & \\ \Pi_1^0 = F & \xrightarrow{\cup} & \Sigma_2^0 = F_\sigma & \xrightarrow{\cap} & \Pi_3^0 = F_{\sigma\delta} & \xrightarrow{\cup} & \Sigma_4^0 = F_{\sigma\delta\sigma} & \xrightarrow{\cap} & \dots & \xrightarrow{\cap} & \Pi_\omega^0 & \xrightarrow{\cup} & \dots \end{array}$$

FIGURE 2.1. The Borel hierarchy for subsets of a topological space.

The placement of a (Borel) set within the Borel hierarchy is a useful notion of “complexity” for sets. Intuitively speaking, if a set is lower down in the Borel hierarchy, then it is in some sense easier to define than a set higher up the hierarchy. Determining where sets occur in the Borel hierarchy (or if they are Borel at all) is a common theme in an area of mathematical logic known as *descriptive set theory*. We will largely not concern ourselves with such problems in this course, but some suggested additional reading appears at the end of this chapter for those who are interested.

In our development of the abstract theory of measures (where we may not even have a topology to work with), our object of study will be arbitrary sets X equipped with a σ -algebra.

DEFINITION 2.6

A *measurable space* is a pair (X, \mathcal{B}) , where X is a set and \mathcal{B} is a σ -algebra on X . Elements of the σ -algebra \mathcal{B} are called *measurable sets*.

2. Measurable Functions

Recall that a function $f : X \rightarrow Y$ from one topological space to another is continuous if the preimage of every open set in Y is open in X . Measurable functions are defined analogously, but with “open” replaced by “measurable.”

DEFINITION 2.7

Let (X, \mathcal{B}) and (Y, \mathcal{C}) be measurable spaces. A function $f : X \rightarrow Y$ is *measurable* if for every $C \in \mathcal{C}$, one has $f^{-1}(C) \in \mathcal{B}$.

Some basic properties of measurable functions that will be used frequently are as follows:

PROPOSITION 2.8

- (1) Let (X, \mathcal{B}) , (Y, \mathcal{C}) , and (Z, \mathcal{D}) be measurable spaces. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be measurable functions. Then $g \circ f : X \rightarrow Z$ is measurable.
- (2) Let (X, \mathcal{B}) and (Y, \mathcal{C}) be measurable spaces, and let $f : X \rightarrow Y$. Suppose $\mathcal{S} \subseteq \mathcal{P}(Y)$ is a family of sets such that $\sigma(\mathcal{S}) = \mathcal{C}$. If $f^{-1}(S) \in \mathcal{B}$ for every $S \in \mathcal{S}$, then f is a measurable function.
- (3) Suppose X and Y are topological spaces and $\mathcal{B} = \text{Borel}(X)$ and $\mathcal{C} = \text{Borel}(Y)$ are the Borel σ -algebras on X and Y respectively. Then every continuous function $f : X \rightarrow Y$ is measurable.

PROOF. (1) Let $D \in \mathcal{D}$. Since g is measurable, we have $C = g^{-1}(D) \in \mathcal{C}$. Then since f is measurable, $B = f^{-1}(C) \in \mathcal{B}$. But $B = f^{-1}(g^{-1}(D)) = (g \circ f)^{-1}(D)$, so $g \circ f$ is measurable.

(2) Let $\mathcal{F} = \{E \subseteq Y : f^{-1}(E) \in \mathcal{B}\}$. We claim that \mathcal{F} is a σ -algebra. Then since $\mathcal{S} \subseteq \mathcal{F}$, we conclude that $\mathcal{C} = \sigma(\mathcal{S}) \subseteq \mathcal{F}$, so f is measurable. Let us now prove the claim:

- $f^{-1}(Y) = X \in \mathcal{B}$, so $Y \in \mathcal{F}$.
- Suppose $E \in \mathcal{F}$. Then $f^{-1}(Y \setminus E) = X \setminus \underbrace{f^{-1}(E)}_{\in \mathcal{B}} \in \mathcal{B}$, so $Y \setminus E \in \mathcal{F}$.
- Suppose $E_1, E_2, \dots \in \mathcal{F}$, and let $E = \bigcup_{n \in \mathbb{N}} E_n$. Then

$$f^{-1}(E) = \bigcup_{n \in \mathbb{N}} \underbrace{f^{-1}(E_n)}_{\in \mathcal{B}} \in \mathcal{B},$$

so $E \in \mathcal{F}$.

This proves that \mathcal{F} is a σ -algebra on Y .

(3) This follows from (1) by taking \mathcal{S} to be the collection of open sets in Y . □

3. The Extended Real Numbers and Extended Real-Valued Functions

One obtains an important class of measurable functions when one considers functions defined on a measurable space taking real values. For many applications and in order to account more fully for limits of functions, it is often convenient to work with the slightly more general concept of *extended* real-valued functions.

DEFINITION 2.9

The *extended real numbers* are the set $[-\infty, \infty] = \mathbb{R} \cup \{\infty, -\infty\}$ with the following topological and algebraic properties:

- The topology on $[-\infty, \infty]$ is generated by open intervals (a, b) with $a, b \in \mathbb{R}$ and sets of the form $(a, \infty] = (a, \infty) \cup \{\infty\}$ and $[-\infty, b) = (-\infty, b) \cup \{-\infty\}$ for $a, b \in \mathbb{R}$.

- Addition is extended as a commutative operation with $\infty + x = \infty$ and $-\infty + x = -\infty$ for real numbers $x \in \mathbb{R}$. For addition of two infinite quantities, we define $\infty + \infty = \infty$ and $-\infty + (-\infty) = -\infty$. However, $-\infty + \infty$ is undefined.
- Multiplication is also extended as a commutative operation with the properties

$$x \in (0, \infty) \implies \infty \cdot x = \infty \quad \text{and} \quad -\infty \cdot x = -\infty;$$

$$x \in (-\infty, 0) \implies \infty \cdot x = -\infty \quad \text{and} \quad -\infty \cdot x = \infty.$$

By convention, we define $\infty \cdot 0 = -\infty \cdot 0 = 0$. Multiplication of infinities is defined by $\infty \cdot \infty = (-\infty) \cdot (-\infty) = \infty$, and $-\infty \cdot \infty = -\infty$.

The topology we have defined on $[-\infty, \infty]$ is the *two-point compactification* of \mathbb{R} . You will check in the exercises (Exercise 2.5) that $[-\infty, \infty]$ is indeed a compact space (that is homeomorphic to a closed interval, say $[0, 1]$). The algebraic operations on $[-\infty, \infty]$ are all as one would expect, with one exception: $\infty \cdot 0$ is often considered as an “indeterminate form”, but here we have given it a definite value of 0. The reason for this convention is the following proposition, which you will also prove in the exercises:

PROPOSITION 2.10

Let $(x_n)_{n \in \mathbb{N}}$ be a sequence in $[-\infty, \infty]$, and let $c \in \mathbb{R}$. If $(x_n)_{n \in \mathbb{N}}$ converges to an extended real number, then the sequence $(cx_n)_{n \in \mathbb{N}}$ also converges, and

$$\lim_{n \rightarrow \infty} (cx_n) = c \cdot \lim_{n \rightarrow \infty} x_n. \quad (2.1)$$

PROOF. Exercise 2.6. □

In order to have the desirable property (2.1), one has no choice but to define $\infty \cdot 0 = 0$: by taking the sequence $x_n = n$, we have

$$0 \cdot \infty = 0 \cdot \lim_{n \rightarrow \infty} n = \lim_{n \rightarrow \infty} (0 \cdot n) = 0.$$

WARNING: Property (2.1) does not hold for $c \in \{\infty, -\infty\}$, as can be seen by taking a sequence $(x_n)_{n \in \mathbb{N}}$ that converges to 0.

We say that an extended real-valued function $f : X \rightarrow [-\infty, \infty]$ defined on a measurable space (X, \mathcal{B}) is *\mathcal{B} -measurable* (or simply *measurable*) if it is measurable as a function between the measurable spaces (X, \mathcal{B}) and $([-\infty, \infty], \text{Borel}([-\infty, \infty]))$. Since we will always take the same σ -algebra on $[-\infty, \infty]$, we omit explicit reference to the Borel σ -algebra when discussing measurable extended real-valued functions.

PROPOSITION 2.11

Let (X, \mathcal{B}) be a measurable space.

- (1) Let $f : X \rightarrow [-\infty, \infty]$. The following are equivalent:
 - f is measurable;
 - for every $c \in \mathbb{R}$, $f^{-1}((c, \infty]) \in \mathcal{B}$;
 - for every $c \in \mathbb{R}$, $f^{-1}([c, \infty]) \in \mathcal{B}$;
 - for every $c \in \mathbb{R}$, $f^{-1}((-\infty, c)) \in \mathcal{B}$;
 - for every $c \in \mathbb{R}$, $f^{-1}((-\infty, c]) \in \mathcal{B}$.
- (2) Suppose $(f_n)_{n \in \mathbb{N}}$ is a sequence of measurable functions from X to $[-\infty, \infty]$. The following functions are also measurable:

- (a) $\sup_{n \in \mathbb{N}} f_n$;
- (b) $\inf_{n \in \mathbb{N}} f_n$;
- (c) $\limsup_{n \rightarrow \infty} f_n$;
- (d) $\liminf_{n \rightarrow \infty} f_n$.

(3) Suppose $f, g : X \rightarrow \mathbb{R}$ are measurable functions. Then $f + g$ and $f \cdot g$ are measurable.

NOTATION. For convenience, we will often write sets of the form $f^{-1}((c, \infty])$ as $\{f > c\}$ and similarly for $\{f \geq c\}$, $\{f < c\}$, and $\{f \leq c\}$.

PROOF OF PROPOSITION 2.11. (1) By Proposition 2.8(2), it suffices to check that each of the relevant collections of intervals generates the Borel σ -algebra on $[-\infty, \infty]$. Let us show that the collection of intervals $(c, \infty]$ for $c \in \mathbb{R}$ generates the Borel σ -algebra. All of the other proofs are similar, so we omit them.

Let $\mathcal{S} = \{(c, \infty] : c \in \mathbb{R}\}$. Note that every element of \mathcal{S} is open in $[-\infty, \infty]$, so $\sigma(\mathcal{S}) \subseteq \text{Borel}([-\infty, \infty])$. On the other hand, we can write $(a, b] = (a, \infty] \setminus (b, \infty]$ for $a, b \in \mathbb{R}, a < b$. Every open set in \mathbb{R} is a countable (disjoint) union of such intervals, so every open subset of \mathbb{R} is contained in $\sigma(\mathcal{S})$. We obtain the additional open sets in $[-\infty, \infty]$ from the rays $(c, \infty] \in \mathcal{S}$ and

$$[-\infty, c) = \bigcup_{n \in \mathbb{N}} \left[-\infty, c - \frac{1}{n} \right] = \bigcup_{n \in \mathbb{N}} \left([-\infty, \infty] \setminus \left(c - \frac{1}{n}, \infty \right] \right) \in \sigma(\mathcal{S}).$$

Thus, $\text{Borel}([-\infty, \infty]) \subseteq \sigma(\mathcal{S})$.

(2) We will use (1).

(a) Let $f = \sup_{n \in \mathbb{N}} f_n$. Note that $\{f > c\} = \bigcup_{n \in \mathbb{N}} \{f_n > c\}$. Each of the sets $\{f_n > c\}$ belongs to \mathcal{B} , so $\{f > c\} \in \mathcal{B}$.

(b) Similarly to (a), letting $f = \inf_{n \in \mathbb{N}} f_n$, we may express $\{f < c\} = \bigcup_{n \in \mathbb{N}} \underbrace{\{f_n < c\}}_{\in \mathcal{B}} \in \mathcal{B}$.

(c) Recall that $\limsup_{n \rightarrow \infty} f_n = \inf_{k \in \mathbb{N}} \sup_{n \geq k} f_n$, so measurability of $\limsup_{n \rightarrow \infty} f_n$ follows from (a) and (b).

(d) Similar to (c): $\liminf_{n \rightarrow \infty} f_n = \sup_{k \in \mathbb{N}} \inf_{n \geq k} f_n$.

(3) Let $A : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $M : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the maps $A(x, y) = x + y$ and $M(x, y) = xy$. Both of the maps A and M are continuous and therefore (Borel) measurable. Moreover, $(f + g)(x) = A(f(x), g(x))$ and $(f \cdot g)(x) = M(f(x), g(x))$. Since the composition of measurable maps is measurable (see Proposition 2.8(1)), it suffices to prove $h : x \mapsto (f(x), g(x))$ is a measurable function from X to \mathbb{R}^2 . By Proposition 2.8(2), we only need to check preimages of sets generating the Borel σ -algebra on \mathbb{R}^2 . For convenience, we will take the boxes $[a, b] \times [c, d]$ (the first homework problem was to show that every open set in \mathbb{R}^2 is a countable (disjoint) union of such boxes, so they generate the Borel σ -algebra). Observe that

$$h^{-1}([a, b] \times [c, d]) = f^{-1}([a, b]) \cap g^{-1}([c, d]) \in \mathcal{B},$$

since f and g are measurable, so h is indeed a measurable function. □

EXAMPLE 2.12

Let (X, \mathcal{B}) be a measurable space and $E \subseteq X$. The function $\mathbb{1}_E$ is measurable if and only if $E \in \mathcal{B}$.

4. Measures

We are now prepared to define measures on abstract measurable spaces.

DEFINITION 2.13

Let (X, \mathcal{B}) be a measurable space. A *measure* on (X, \mathcal{B}) is a function $\mu : \mathcal{B} \rightarrow [0, \infty]$ such that

- $\mu(\emptyset) = 0$;
- COUNTABLE ADDITIVITY: for any sequence $(E_n)_{n \in \mathbb{N}}$ of pairwise disjoint elements of \mathcal{B} , one has $\mu(\bigsqcup_{n \in \mathbb{N}} E_n) = \sum_{n \in \mathbb{N}} \mu(E_n)$.

The triple (X, \mathcal{B}, μ) is called a *measure space*.

Nontrivial examples of measures take some effort to construct, and we will spend significant portions of the course discussing different methods for constructing interesting measures. However, there are a few immediate examples that do not require complicated constructions.

EXAMPLE 2.14

Examples of measures include:

- For any set X , the *counting measure* is a measure defined on the σ -algebra $\mathcal{P}(X)$ by $\mu(E) = |E|$ if E is a finite set and $\mu(E) = \infty$ if E is an infinite set.
- Given a point $x \in X$, the *Dirac measure* defined on $\mathcal{P}(X)$ is the measure $\delta_x(E) = 1$ if $x \in E$ and $\delta_x(E) = 0$ if $x \notin E$.

We will use the following basic properties of measures frequently throughout this course:

PROPOSITION 2.15

Let (X, \mathcal{B}, μ) be a measure space.

- (1) MONOTONICITY: For any $A, B \in \mathcal{B}$, if $A \subseteq B$, then $\mu(A) \leq \mu(B)$.
- (2) COUNTABLE SUB-ADDITIONITY: For any sequence $(E_n)_{n \in \mathbb{N}}$ in \mathcal{B} ,

$$\mu\left(\bigcup_{n \in \mathbb{N}} E_n\right) \leq \sum_{n \in \mathbb{N}} \mu(E_n).$$

- (3) CONTINUITY FROM BELOW: If $E_1 \subseteq E_2 \subseteq \dots \in \mathcal{B}$, then

$$\mu\left(\bigcup_{n \in \mathbb{N}} E_n\right) = \lim_{n \rightarrow \infty} \mu(E_n).$$

- (4) CONTINUITY FROM ABOVE: If $E_1 \supseteq E_2 \supseteq \dots \in \mathcal{B}$ and $\mu(E_1) < \infty$, then

$$\mu\left(\bigcap_{n \in \mathbb{N}} E_n\right) = \lim_{n \rightarrow \infty} \mu(E_n).$$

PROOF. (1) Write $B = A \sqcup (B \setminus A)$. Then $\mu(B) = \mu(A) + \mu(B \setminus A) \geq \mu(A)$, since μ takes nonnegative values.

(2) Define a new sequence of sets E'_n by $E'_1 = E_1$ and $E'_n = E_n \setminus \bigcup_{j=1}^{n-1} E_j$ for $n \geq 2$. Then the sets $(E'_n)_{n \in \mathbb{N}}$ are pairwise disjoint and satisfy $E'_n \subseteq E_n$ and $\bigsqcup_{n \in \mathbb{N}} E'_n = \bigcup_{n \in \mathbb{N}} E_n$. Therefore,

$$\mu \left(\bigcup_{n \in \mathbb{N}} E_n \right) = \mu \left(\bigsqcup_{n \in \mathbb{N}} E'_n \right) = \sum_{n \in \mathbb{N}} \mu(E'_n) \leq \sum_{n \in \mathbb{N}} \mu(E_n),$$

where in the last step we have applied monotonicity of μ (property (1)).

(3) Let $E'_1 = E_1$ and $E'_n = E_n \setminus E_{n-1}$ for $n \geq 2$. For convenience, we will set $E_0 = \emptyset$ so that we also have $E'_1 = E_1 \setminus E_0$. Then

$$\mu \left(\bigcup_{n \in \mathbb{N}} E_n \right) = \mu \left(\bigsqcup_{n \in \mathbb{N}} E'_n \right) = \sum_{n \in \mathbb{N}} \mu(E'_n) \stackrel{(*)}{=} \sum_{n \in \mathbb{N}} (\mu(E_n) - \mu(E_{n-1})) \stackrel{(**)}{=} \lim_{n \rightarrow \infty} \mu(E_n).$$

The step $(*)$ uses additivity of μ , and $(**)$ comes from the telescoping of the sum.

(4) Define a new sequence $A_n = E_1 \setminus E_n$. Then $\emptyset = A_1 \subseteq A_2 \subseteq \dots$, so

$$\mu \left(\bigcup_{n \in \mathbb{N}} A_n \right) = \lim_{n \rightarrow \infty} \mu(A_n)$$

by (3). But $\bigcup_{n \in \mathbb{N}} A_n = E_1 \setminus \bigcap_{n \in \mathbb{N}} E_n$, so

$$\mu(E_1) - \mu \left(\bigcap_{n \in \mathbb{N}} E_n \right) = \mu \left(\bigcup_{n \in \mathbb{N}} A_n \right) = \lim_{n \rightarrow \infty} \mu(A_n) = \mu(E_1) - \lim_{n \rightarrow \infty} \mu(E_n),$$

whence we deduce that (4) holds, since $\mu(E_1) < \infty$. □

EXAMPLE 2.16

Property (4) may fail if $\mu(E_1) = \infty$. Let $X = \mathbb{N}$, $\mathcal{B} = \mathcal{P}(\mathbb{N})$, and let μ be the counting measure. Let $E_n = \{m \in \mathbb{N} : m \geq n\}$. Then $\mu(E_n) = \infty$ for every $n \in \mathbb{N}$, but $\bigcap_{n \in \mathbb{N}} E_n = \emptyset$, so

$$\mu \left(\bigcap_{n \in \mathbb{N}} E_n \right) = 0 \neq \infty = \lim_{n \rightarrow \infty} \mu(E_n).$$

Chapter Notes

The content of this chapter is common to every text on abstract measure theory, though the order of presentation differs. We have elected to follow more or less the order of presentation from Rudin's *Real and Complex Analysis* [7, Chapter 1]. Alternative presentations can be found in [2, Sections 1.2, 1.3, and 2.1], and [10, Section 1.4].

Introductory texts on measure theory tend not to give much treatment to the Borel hierarchy or other topics in descriptive set theory (and we will also not expand on such topics within these lecture notes). Those interested in learning more can take a look at the book of Kechris [4] and/or the lecture notes of Tserunyan [11], which draw quite heavily on [4].

Exercises

2.1 Let X be a set. A family of subsets $\mathcal{S} \subseteq \mathcal{P}(X)$ is a *semi-algebra* if

- $\emptyset, X \in \mathcal{S}$;
- if $A, B \in \mathcal{S}$, then $A \cap B \in \mathcal{S}$;
- if $A, B \in \mathcal{S}$, then $A \setminus B = \bigsqcup_{i=1}^n C_i$ for some $C_1, \dots, C_n \in \mathcal{S}$.

Show that if \mathcal{S} is a semi-algebra, then the algebra generated by \mathcal{S} is

$$\mathcal{A}(\mathcal{S}) = \left\{ \bigcup_{i=1}^n A_i : n \in \mathbb{N}, A_i \in \mathcal{S} \right\}.$$

Can \bigcup be replaced by \bigsqcup ?

2.2 Suppose \mathcal{B} is an infinite σ -algebra (on an infinite set X).

- Show that \mathcal{B} contains an infinite sequence $(E_n)_{n \in \mathbb{N}}$ of pairwise disjoint sets.
- Deduce that \mathcal{B} has at least the cardinality of the continuum.

2.3 Prove that the following sets are Borel sets in \mathbb{R} :

- The set of points of continuity

$$C_f = \{x \in \mathbb{R} : f \text{ is continuous at } x\}$$

for an arbitrary function $f : \mathbb{R} \rightarrow \mathbb{R}$.

- The set of points of convergence

$$\text{Conv} = \{x \in \mathbb{R} : \lim_{n \rightarrow \infty} f_n(x) \text{ exists}\}$$

for an arbitrary sequence of continuous functions $f_n : \mathbb{R} \rightarrow \mathbb{R}$.

2.4 Let X, Y be sets and $f : X \rightarrow Y$ any function.

- Prove that if $\mathcal{C} \subseteq \mathcal{P}(Y)$ is a σ -algebra on Y , then $\mathcal{B} = \{f^{-1}(C) : C \in \mathcal{C}\}$ is a σ -algebra on X .
- Prove that for any family of sets $\mathcal{S} \subseteq \mathcal{P}(Y)$, we have $\sigma(f^{-1}(\mathcal{S})) = f^{-1}(\sigma(\mathcal{S}))$.

2.5 Prove that the extended real line $[-\infty, \infty]$ is homeomorphic to the closed unit interval $[0, 1]$.

2.6 Prove Proposition 2.10.

2.7 Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [0, \infty]$. Prove that μ is a measure if and only if it satisfies the following three properties:

- $\mu(\emptyset) = 0$;
- FINITE ADDITIVITY: for any disjoint sets $A, B \in \mathcal{B}$,

$$\mu(A \sqcup B) = \mu(A) + \mu(B);$$

- CONTINUITY FROM BELOW: if $E_1 \subseteq E_2 \subseteq \dots \in \mathcal{B}$, then

$$\mu\left(\bigcup_{n \in \mathbb{N}} E_n\right) = \lim_{n \rightarrow \infty} \mu(E_n).$$

2.8 Let (X, \mathcal{B}, μ) be a finite measure space, and let $\mathcal{A} \subseteq \mathcal{P}(X)$ be an algebra on X such that $\sigma(\mathcal{A}) = \mathcal{B}$. Show that for every $B \in \mathcal{B}$ and every $\varepsilon > 0$, there exists $A \in \mathcal{A}$ such that $\mu(A \Delta B) < \varepsilon$.

2.9 Prove the Borel–Cantelli lemma: if $(A_n)_{n \in \mathbb{N}}$ is a family of measurable subsets of a measure space (X, \mathcal{B}, μ) and $\sum_{n \in \mathbb{N}} \mu(A_n) < \infty$, then

$$\mu(\{x \in X : x \in A_n \text{ for infinitely many } n \in \mathbb{N}\}) = 0.$$

CHAPTER 3

Integration Against a Measure

Our next task is to develop an integration theory for integrating measurable functions on abstract measure spaces. In the Riemann–Darboux approach to integration, we approximate a function $f : [a, b] \rightarrow [0, \infty)$ by step functions, for which we can easily define the integral. For the Lebesgue theory of integration, we will use a similar idea but with a more general class of functions: so-called simple functions.

1. Integration of Simple Functions

DEFINITION 3.1

Let (X, \mathcal{B}) be a measurable space. A *simple function* is a measurable function $s : X \rightarrow \mathbb{C}$ taking only finitely many values.

Partitioning X into finitely many pieces corresponding to the values of a simple function s , we may write simple functions as linear combinations of indicator functions of measurable sets. That is, $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$ for some numbers $c_j \in \mathbb{C}$ and measurable sets $E_j \in \mathcal{B}$. Given a measure μ on (X, \mathcal{B}) , we define the integral of a simple function in the obvious way. To avoid issues with adding and subtracting infinities, we will deal for now only with nonnegative functions.

DEFINITION 3.2

Let (X, \mathcal{B}, μ) be a measure space and $s : X \rightarrow [0, \infty)$ a simple function. Write $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$ with $c_j \geq 0$ and $E_j \in \mathcal{B}$. The *integral of s with respect to μ* is given by

$$\int_X s \, d\mu = \sum_{j=1}^n c_j \mu(E_j).$$

PROPOSITION 3.3

The integral of a nonnegative simple function is well-defined. That is, the value of the integral of a simple function s does not depend on the representation of s as a linear combination of indicator functions of measurable sets.

PROOF. Suppose $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$. Let a_1, \dots, a_m be the finite collection of values taken by s , and let $A_k = \{s = a_k\}$ for $k = 1, \dots, m$. Then the sets A_1, \dots, A_k partition X , and $s = \sum_{k=1}^m a_k \mathbb{1}_{A_k}$. We will show $\sum_{j=1}^n c_j \mu(E_j) = \sum_{k=1}^m a_k \mu(A_k)$.

Define a new collection of sets $E'_J = \bigcap_{j \in J} E_j \setminus \bigcup_{i \notin J} E_j$ for $J \subseteq \{1, \dots, n\}$. In other words, $x \in E'_J$ means that $x \in E_j$ if and only if $j \in J$. This defines a partition of X . Note that the

value of s on the set E'_J is $c'_J = \sum_{j \in J} c_j$. We can therefore relate the sets E'_J to the sets A_k by

$$A_k = \bigsqcup_{J \subseteq \{1, \dots, n\}, c'_J = a_k} E'_J.$$

Then on the one hand,

$$\sum_{k=1}^m a_k \mu(A_k) = \sum_{k=1}^m a_k \sum_{J \subseteq \{1, \dots, n\}, c'_J = a_k} \mu(E'_J) = \sum_{J \subseteq \{1, \dots, n\}} c'_J \mu(E'_J).$$

On the other hand,

$$\sum_{j=1}^n c_j \mu(E_j) = \sum_{j=1}^n c_j \sum_{\{j\} \subseteq J \subseteq \{1, \dots, n\}} \mu(E'_J) = \sum_{J \subseteq \{1, \dots, n\}} \sum_{j \in J} c_j \mu(E'_J) = \sum_{J \subseteq \{1, \dots, n\}} c'_J \mu(E'_J).$$

This completes the proof. \square

We used a particular representation of a simple function in the previous proof that will continue to be convenient to work with. Say that $\sum_{j=1}^n c_j \mathbb{1}_{E_j}$ is the *standard representation* of a simple function s if $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$, and the sets E_1, \dots, E_n partition X (that is, they are pairwise disjoint and their union is X).

PROPOSITION 3.4

Let (X, \mathcal{B}, μ) be a measure space, let $s, t : X \rightarrow [0, \infty)$ be simple functions, and let $c \in \mathbb{R}$, $c \geq 0$. Then

- (1) $\int_X cs \, d\mu = c \cdot \int_X s \, d\mu$;
- (2) $\int_X (s + t) \, d\mu = \int_X s \, d\mu + \int_X t \, d\mu$;
- (3) if $s \leq t$, then $\int_X s \, d\mu \leq \int_X t \, d\mu$.

PROOF. (1) Let $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$. Then $cs = \sum_{j=1}^n (cc_j) \mathbb{1}_{E_j}$, so

$$\int_X cs \, d\mu = \sum_{j=1}^n (cc_j) \mu(E_j) = c \cdot \sum_{j=1}^n c_j \mu(E_j) = c \cdot \int_X s \, d\mu.$$

For (2) and (3), it will be helpful to work with the standard representation, so let $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$ and $t = \sum_{k=1}^m d_k \mathbb{1}_{F_k}$ be the standard representations. Define sets $A_{j,k} = E_j \cap F_k$ for $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$. Then $E_j = \bigsqcup_{k=1}^m A_{j,k}$ and $F_k = \bigsqcup_{j=1}^n A_{j,k}$.

(2) The function $s + t$ takes the value $c_j + d_k$ on $A_{j,k}$, so

$$\int_X (s + t) \, d\mu = \sum_{j,k} (c_j + d_k) \mu(A_{j,k}) = \sum_{j=1}^n c_j \underbrace{\sum_{k=1}^m \mu(A_{j,k})}_{\mu(E_j)} + \sum_{k=1}^m d_k \underbrace{\sum_{j=1}^n \mu(A_{j,k})}_{\mu(F_k)} = \int_X s \, d\mu + \int_X t \, d\mu.$$

(3) By assumption, if $A_{j,k} \neq \emptyset$, then $c_j \leq d_k$. Thus,

$$\int_X s \, d\mu = \sum_{j=1}^n c_j \mu(E_j) = \sum_{j,k} c_j \mu(A_{j,k}) \leq \sum_{j,k} d_k \mu(A_{j,k}) = \sum_{k=1}^m d_k \mu(F_k) = \int_X t \, d\mu.$$

□

DEFINITION 3.5

Let (X, \mathcal{B}, μ) be a measure space, $s : X \rightarrow [0, \infty)$ a simple function, and $E \in \mathcal{B}$ a measurable set. The *integral of s with respect to μ over E* is given by

$$\int_E s \, d\mu = \int_X s \cdot \mathbb{1}_E \, d\mu.$$

Note that if s is simple, then $s \cdot \mathbb{1}_E$ is also simple, so the above definition makes sense.

PROPOSITION 3.6

Let (X, \mathcal{B}, μ) be a measure space, and let $s : X \rightarrow [0, \infty)$ be a simple function. Then

$$\nu(E) = \int_E s \, d\mu$$

defines a measure on (X, \mathcal{B}) .

PROOF. Note that $s \cdot \mathbb{1}_\emptyset = 0$, so $\nu(\emptyset) = 0$. Suppose $(E_n)_{n \in \mathbb{N}}$ is a pairwise disjoint family of measurable sets, and let $E = \bigsqcup_{n \in \mathbb{N}} E_n$. Write $s = \sum_{j=1}^m a_j \mathbb{1}_{A_j}$. Then $s \cdot \mathbb{1}_E = \sum_{j=1}^m a_j \mathbb{1}_{A_j \cap E}$, so

$$\nu(E) = \sum_{j=1}^m a_j \mu(A_j \cap E) = \sum_{j,n} a_j \mu(A_j \cap E_n) = \sum_{n \in \mathbb{N}} \int_X s \cdot \mathbb{1}_{E_n} \, d\mu = \sum_{n \in \mathbb{N}} \nu(E_n).$$

Note that the sum over n is an infinite sum so reordering requires some justification. Fortunately, all of the values $a_j \mu(A_j \cap E_n)$ are nonnegative, so the sum can be computed in any order without changing the value. □

2. Integration of Nonnegative Measurable Functions

We now want to extend the definition of the integral against a measure to all nonnegative measurable functions. The next proposition shows that simple functions are a sufficiently general class to approximate arbitrary measurable functions.

PROPOSITION 3.7

Let (X, \mathcal{B}) be a measurable space, and let $f : X \rightarrow [0, \infty]$ be measurable. Then there exists a sequence $(s_n)_{n \in \mathbb{N}}$ of simple functions such that $0 \leq s_1 \leq s_2 \leq \dots \leq f$, and $s_n \rightarrow f$ pointwise.

PROOF. For $n \in \mathbb{N}$, define

$$s_n(x) = \begin{cases} \frac{a}{2^n}, & \text{if } \frac{a}{2^n} \leq f(x) < \frac{a+1}{2^n} \text{ and } a < n \cdot 2^n. \\ n, & \text{if } f(x) \geq n. \end{cases}$$

□

It is therefore reasonable to define the integral of an arbitrary nonnegative measurable function as follows.

DEFINITION 3.8

Let (X, \mathcal{B}, μ) be a measure space, and let $f : X \rightarrow [0, \infty]$ be measurable. We define the *integral of f with respect to μ* as

$$\int_X f \, d\mu = \sup \left\{ \int_X s \, d\mu : s \text{ simple and } 0 \leq s \leq f \right\}.$$

Given a measurable set $E \in \mathcal{B}$, the *integral of f with respect to μ over E* is defined by

$$\int_E f \, d\mu = \int_X f \cdot \mathbb{1}_E \, d\mu.$$

One may object at this point and suggest an alternative definition. Since $f : X \rightarrow [0, \infty]$ can be obtained as $f = \lim_{n \rightarrow \infty} s_n$ for an increasing sequence of simple functions $0 \leq s_1 \leq s_2 \leq \dots$, why not define $\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X s_n \, d\mu$? As we will see shortly, this is in fact an equivalent definition that is extremely useful for many applications. However, *as a definition*, it has two serious defects: why should the limit exist? and why should the value be the same for all possible approximations by simple functions? This is why we prefer Definition 3.8 above (and why this is the standard definition across measure theory textbooks).

PROPOSITION 3.9

Let (X, \mathcal{B}, μ) be a measure space, and let $f, g : X \rightarrow [0, \infty]$ be measurable. If $f \leq g$, then

$$\int_X f \, d\mu \leq \int_X g \, d\mu.$$

PROOF. It suffices to observe $\{s \text{ simple function} : 0 \leq s \leq f\} \subseteq \{s \text{ simple function} : 0 \leq s \leq g\}$. \square

THEOREM 3.10: MONOTONE CONVERGENCE THEOREM

Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions $0 \leq f_1 \leq f_2 \leq \dots$, and let $f = \lim_{n \rightarrow \infty} f_n$. Then

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

REMARK. Note that a consequence of the monotone convergence theorem is that $\int_X f \, d\mu$ can be computed by taking a sequence of simple functions $0 \leq s_1 \leq s_2 \leq \dots \rightarrow f$ and computing $\lim_{n \rightarrow \infty} \int_X s_n \, d\mu$.

PROOF OF MONOTONE CONVERGENCE THEOREM. First, f is a measurable function by Proposition 2.11. By monotonicity of the integral (Proposition 3.9), the sequence $\int_X f_n \, d\mu$ is increasing, so $\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \sup_{n \in \mathbb{N}} \int_X f_n \, d\mu \in [0, \infty]$ exists as an extended real number. Moreover,

$$\int_X f \, d\mu \geq \lim_{n \rightarrow \infty} \int_X f_n \, d\mu,$$

since the inequality holds for each $n \in \mathbb{N}$. Therefore, it suffices to show

$$\int_X f \, d\mu \leq \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

If $\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \infty$, there is nothing to prove, so assume $\lim_{n \rightarrow \infty} \int_X f_n \, d\mu < \infty$.

Let $c < 1$. Let $s : X \rightarrow [0, \infty)$ be a simple function, $0 \leq s \leq f$. For $n \in \mathbb{N}$, let $E_n = \{f_n \geq cs\}$. Then $E_1 \subseteq E_2 \subseteq \dots$ and $X = \bigcup_{n \in \mathbb{N}} E_n$. By Proposition 3.6, let $\nu : \mathcal{B} \rightarrow [0, \infty]$ be the measure $\nu(E) = \int_E s \, d\mu$. We have

$$\begin{aligned} c \cdot \int_X s \, d\mu &= c \cdot \nu(X) \\ &= c \cdot \lim_{n \rightarrow \infty} \nu(E_n) && \text{(continuity from below)} \\ &= \lim_{n \rightarrow \infty} c \cdot \nu(E_n) && \text{(Proposition 2.10)} \\ &= \lim_{n \rightarrow \infty} \int_{E_n} cs \, d\mu && \text{(Proposition 3.4)} \\ &\leq \lim_{n \rightarrow \infty} \int_X f_n \, d\mu && \text{(monotonicity).} \end{aligned}$$

Taking a supremum over all such simple functions, we conclude

$$c \cdot \int_X f \, d\mu \leq \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

Letting $c \rightarrow 1$ yields the desired result. □

PROPOSITION 3.11

Let (X, \mathcal{B}, μ) be a measure space, and let $f, g : X \rightarrow [0, \infty]$ be measurable functions. Let $c \in [0, \infty)$.

- (1) $\int_X cf \, d\mu = c \cdot \int_X f \, d\mu$.
- (2) $\int_X (f + g) \, d\mu = \int_X f \, d\mu + \int_X g \, d\mu$.

PROOF. (1) This follows quickly from the definition of the integral and Proposition 3.4.

(2) We use the monotone convergence theorem. Let $0 \leq s_1 \leq s_n \leq \dots \leq f$ with $s_n \rightarrow f$ and $0 \leq t_1 \leq t_2 \leq \dots \leq g$ with $t_n \rightarrow g$. Then $0 \leq s_1 + t_1 \leq s_2 + t_2 \leq \dots \leq f + g$ and $s_n + t_n \rightarrow f + g$. Thus,

$$\begin{aligned} \int_X (f + g) \, d\mu &= \lim_{n \rightarrow \infty} \int_X (s_n + t_n) \, d\mu && \text{(MCT)} \\ &= \lim_{n \rightarrow \infty} \int_X s_n \, d\mu + \lim_{n \rightarrow \infty} \int_X t_n \, d\mu && \text{(Proposition 3.4)} \\ &= \int_X f \, d\mu + \int_X g \, d\mu && \text{(MCT).} \end{aligned}$$

□

THEOREM 3.12

Let (X, \mathcal{B}, μ) be a measure space, and let $(f_n)_{n \in \mathbb{N}}$ be a sequence of nonnegative measurable functions, $f_n : X \rightarrow [0, \infty]$. Then

$$\int_X \left(\sum_{n=1}^{\infty} f_n \right) \, d\mu = \sum_{n=1}^{\infty} \int_X f_n \, d\mu.$$

PROOF. We have

$$\begin{aligned}
\int_X \left(\sum_{n=1}^{\infty} f_n \right) d\mu &= \int_X \lim_{N \rightarrow \infty} \left(\sum_{n=1}^N f_n \right) d\mu \\
&= \lim_{N \rightarrow \infty} \int_X \left(\sum_{n=1}^N f_n \right) d\mu && \text{(MCT)} \\
&= \lim_{N \rightarrow \infty} \sum_{n=1}^N \int_X f_n d\mu && \text{(additivity of the integral)} \\
&= \sum_{n=1}^{\infty} \int_X f_n d\mu.
\end{aligned}$$

□

THEOREM 3.13: FATOU'S LEMMA

Let (X, \mathcal{B}, μ) be a measure space. Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions, $f_n : X \rightarrow [0, \infty]$. Then

$$\int_X \liminf_{n \rightarrow \infty} f_n d\mu \leq \liminf_{n \rightarrow \infty} \int_X f_n d\mu.$$

PROOF. Let $f = \liminf_{n \rightarrow \infty} f_n$. Define $F_N = \inf_{n \geq N} f_n$. Then $0 \leq F_1 \leq F_2 \leq \dots$ and $F_N \rightarrow f$. Therefore,

$$\begin{aligned}
\int_X f d\mu &= \lim_{N \rightarrow \infty} \int_X F_N d\mu && \text{(MCT)} \\
&\leq \lim_{N \rightarrow \infty} \inf_{n \geq N} \int_X f_n d\mu && \text{(monotonicity of the integral)} \\
&= \liminf_{N \rightarrow \infty} \int_X f_n d\mu.
\end{aligned}$$

□

3. Integration of Real and Complex-Valued Functions

The method for integrating real and complex-valued functions involves decomposing these functions as linear combinations of nonnegative functions. An important observation is that such a decomposition can be done in a measurable way.

DEFINITION 3.14

Let X be a set and $f : X \rightarrow [-\infty, \infty]$. The *positive part* f^+ and *negative part* f^- of f are defined by

$$f^+ = \max\{f, 0\} \quad \text{and} \quad f^- = \max\{-f, 0\}.$$

Note that $f = f^+ - f^-$ and $|f| = f^+ + f^-$. Moreover, if (X, \mathcal{B}) is a measurable space and $f : X \rightarrow [-\infty, \infty]$ is measurable, then f^+ and f^- are measurable by Proposition 2.11.

DEFINITION 3.15

Let (X, \mathcal{B}, μ) be a measure space.

- An extended real-valued measurable function $f : X \rightarrow [-\infty, \infty]$ is *integrable* if

$$\int_X |f| d\mu < \infty.$$

In this case, the *integral of f* is defined by

$$\int_X f d\mu = \int_X f^+ d\mu - \int_X f^- d\mu.$$

- A complex-valued measurable function $f : X \rightarrow \mathbb{C}$ is *integrable* if

$$\int_X |f| d\mu < \infty,$$

and the *integral of f* is defined by

$$\int_X f d\mu = \int_X \operatorname{Re}(f) d\mu + i \int_X \operatorname{Im}(f) d\mu.$$

- Given a measurable set $E \in \mathcal{B}$, a measurable function f taking extended real or complex values is *integrable over E* if $f \cdot \mathbb{1}_E$ is integrable, and the *integral of f over E* is

$$\int_E f d\mu = \int_X f \cdot \mathbb{1}_E d\mu.$$

REMARK. By monotonicity of the integral (Proposition 3.9), if a function is integrable, then it is also integrable over every measurable subset of X .

4. Integral Identities and Inequalities

PROPOSITION 3.16: TRIANGLE INEQUALITY FOR THE INTEGRAL

Suppose (X, \mathcal{B}, μ) is a measure space and $f : X \rightarrow \mathbb{C}$ is an integrable function. Then

$$\left| \int_X f d\mu \right| \leq \int_X |f| d\mu.$$

PROOF. First, suppose f is real-valued. Then by the triangle inequality and linearity,

$$\left| \int_X f d\mu \right| = \left| \int_X f^+ d\mu - \int_X f^- d\mu \right| \leq \int_X f^+ d\mu + \int_X f^- d\mu = \int_X |f| d\mu.$$

Now suppose f is complex-valued. Let $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ such that $|\int_X f d\mu| = \lambda \int_X f d\mu$. Then

$$\left| \int_X f d\mu \right| = \operatorname{Re} \left(\int_X \lambda f d\mu \right) = \int_X \operatorname{Re}(\lambda f) d\mu \leq \int_X |\operatorname{Re}(\lambda f)| d\mu \leq \int_X |f| d\mu.$$

□

PROPOSITION 3.17: LINEARITY OF THE INTEGRAL

Let (X, \mathcal{B}, μ) be a measure space. Let $f, g : X \rightarrow \mathbb{C}$ be integrable functions, and let $c \in \mathbb{C}$. Then

- (1) $f + g$ is integrable, and $\int_X (f + g) d\mu = \int_X f d\mu + \int_X g d\mu$.

(2) cf is integrable, and $\int_X cf \, d\mu = c \int_X f \, d\mu$.

PROOF. (1) First, by the triangle inequality, we have $|f + g| \leq |f| + |g|$. Therefore,

$$\int_X |f + g| \, d\mu \stackrel{(*)}{\leq} \int_X (|f| + |g|) \, d\mu \stackrel{(**)}{=} \int_X |f| \, d\mu + \int_X |g| \, d\mu < \infty.$$

In step (*), we have used monotonicity of the integral (Proposition 3.9), and in (**), we have used additivity (Proposition 3.11).

Decomposing f and g into their real and imaginary parts, it suffices to prove the identity $\int_X (f + g) \, d\mu = \int_X f \, d\mu + \int_X g \, d\mu$ for real-valued functions f and g . Let $h = f + g$. Then $h = h^+ - h^- = f^+ - f^- + g^+ - g^-$. This can be rearranged to the identity $h^+ + f^- + g^- = h^- + f^+ + g^+$. Then using additivity of the integral for nonnegative functions (Proposition 3.11), we have

$$\begin{aligned} \int_X h^+ \, d\mu + \int_X f^- \, d\mu + \int_X g^- \, d\mu &= \int_X (h^+ + f^- + g^-) \, d\mu \\ &= \int_X (h^- + f^+ + g^+) \, d\mu \\ &= \int_X h^- \, d\mu + \int_X f^+ \, d\mu + \int_X g^+ \, d\mu. \end{aligned} \tag{3.1}$$

Rearranging again,

$$\begin{aligned} \int_X (f + g) \, d\mu &= \int_X h^+ \, d\mu - \int_X h^- \, d\mu && \text{(Definition 3.15)} \\ &= \int_X f^+ \, d\mu - \int_X f^- \, d\mu + \int_X g^+ \, d\mu - \int_X g^- \, d\mu && \text{(by (3.1))} \\ &= \int_X f \, d\mu + \int_X g \, d\mu && \text{(Definition 3.15)} \end{aligned}$$

(2) Note that $|cf| = |c||f|$, so

$$\int_X |cf| \, d\mu = \int_X |c||f| \, d\mu \stackrel{(*)}{=} |c| \int_X |f| \, d\mu < \infty,$$

where (*) follows from Proposition 3.11. Hence, cf is integrable.

For computing the integral of cf , we consider several different cases.

CASE 1. $c \geq 0$

When f is nonnegative, we have

$$\int_X cf \, d\mu = c \int_X f \, d\mu$$

by Proposition 3.11. The identity follows for a general complex-valued function f by decomposing $f = (\operatorname{Re}(f)^+ - \operatorname{Re}(f)^-) + i(\operatorname{Im}(f)^+ - \operatorname{Im}(f)^-)$.

CASE 2. $c = -1$

For real-valued $f : X \rightarrow \mathbb{R}$, we use the identities $(-f)^+ = f^-$ and $(-f)^- = f^+$ to obtain

$$\int_X (-f) \, d\mu = \int_X f^- \, d\mu - \int_X f^+ \, d\mu = - \int_X f \, d\mu.$$

Complex-valued functions can be handled by decomposing into real and imaginary parts.

CASE 3. $c = i$

Noting that $\operatorname{Re}(if) = -\operatorname{Im}(f)$ and $\operatorname{Im}(if) = \operatorname{Re}(f)$, we have

$$\begin{aligned} \int_X if \, d\mu &= \int_X (-\operatorname{Im}(f)) \, d\mu + i \int_X \operatorname{Re}(f) \, d\mu && \text{(Definition 3.15)} \\ &= - \int_X \operatorname{Im}(f) \, d\mu + i \int_X \operatorname{Re}(f) \, d\mu && \text{(Case 2)} \\ &= i \left(\int_X \operatorname{Re}(f) \, d\mu + i \int_X \operatorname{Im}(f) \, d\mu \right) \\ &= i \int_X f \, d\mu && \text{(Definition 3.15)} \end{aligned}$$

CASE 4. $c \in \mathbb{R}$

Combine Case 1 and Case 2.

CASE 5. $c \in \mathbb{C}$

Write $c = a + ib$ with $a, b \in \mathbb{R}$. Then

$$\begin{aligned} \int_X cf \, d\mu &= \int_X (af + ibf) \, d\mu \\ &= \int_X af \, d\mu + \int_X ibf \, d\mu && \text{(by (1))} \\ &= \int_X af \, d\mu + i \int_X bf \, d\mu && \text{(Case 3)} \\ &= a \int_X f \, d\mu + ib \int_X f \, d\mu && \text{(Case 4)} \\ &= c \int_X f \, d\mu. \end{aligned}$$

□

Let (X, \mathcal{B}, μ) be a measure space, and denote by $L^1(\mu)$ the set of integrable functions. Proposition 3.17 shows that $L^1(\mu)$ is a (complex) vector space. Moreover, in the course of the proof, we showed

$$\int_X |cf| \, d\mu = |c| \int_X |f| \, d\mu \quad \text{and} \quad \int_X |f + g| \, d\mu \leq \int_X |f| \, d\mu + \int_X |g| \, d\mu.$$

In other words, if we let

$$\|f\|_1 = \int_X |f| \, d\mu,$$

then $\|\cdot\|_1$ defines a *seminorm* on the vector space of integrable functions on (X, \mathcal{B}, μ) .

DEFINITION 3.18

Let V be a real or complex vector space. A function $\|\cdot\| : V \rightarrow [0, \infty)$ is a *seminorm* if it satisfies:

- TRIANGLE INEQUALITY: $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in V$, and
- ABSOLUTE HOMOGENEITY: $\|cv\| = |c| \|v\|$ for all $v \in V$ and all scalars c .

A seminorm is a *norm* if it satisfies the additional property

- POSITIVE DEFINITE: if $v \in V$ and $\|v\| = 0$, then $v = 0$.

The seminorm $\|\cdot\|_1$ on the space of integrable functions may not be a norm in general, but a small modification will turn it into a norm. This will be discussed in greater detail later in the course, in the context of so-called L^p spaces. One of the important ingredients is a deeper understanding of *null sets*, which we will discuss now.

5. Sets of Measure Zero

DEFINITION 3.19

Let (X, \mathcal{B}, μ) be a measure space.

- A measurable set $N \in \mathcal{B}$ is a *null set* if $\mu(N) = 0$.
- We say that a property holds *almost everywhere* if there exists a null set $N \in \mathcal{B}$ such that the property holds for every point $x \in X \setminus N$.

REMARK. An easy consequence of countable additivity and monotonicity of measures is that the family \mathcal{N} of null sets forms a σ -ideal of \mathcal{B} :

- $\emptyset \in \mathcal{N}$;
- if $A \in \mathcal{N}$ and $B \in \mathcal{B}$ with $B \subseteq A$, then $B \in \mathcal{N}$; and
- if $(N_n)_{n \in \mathbb{N}}$ is a countable family of null sets, then $\bigcup_{n \in \mathbb{N}} N_n \in \mathcal{N}$.

NOTATION. The phrases “almost everywhere” or “almost every” are often abbreviated by a.e. or μ -a.e. if the measure needs to be specified. In a statement of the form “Property P holds a.e.,” we interpret a.e. as “almost everywhere.” For a statement of the form “Property P holds for a.e. $x \in X$,” we read a.e. as “almost every,” and the meaning is the same as in the previous example statement.

Null sets naturally arise and play an important role in integration theory. Some examples are provided by the next three propositions.

PROPOSITION 3.20

Let (X, \mathcal{B}, μ) be a measure space. Suppose $f : X \rightarrow [-\infty, \infty]$ is an integrable function. Then $f(x) \in \mathbb{R}$ for μ -a.e. $x \in X$.

PROOF. Let $N = \{x \in \mathbb{R} : |f(x)| = \infty\}$. We want to show that N is a null set. By monotonicity of the integral (Proposition 3.9),

$$\int_X |f| d\mu \geq \int_N |f| d\mu = \infty \cdot \mu(N).$$

On the other hand, by integrability of f ,

$$\int_X |f| d\mu < \infty.$$

Thus, $\infty \cdot \mu(N) < \infty$, so $\mu(N) = 0$. □

COROLLARY 3.21: BOREL–CANTELLI LEMMA

Let (X, \mathcal{B}, μ) be a measure space. Suppose $(E_n)_{n \in \mathbb{N}}$ is a sequence of measurable sets and $\sum_{n=1}^{\infty} \mu(E_n) < \infty$. Then

$$\mu(\{x \in X : x \in E_n \text{ for infinitely many } n \in \mathbb{N}\}) = 0.$$

PROOF. One possible proof uses continuity from above and was given in the exercises (see Exercise 2.9). We will now give a different proof using integration.

Let $f = \sum_{n=1}^{\infty} \mathbb{1}_{E_n}$. Note that $f(x) = \infty$ if and only if $x \in E_n$ for infinitely many $n \in \mathbb{N}$. By Theorem 3.12,

$$\int_X f d\mu = \sum_{n=1}^{\infty} \underbrace{\int_X \mathbb{1}_{E_n}}_{\mu(E_n)} < \infty.$$

So by Proposition 3.20, $f < \infty$ a.e. That is,

$$\mu(\{x \in X : x \in E_n \text{ for infinitely many } n \in \mathbb{N}\}) = \mu(\{f = \infty\}) = 0.$$

□

PROPOSITION 3.22

Let (X, \mathcal{B}, μ) be a measure space, and let $f, g : X \rightarrow \mathbb{C}$ be measurable functions. Suppose $f = g$ a.e. Then f is integrable if and only if g is integrable. Moreover, if f and g are integrable, then

$$\int_X f d\mu = \int_X g d\mu.$$

PROOF. Let $N = \{x \in X : f(x) \neq g(x)\}$. By assumption, N is a null set.

STEP 1. Integrability

Suppose f is integrable. Then

$$\begin{aligned} \int_X |g| d\mu &= \int_{X \setminus N} |f| d\mu + \int_N |g| d\mu && \text{(linearity of the integral)} \\ &\leq \int_X |f| d\mu + \underbrace{\infty \cdot \mu(N)}_0 && \text{(monotonicity of the integral)} \\ &= \int_X |f| d\mu < \infty, \end{aligned}$$

so g is integrable. Reversing the roles of f and g proves the converse.

STEP 2. Integral

Assume f and g are integrable. Then

$$\begin{aligned}
 \left| \int_X g \, d\mu - \int_X f \, d\mu \right| &= \left| \int_X (g - f) \, d\mu \right| && \text{(linearity of the integral)} \\
 &\leq \int_X |g - f| \, d\mu && \text{(triangle inequality for the integral)} \\
 &= \int_{X \setminus N} 0 \, d\mu + \int_N |g - f| \, d\mu && \text{(linearity of the integral)} \\
 &\leq 0 \cdot \mu(X \setminus N) + \infty \cdot \mu(N) = 0.
 \end{aligned}$$

□

PROPOSITION 3.23

Let (X, \mathcal{B}, μ) be a measure space, and let $f : X \rightarrow [0, \infty]$ be a measurable function. Then $\int_X f \, d\mu = 0$ if and only if $f = 0$ a.e.

PROOF. If $f = 0$ a.e., then by Proposition 3.22, f is integrable and

$$\int_X f \, d\mu = \int_X 0 \, d\mu = 0 \cdot \mu(X) = 0.$$

Conversely, suppose $\int_X f \, d\mu = 0$. Then by Markov's inequality (Exercise 3.2),

$$\mu(\{f > c\}) \leq \frac{1}{c} \int_X f \, d\mu = 0$$

for every $c > 0$. Therefore, by continuity of μ from below,

$$\mu(\{f \neq 0\}) = \mu\left(\bigcup_{n \in \mathbb{N}} \left\{f > \frac{1}{n}\right\}\right) = \lim_{n \rightarrow \infty} \mu\left(\left\{f > \frac{1}{n}\right\}\right) = 0.$$

That is, $f = 0$ a.e. □

The examples above (especially Proposition 3.22) show that null sets are negligible from the point of view of integration, and we can very often ignore modifications that happen on null sets. There is one subtle issue that requires care, however: in general, a subset of a null set may not be measurable and non-measurable modifications on null sets may create issues. For this reason, it is often convenient to work with *complete* measure spaces, as defined below.

DEFINITION 3.24

A measure space (X, \mathcal{B}, μ) is *complete* if every subset of every null set is measurable. That is, if $E \subseteq X$ and there exists $N \in \mathcal{B}$ with $E \subseteq N$ and $\mu(N) = 0$, then $E \in \mathcal{B}$.

The following proposition is a useful tool for passing to complete measure spaces.

PROPOSITION 3.25

Let (X, \mathcal{B}, μ) be a measure space. Let $\mathcal{N} = \{N \in \mathcal{B} : \mu(N) = 0\}$ be the σ -ideal of μ -null sets. Then the family $\bar{\mathcal{B}} = \{E \cup F : E \in \mathcal{B}, F \subseteq N \in \mathcal{N}\}$ is a σ -algebra, and there is a unique extension $\bar{\mu}$ of μ to $\bar{\mathcal{B}}$.

PROOF. Exercise 3.7. □

DEFINITION 3.26

The *completion* of a measure space (X, \mathcal{B}, μ) is the space $(X, \bar{\mathcal{B}}, \bar{\mu})$, where $\bar{\mathcal{B}}$ and $\bar{\mu}$ are as defined in Proposition 3.25.

6. The Dominated Convergence Theorem

We have already seen two fundamental convergence theorems for integration against a measure: the monotone convergence theorem and Fatou's lemma. We are nearly ready to state another fundamental result about integration: the dominated convergence theorem. First, we need to introduce the two notions of convergence that will be related by the dominated convergence theorem.

DEFINITION 3.27

Let (X, \mathcal{B}, μ) be a measure space.

- We say that a sequence $(f_n)_{n \in \mathbb{N}}$ of functions on X *converges almost everywhere* to a function f if $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for almost every $x \in X$.
- A sequence $(f_n)_{n \in \mathbb{N}}$ of integrable functions *converges in L^1* to $f \in L^1(\mu)$ if

$$\|f_n - f\|_1 = \int_X |f_n - f| \, d\mu \rightarrow 0$$

in \mathbb{R} as $n \rightarrow \infty$.

The dominated convergence theorem says that any sequence that converges almost everywhere and is “ L^1 -dominated” will converge in L^1 . The precise mathematical formulation is as follows:

THEOREM 3.28: DOMINATED CONVERGENCE THEOREM

Let (X, \mathcal{B}, μ) be a measure space. Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of integrable functions, $f_n : X \rightarrow \mathbb{C}$, and let $f : X \rightarrow \mathbb{C}$ be measurable. Suppose

- $f_n \rightarrow f$ a.e., and
- there is an integrable function $g : X \rightarrow [0, \infty)$ such that $\sup_{n \in \mathbb{N}} |f_n| \leq g$ a.e.

Then f is integrable and $f_n \rightarrow f$ in $L^1(\mu)$. In particular,

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

PROOF. First, $|f| \leq |g|$ a.e., so f is integrable.

Observe:

$$\begin{aligned} \int_X 2g \, d\mu - \limsup_{n \rightarrow \infty} \int_X |f - f_n| \, d\mu &= \liminf_{n \rightarrow \infty} \int_X (2g - |f - f_n|) \, d\mu \\ &\geq \int_X \liminf_{n \rightarrow \infty} (2g - |f - f_n|) \, d\mu && \text{(Fatou's lemma)} \\ &= \int_X 2g \, d\mu && (f_n \rightarrow f) \end{aligned}$$

Rearranging, we conclude

$$\limsup_{n \rightarrow \infty} \int_X |f - f_n| \, d\mu \leq 0.$$

Using the triangle inequality for the integral,

$$\left| \int_X f \, d\mu - \int_X f_n \, d\mu \right| \leq \int_X |f - f_n| \, d\mu \rightarrow 0,$$

so

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

□

The assumption that the sequence $(f_n)_{n \in \mathbb{N}}$ is “dominated” by an integrable function g is a necessary assumption to avoid “escape of mass to infinity,” as the following example demonstrates.

EXAMPLE 3.29

Let $X = \mathbb{Z}$, $\mathcal{B} = \mathcal{P}(\mathbb{Z})$, and let μ be the counting measure. Let $f_n = \mathbb{1}_{\{n\}}$. Then $f_n(x) \rightarrow 0$ for every $x \in X$. However,

$$\int_X f_n \, d\mu = 1$$

for every $n \in \mathbb{N}$, while

$$\int_X \lim_{n \rightarrow \infty} f_n \, d\mu = \int_X 0 \, d\mu = 0 \neq 1.$$

Chapter Notes

For other presentations of integration on abstract measures spaces, see [2, Section 2.1–2.3], [7, Chapter 1], [9, Sections 2.1 and 6.2], and/or [10, Section 1.3 and Subsection 1.4.4]. The development of integration in the books of Folland [2] and Rudin [7] is very similar to the presentation in these notes. By contrast, Stein and Shakarchi [9] and Tao [10] first develop integration in the special case of the Lebesgue measure before moving to abstract spaces. The book of Stein and Shakarchi [9] also proves the fundamental convergence theorems in a different order, starting with a special case of the dominated convergence theorem known as the *bounded convergence theorem*, and then deducing Fatou’s lemma, the monotone convergence theorem, and the general case of the dominated convergence theorem.

There is a very nice book of Oxtoby [6] that develops useful analogies between measure spaces and topological spaces and includes a discussion of null sets in relation to a σ -ideal of “topologically negligible” sets called *meager* sets or sets of *first category*.

Exercises

3.1 Let (X, \mathcal{B}, μ) be a measure space. Suppose $f : X \rightarrow [0, \infty]$ is a measurable function. Define $\nu : \mathcal{B} \rightarrow [0, \infty]$ by

$$\nu(E) = \int_E f \, d\mu.$$

Prove that ν is a measure.

3.2 Let (X, \mathcal{B}, μ) be a measure space and f a measurable function. Prove Markov’s inequality: for any $c > 0$,

$$\mu(\{|f| \geq c\}) \leq \frac{1}{c} \int_{\{|f| > c\}} |f| \, d\mu \leq \frac{1}{c} \int_X |f| \, d\mu.$$

3.3 Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be measure space, and let $T : X \rightarrow Y$ be a measurable function. Define $T\mu : \mathcal{C} \rightarrow [0, \infty]$ by $(T\mu)(A) = \mu(T^{-1}(A))$. Prove $T\mu = \nu$ if and only if for every integrable

function $f : Y \rightarrow \mathbb{C}$,

$$\int_Y f \, d\nu = \int_X f \circ T \, d\mu.$$

3.4 Let (X, \mathcal{B}, μ) be a probability space. Let $(A_n)_{n \in \mathbb{N}}$ be a family of measurable sets with $a = \inf_{n \in \mathbb{N}} \mu(A_n) > 0$. Show that there is a set $E \subseteq \mathbb{N}$ such that $\bar{d}(E) := \limsup_{N \rightarrow \infty} \frac{|E \cap \{1, \dots, N\}|}{N} \geq a$, and for any finite set $F \subseteq E$, $F \neq \emptyset$, one has $\mu(\bigcap_{n \in F} A_n) > 0$ by proving the following intermediate steps:

(a) Justify that we can assume without loss of generality that $\bigcap_{n \in F} A_n \neq \emptyset$ if and only if $\mu(\bigcap_{n \in F} A_n) > 0$ for every finite set $F \subseteq \mathbb{N}$. It may help to define the countable set

$$\mathcal{F} = \left\{ F \subseteq \mathbb{N} : |F| < \infty, \bigcap_{n \in F} A_n \neq \emptyset, \text{ and } \mu\left(\bigcap_{n \in F} A_n\right) = 0 \right\}.$$

(b) Prove

$$\int_X \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{A_n} \, d\mu \geq a.$$

(c) Define $E = \{n \in \mathbb{N} : x \in A_n\}$ for a suitable choice of $x \in X$.

3.5 Let (X, \mathcal{B}, μ) be a measure space, and let $f : X \rightarrow \mathbb{C}$ be an integrable function. Prove that for any $\varepsilon > 0$, there exists $\delta > 0$ with the following property: if $E \in \mathcal{B}$ and $\mu(E) < \delta$, then $|\int_E f \, d\mu| < \varepsilon$.

3.6 Let (X, \mathcal{B}, μ) be a measure space, and let $f : X \rightarrow \mathbb{C}$ be an integrable function. Show that $f = 0$ a.e. if and only if $\int_E f \, d\mu = 0$ for every $E \in \mathcal{B}$.

3.7 Prove Proposition 3.25.

3.8 Show that a measure space (X, \mathcal{B}, μ) is complete if and only if it satisfies the following property: for functions $f, g : X \rightarrow \mathbb{C}$, if f is measurable and $f = g$ a.e., then g is measurable.

3.9 Show that simple functions are dense in L^1 . That is, if (X, \mathcal{B}, μ) is a measure space and $f \in L^1(\mu)$, then for every $\varepsilon > 0$, there exists a simple function $s : X \rightarrow \mathbb{C}$ such that $\|f - s\|_1 < \varepsilon$.

3.10 Let (X, \mathcal{B}, μ) be a measure space and $E \in \mathcal{B}$. If $(E_n)_{n \in \mathbb{N}}$ is a sequence of measurable sets and $E = \bigcup_{n \in \mathbb{N}} E_n$, prove that for every integrable function $f \in L^1(\mu)$,

$$\lim_{n \rightarrow \infty} \int_{E_n} f \, d\mu = \int_E f \, d\mu.$$

State and prove an analogous result for decreasing sequences.

3.11 Prove

$$\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = \sum_{k \geq 0} \frac{x^k}{k!}.$$

3.12 Let (X, \mathcal{B}, μ) be a measure space, and let $(f_n)_{n \in \mathbb{N}}$ be a sequence of measurable functions $f_n : X \rightarrow \mathbb{C}$. Suppose

$$\sum_{n \in \mathbb{N}} \int_X |f_n| \, d\mu < \infty.$$

Prove that $\sum_{n \in \mathbb{N}} f_n$ converges a.e. to an integrable function $f \in L^1(\mu)$, and

$$\int_X f \, d\mu = \sum_{n \in \mathbb{N}} \int_X f_n \, d\mu.$$

3.13 In this exercise, we will use measure-theoretic tools in order to carry out computations with Riemann integrals. Assume for the purposes of this exercise that there is a measure λ on the Borel subsets of \mathbb{R} with the property: if $f : [a, b] \rightarrow \mathbb{R}$ is a Riemann integrable function, then

$$\int_{[a,b]} f \, d\lambda = \int_a^b f(x) \, dx,$$

where the integral on the left is the measure-theoretic integral and the integral on the right is the Riemann integral. (We will discuss multiple methods of constructing such a measure λ in future lectures.)

(a) Compute

$$\lim_{n \rightarrow \infty} \int_0^\infty \frac{n \sin(\frac{x}{n})}{x(1+x^2)} \, dx.$$

(b) Show that for $a > -1$,

$$\int_0^1 \frac{x^a \log x}{1-x} \, dx = - \sum_{k=1}^{\infty} \frac{1}{(a+k)^2}.$$


Part 2

Constructions of Measures

CHAPTER 4

Taxonomy of Measures

In this short chapter, we give a taxonomy of measures based on various properties that may be desirable or undesirable in certain circumstances. Some of these properties have been seen in previous chapters, while others are introduced for the first time here. The taxonomy is summarized in Figure 4.1, and precise definitions are given below.



1. Properties of Measures

DEFINITION 4.1

Let (X, \mathcal{B}, μ) be a measure space. A set E is

- *locally measurable* if $E \cap K \in \mathcal{B}$ for every set $K \in \mathcal{B}$ with $\mu(K) < \infty$;
- an *atom* if $\mu(E) > 0$ and every measurable subset $F \subseteq E$, $F \in \mathcal{B}$ satisfies either $\mu(F) = 0$ or $\mu(E \setminus F) = 0$.

DEFINITION 4.2

Let (X, \mathcal{B}) be a measurable space. A measure μ on (X, \mathcal{B}) is

- a *probability measure* if $\mu(X) = 1$;
- *finite* if $\mu(X) < \infty$;
- *σ -finite* if there is a countable sequence of measurable sets $(X_n)_{n \in \mathbb{N}}$ in \mathcal{B} such that $X = \bigcup_{n \in \mathbb{N}} X_n$ and $\mu(X_n) < \infty$ for each $n \in \mathbb{N}$;
- *s-finite* if μ is a countable sum $\mu = \sum_{n \in \mathbb{N}} \mu_n$ of finite measures $\mu_n : \mathcal{B} \rightarrow [0, \infty)$;
- *semi-finite* if every set of infinite measure contains a subset of positive finite measure, i.e. if $E \in \mathcal{B}$ and $\mu(E) = \infty$, then there exists $F \in \mathcal{B}$ with $F \subseteq E$ and $\mu(F) < \infty$;
- *complete* if every subset of every null set is measurable, i.e. if $E \subseteq X$ and there exists $N \in \mathcal{B}$ with $E \subseteq N$ and $\mu(N) = 0$, then $E \in \mathcal{B}$;
- *saturated* if every locally measurable set is measurable;
- *discrete* if μ is a combination of Dirac measures, $\mu = \sum_{x \in X} c_x \delta_x$ for some coefficients $c_x \in [0, \infty]$;
- *continuous* if μ has no point masses, i.e. $\mu(\{x\}) = 0$ for every $x \in X$;
- *(purely) atomic* if every set of positive measure contains an atom;
- *non-atomic* or *diffuse* if there are no atoms.

If X is a topological space and $\mathcal{B} = \text{Borel}(X)$, then μ is

- *locally finite* if every compact set has finite measure;
- *outer regular* if $\mu(E) = \inf\{\mu(U) : U \supseteq E \text{ open}\}$ for every $E \in \mathcal{B}$;
- *inner regular* if $\mu(E) = \sup\{\mu(K) : K \subseteq E \text{ compact}\}$ for every $E \in \mathcal{B}$.

The relationships between the various properties in Definition 4.2 are displayed in Figure 4.1.

2. Finiteness Properties

As shown in Figure 4.1, every probability measure is finite, every finite measure is σ -finite, and every σ -finite measure is both s-finite and semi-finite. The next example shows that s-finiteness and semi-finiteness are rather different notions from one another, neither one implying the other.

EXAMPLE 4.3

AN S-FINITE MEASURE THAT IS NOT SEMI-FINITE: Let X be a non-empty set, and let $x \in X$. Define $\mu(E) = \infty$ if $x \in E$ and $\mu(E) = 0$ if $x \notin E$. Then $\mu = \sum_{n=1}^{\infty} \delta_x$, so μ is s-finite. However, the set $\{x\}$ has infinite measure and no subsets of non-zero finite measure, so μ is not semi-finite.

A SEMI-FINITE MEASURE THAT IS NOT S-FINITE: Let X be an uncountable set, and let $\mu : \mathcal{P}(X) \rightarrow [0, \infty]$ be the counting measure on X . If $E \subseteq X$ and $\mu(E) = \infty$, then taking any point $x \in E$, we have $\mu(\{x\}) = 1 < \infty$, so μ is a semi-finite measure. However, since

X is uncountable, μ cannot be expressed as a countable sum of finite measures, so μ is not s-finite.

Most texts on measure theory focus on σ -finite measures and omit mention of the more general concept of s-finite measures. However, as we will see, s-finite measures are the natural class of measures for many important results in measure theory. One reason to appreciate the generality of s-finite measures is provided by the next theorem. We will encounter other advantages of working with s-finite measures later on in the course.

THEOREM 4.4

- (1) Let (X, \mathcal{B}, μ) be an s-finite measure space, and let (Y, \mathcal{C}) be a measurable space. Suppose $\pi : X \rightarrow Y$ is a measurable map. Then the measure $\pi_*\mu : \mathcal{C} \rightarrow [0, \infty]$ defined by $\pi_*\mu(C) = \mu(\pi^{-1}(C))$ is s-finite.
- (2) There exists a σ -finite measure space (X, \mathcal{B}, μ) , a measurable space (Y, \mathcal{C}) , and a measurable map $\pi : X \rightarrow Y$ such that $\pi_*\mu$ is not σ -finite.

PROOF. (1) First note that the projection of a finite measure is finite. Indeed, $\pi_*\mu(Y) = \mu(\pi^{-1}(Y)) = \mu(X)$. Noting that $\pi_*(\sum_{n=1}^{\infty} \mu_n) = \sum_{n=1}^{\infty} \pi_*\mu_n$ then completes the proof.

(2) Let $X = \mathbb{Z}^2$, $\mathcal{B} = \mathcal{P}(\mathbb{Z}^2)$, and let $\mu : \mathcal{B} \rightarrow [0, \infty]$ be the counting measure. Let $Y = \mathbb{Z}$ and $\mathcal{C} = \mathcal{P}(\mathbb{Z})$, and let $\pi : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ be the projection onto the first coordinate, i.e. $\pi(n, m) = n$ for $(n, m) \in \mathbb{Z}^2$. Then $\pi_*\mu(E)$ counts the number of points in \mathbb{Z}^2 whose first coordinate belongs to E . Hence, $\pi_*\mu(E) = \infty$ whenever $E \neq \emptyset$. Therefore, $\pi_*\mu$ is not σ -finite (nor even semi-finite). \square

3. Decompositions of Measures

When we say that two notions are “complementary,” we mean that they are mutually exclusive and every (σ -finite) measure can be decomposed into pieces satisfying one or the other property. Namely, for the complementary notions shown in Figure 4.1, we have the following decomposition result:

PROPOSITION 4.5

- (1) Let (X, \mathcal{B}, μ) be a σ -finite measure space. Then there is a unique decomposition $\mu = \mu_a + \mu_{na}$ as a sum of a purely atomic measure μ_a and a non-atomic measure μ_{na} .
- (2) Let (X, \mathcal{B}, μ) be an s-finite measure space, and suppose $\{x\} \in \mathcal{B}$ for every $x \in X$. Then there is a unique decomposition $\mu = \mu_d + \mu_c$ as a sum of a discrete measure μ_d and a continuous measure μ_c .

In general, *atomic* and *discrete* are different notions.

EXAMPLE 4.6

Let X be an uncountable set, and let $\mathcal{B} = \{E \subseteq X : E \text{ is countable or } X \setminus E \text{ is countable}\}$. Define a probability measure $\mu : \mathcal{B} \rightarrow [0, 1]$ by $\mu(E) = 0$ if E is countable and $\mu(E) = 1$ if $X \setminus E$ is uncountable. Then E is atomic (each co-countable set is an atom) but also continuous.

However, in many frequently-encountered situations, atomic and discrete measures coincide.

THEOREM 4.7

Let X be a separable metric space. Suppose μ is a locally finite Borel measure on X . If $A \in \text{Borel}(X)$ is an atom of μ , then there is a point $x \in A$ such that $\mu(\{x\}) = \mu(A) > 0$. Hence, every atomic locally finite Borel measure on X is discrete.

Proving the decomposition of a σ -finite measure into atomic and non-atomic components is a bit lengthy, so we will prove only part (2) of Proposition 4.5. Because of Theorem 4.7, the decomposition into discrete and continuous parts is sufficient for most purposes.

PROOF OF PROPOSITION 4.5(2).

STEP 1. Existence.

Since μ is s-finite, we may write $\mu = \sum_{n=1}^{\infty} \mu_n$ for some finite measures $\mu_n : \mathcal{B} \rightarrow [0, \infty)$. For each $k \in \mathbb{N}$, let $X_{n,k} = \{x \in X : \mu_n(\{x\}) \geq \frac{1}{k}\}$. Note that $X_{n,k}$ has at most $k\mu_n(X)$ elements for each $n, k \in \mathbb{N}$. Therefore, $X_0 = \{x \in X : \mu(\{x\}) > 0\} = \bigcup_{n \in \mathbb{N}} \bigcup_{k \in \mathbb{N}} X_{n,k}$ is a countable set.

For $x \in X_0$, let $c_x = \mu(\{x\})$. Define $\mu_d = \sum_{x \in X_0} c_x \delta_x$, and let $\mu_c : \mathcal{B} \rightarrow [0, \infty)$ be the measure $\mu_c(E) = \mu(E \setminus X_0)$ for $E \in \mathcal{B}$. Then μ_d is manifestly a discrete measure. Moreover, for any $x \in X$,

$$\mu_c(\{x\}) = \mu(\{x\} \setminus X_0) = \begin{cases} \mu(\{x\}), & \text{if } x \notin X_0; \\ 0, & \text{if } x \in X_0. \end{cases}$$

Since X_0 is the set of all point masses for μ , it follows that $\mu_c(\{x\}) = 0$ for every $x \in X$; that is, μ_c is continuous. Finally, for any $E \in \mathcal{B}$,

$$\mu(E) = \mu(E \cap X_0) + \mu_c(E)$$

and

$$\mu(E \cap X_0) = \sum_{x \in E \cap X_0} \mu(\{x\}) = \sum_{x \in X_0} c_x \delta_x(E) = \mu_d(E).$$

STEP 2. Uniqueness.

Let $\mu = \mu_d + \mu_c$ be the decomposition obtained in Step 1. Suppose $\mu = \mu'_d + \mu'_c$ is another decomposition into a discrete measure μ'_d and a continuous measure μ'_c . We want to show $\mu'_d = \mu_d$ and $\mu'_c = \mu_c$.

Let $x \in X_0$. Since μ'_c is continuous, we have $\mu'_c(\{x\}) = 0$, so $\mu'_d(\{x\}) = \mu(\{x\}) = c_x$. On the other hand, if $x \in X$ is any point and $\mu'_d(\{x\}) > 0$, then $\mu(\{x\}) \geq \mu'_d(\{x\}) > 0$, so $x \in X_0$. Therefore, the point masses of μ'_d are exactly the elements of X_0 , and $\mu'_d(\{x\}) = c_x$ for $x \in X_0$. Since μ'_d is discrete, it can thus be represented as $\mu'_d = \sum_{x \in X_0} c_x \delta_x$. That is, $\mu'_d = \mu_d$, and it follows that we also have $\mu'_c = \mu_c$.

□

The condition of semi-finiteness also leads to a decomposition result.

PROPOSITION 4.8

Let (X, \mathcal{B}, μ) be a measure space. There exists a decomposition $\mu = \mu_{\text{sf}} + \mu_{\text{inf}}$ such that μ_{sf} is semi-finite and μ_{inf} takes only the values 0 and ∞ .

Unlike the decompositions in Proposition 4.5, the decomposition in Proposition 4.8 is not unique in general. One way of obtaining the decomposition is to define

$$\mu_{\text{sf}}(E) = \sup \{ \mu(F) : F \in \mathcal{B}, F \subseteq E, \text{ and } \mu(F) < \infty \},$$

and

$$\mu_{\text{inf}}(E) = \begin{cases} 0, & \text{if } E \text{ is semi-finite;} \\ \infty, & \text{if } E \text{ is not semi-finite.} \end{cases}$$

Here, we say that a measurable set E is semi-finite if the measure $\mu_E : \mathcal{B} \rightarrow [0, \infty]$ defined by $\mu_E(A) = \mu(A \cap E)$ is a semi-finite measure. In other words, $E \in \mathcal{B}$ is semi-finite if every subset of E of infinite measure has a further subset of positive finite measure.

Exercises

4.1 Show that every σ -finite measure is s-finite and semi-finite.

4.2 In this exercise, we will construct a *saturation* of a measure space similarly to how we constructed the completion of a measure space.

Let (X, \mathcal{B}, μ) be a measure space.

(a) Show that the family

$$\tilde{\mathcal{B}} = \{ E \subseteq X : E \cap F \in \mathcal{B} \text{ for every } F \in \mathcal{B} \text{ with } \mu(F) < \infty \}$$

of locally measurable sets is a σ -algebra on X .

(b) Define a function $\tilde{\mu} : \tilde{\mathcal{B}} \rightarrow [0, \infty]$ by

$$\tilde{\mu}(E) = \begin{cases} \mu(E), & \text{if } E \in \mathcal{B}; \\ \infty, & \text{if } E \notin \mathcal{B}. \end{cases}$$

Show that $\tilde{\mu}$ is a measure and $(X, \tilde{\mathcal{B}}, \tilde{\mu})$ is saturated.

4.3 Suppose (X, \mathcal{B}, μ) is a non-atomic probability space. The goal of this problem is to prove Sierpiński's theorem: if $t \in [0, 1]$, then there exists a set $E \in \mathcal{B}$ with $\mu(E) = t$.

(a) Show for any $s \in (0, 1)$, there exists a set $E \in \mathcal{B}$ such that $0 < \mu(E) < s$.

(b) Let $t \in (0, 1)$. Construct a sequence of disjoint measurable sets $(E_n)_{n \in \mathbb{N}}$ such that

(i) for each $n \in \mathbb{N}$, $\mu(E_1 \cup \dots \cup E_n) < t$, and

(ii) if possible, E_n is chosen so that $\mu(E_n) > \frac{1}{n}$.

Show that $\mu(\bigcup_{n \in \mathbb{N}} E_n) = t$.

CHAPTER 5

Lebesgue–Stieltjes Measures

Let us rephrase (an instance of) the problem of measurement using the language of abstract measure theory developed in Part 1.

PROBLEM 5.1: PROBLEM OF MEASUREMENT IN ONE DIMENSION

Construct a measure $\lambda : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ such that $\lambda(I) = \text{length}(I)$ for every interval $I \subseteq \mathbb{R}$. Is there a unique such measure? Can the measure be defined on all subsets of \mathbb{R} ?

We will address this problem in a more general framework, where we allow for different assignments of measure to intervals.

DEFINITION 5.2

A Borel measure $\mu : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ is *locally finite* if $\mu(K) < \infty$ for every compact set $K \subseteq \mathbb{R}$. The *distribution function* of a locally finite Borel measure μ is the function $F_\mu : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$F_\mu(x) = \begin{cases} \mu((0, x]), & \text{if } x > 0; \\ 0, & \text{if } x = 0; \\ -\mu((x, 0]), & \text{if } x < 0. \end{cases}$$

By monotonicity of the measure μ , its distribution function F_μ is necessarily increasing. Moreover, by continuity from above and below, F_μ is a right-continuous function. The goal of this chapter is to prove the following theorem:

THEOREM 5.3: EXISTENCE AND UNIQUENESS OF LEBESGUE–STIELTJES MEASURES

Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing, right-continuous function with $F(0) = 0$. There exists a σ -algebra \mathcal{M}_F containing the Borel subsets of \mathbb{R} and a complete measure $\mu_F : \mathcal{M}_F \rightarrow [0, \infty]$ such that $F = F_{\mu_F}$. Moreover, if $\nu : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ is a Borel measure satisfying $F_\nu = F$, then $\nu = \mu_F|_{\text{Borel}(\mathbb{R})}$, and μ_F is the completion of ν .

DEFINITION 5.4

Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing, right-continuous function. The unique complete measure μ_F given by Theorem 5.3 is called the *Lebesgue–Stieltjes measure* associated to F .

PROPOSITION 5.5

Let F be an increasing, right-continuous function, and let μ_F be the Lebesgue–Stieltjes measure associated to F . Then

$$\lim_{x \rightarrow \infty} F(x) = \sup_{x \in \mathbb{R}} F(x) = \mu_F((0, \infty)) \quad \text{and} \quad \lim_{x \rightarrow -\infty} F(x) = \inf_{x \in \mathbb{R}} F(x) = -\mu_F((-\infty, 0]).$$

PROOF. This is an application of continuity from below of the measure μ_F . □

NOTATION. Given an increasing, right-continuous function, we will write $F(\infty)$ for the value $\lim_{x \rightarrow \infty} F(x)$ and $F(-\infty) = \lim_{x \rightarrow -\infty} F(x)$. In general, $F(\pm\infty)$ is an extended real number.

1. The π - λ Theorem and Uniqueness of Lebesgue–Stieltjes Measures

Before constructing Lebesgue–Stieltjes measures, let us prove that every locally finite Borel measure is uniquely determined by its distribution function. The key tool will be the π - λ theorem, for which we need a new definition.

DEFINITION 5.6

Let X be a set.

- A family $\mathcal{P} \subseteq \mathcal{P}(X)$ of subsets of X is a *π -system* if \mathcal{P} is closed under finite intersections.
- A family $\mathcal{L} \subseteq \mathcal{P}(X)$ is a *λ -system* if $\emptyset \in \mathcal{L}$ and \mathcal{L} is closed under complements and countable disjoint unions.

EXAMPLE 5.7

The following are examples of π systems:

- the collection $\mathcal{P} = \{(a, b] : a, b \in \mathbb{R}\}$ of half-open intervals in \mathbb{R} ;
- the family of open sets of any topological space;
- given a measure space (X, \mathcal{B}, μ) , the family $\mathcal{P} = \{E \in \mathcal{B} : \mu(X \setminus E) = 0\}$ of co-null sets;
- given two measurable spaces (X, \mathcal{B}) and (Y, \mathcal{C}) , the family $\mathcal{P} = \{B \times C : B \in \mathcal{B}, C \in \mathcal{C}\}$ of “rectangles” in $X \times Y$.

Examples of λ -systems include:

- for two probability measures μ, ν on a measurable space (X, \mathcal{B}) , the family $\mathcal{L} = \{E \in \mathcal{B} : \mu(E) = \nu(E)\}$.

Another characterization of λ -systems is given by the following proposition:

PROPOSITION 5.8

Let X be a set. A family $\mathcal{L} \subseteq \mathcal{P}(X)$ is a λ -system if and only if it satisfies the following three properties:

- (1) $X \in \mathcal{L}$;
- (2) if $A, B \in \mathcal{L}$ and $A \subseteq B$, then $B \setminus A \in \mathcal{L}$;
- (3) if $A_1 \subseteq A_2 \subseteq \dots$ is an increasing sequence in \mathcal{L} , then $\bigcup_{n \in \mathbb{N}} A_n \in \mathcal{L}$.

PROOF. Suppose \mathcal{L} is a λ -system. We check that \mathcal{L} satisfies properties (1)–(3).

(1) Since $\emptyset \in \mathcal{L}$ and \mathcal{L} is closed under complements, we have $X \in \mathcal{L}$.

(2) Let $A, B \in \mathcal{L}$ with $A \subseteq B$. Then $B \setminus A = B \cap A^c = (B^c \cup A)^c$. The assumption $A \subseteq B$ means $B^c \cap A = \emptyset$, so we have represented $B \setminus A$ in terms of A and B using complementation and disjoint union. Hence, $B \setminus A \in \mathcal{L}$.

(3) Let $A_1 \subseteq A_2 \subseteq \dots$ be an increasing sequence in \mathcal{L} . Let $B_1 = A_1$ and $B_n = A_n \setminus A_{n-1}$ for $n \geq 2$. By property (ii), $B_n \in \mathcal{L}$ for every $n \in \mathbb{N}$. Therefore, $\bigcup_{n \in \mathbb{N}} A_n = \bigsqcup_{n \in \mathbb{N}} B_n \in \mathcal{L}$.

Conversely, suppose $\mathcal{L} \subseteq \mathcal{P}(X)$ is a family of sets satisfying (1), (2), and (3).

Applying property (2) with $A = B = X$, we have $\emptyset = X \setminus X \in \mathcal{L}$.

Let $A \in \mathcal{L}$. Combining (1) and (2), $A^c = X \setminus A \in \mathcal{L}$.

Finally, let $(A_n)_{n \in \mathbb{N}}$ be a pairwise disjoint sequence of sets in \mathcal{L} . Then $B_n = A_1 \sqcup \dots \sqcup A_n$ forms an increasing sequence, so by property (3), it suffices to prove that $B_n \in \mathcal{L}$. By induction, this reduces to showing that the disjoint union of two sets in \mathcal{L} is an element of \mathcal{L} . Let $C, D \in \mathcal{L}$ with $C \cap D = \emptyset$. Then $C \sqcup D = (C^c \cap D^c)^c = (C^c \setminus D)^c$. We have already checked that \mathcal{L} is closed under complementation. The disjointness of C and D implies $D \subseteq C^c$, so $C^c \setminus D \in \mathcal{L}$ by property (2). Thus, $C \sqcup D \in \mathcal{L}$. \square

THEOREM 5.9: π - λ THEOREM (SIERPIŃSKI–DYNKIN)

Let X be a set, and suppose $\mathcal{P} \subseteq \mathcal{P}(X)$ is a π -system. If $\mathcal{L} \subseteq \mathcal{P}(X)$ is a λ -system and $\mathcal{P} \subseteq \mathcal{L}$, then $\sigma(\mathcal{P}) \subseteq \mathcal{L}$.

We will prove the π - λ theorem with the help of several lemmas.

LEMMA 5.10

Let X be a set. A family $\mathcal{B} \subseteq \mathcal{P}(X)$ of subsets of X is a σ -algebra if and only if \mathcal{B} is both a π -system and a λ -system.

PROOF. The definition of a λ -system is the same as the definition of a σ -algebra, except that one is only allowed to take unions of *disjoint* sets in the definition of a λ -system. It therefore suffices to check that being a π -system as well allows for taking countable unions of not necessarily disjoint sets.

Suppose $E_1, E_2, \dots \in \mathcal{B}$. Define $E'_1 = E_1$, $E'_2 = E_2 \setminus E_1$, \dots , $E'_n = E_n \setminus \bigcup_{i=1}^{n-1} E_i$. Then E'_1, E'_2, \dots are pairwise disjoint and satisfy $\bigsqcup_{n \in \mathbb{N}} E'_n = \bigcup_{n \in \mathbb{N}} E_n$, so it suffices to check that $E'_n \in \mathcal{B}$ for each $n \in \mathbb{N}$. But this is clear upon rewriting $E'_n = E_n \cap \bigcap_{i=1}^{n-1} E_i^c$, since $E_i^c = X \setminus E_i \in \mathcal{B}$ (by the axioms of a λ -system) and a finite intersection of sets from \mathcal{B} belongs to \mathcal{B} (by the axioms of a π -system). \square

LEMMA 5.11

Let X be a set, and suppose $(\mathcal{L}_i)_{i \in I}$ is a collection of λ -systems $\mathcal{L}_i \subseteq \mathcal{P}(X)$. Then $\bigcap_{i \in I} \mathcal{L}_i$ is a λ -system.

PROOF. The proof is the same as the proof of Proposition 2.3, except we only allow disjoint unions. \square

DEFINITION 5.12

Let X be a set and $\mathcal{S} \subseteq \mathcal{P}(X)$ a family of subsets of X . The λ -system generated by \mathcal{S} is the smallest λ -system containing \mathcal{S} :

$$\lambda(\mathcal{S}) = \bigcap \{\mathcal{L} \subseteq \mathcal{P}(X) : \mathcal{L} \text{ is a } \lambda\text{-system, } \mathcal{S} \subseteq \mathcal{L}\}.$$

LEMMA 5.13

Let X be a set, and let $\mathcal{P} \subseteq \mathcal{P}(X)$ be a π -system. The λ -system $\lambda(\mathcal{P})$ generated by \mathcal{P} is a σ -algebra.

PROOF. By Lemma 5.10, it suffices to show that $\lambda(\mathcal{P})$ is a π -system.

CLAIM 1. For any set $A \in \lambda(\mathcal{P})$, the family $\mathcal{L}_A := \{B \subseteq X : A \cap B \in \lambda(\mathcal{P})\}$ is a λ -system.

Since $A \in \lambda(\mathcal{P})$, we see that $X \in \mathcal{L}_A$.

Suppose $B_1, B_2 \in \mathcal{L}_A$ and $B_1 \subseteq B_2$. Then

$$A \cap (B_2 \setminus B_1) = \underbrace{(A \cap B_2)}_{\in \lambda(\mathcal{P})} \setminus \underbrace{(A \cap B_1)}_{\in \lambda(\mathcal{P})} \in \lambda(\mathcal{P}),$$

so $B_2 \setminus B_1 \in \mathcal{L}_A$.

Finally, suppose $B_1 \subseteq B_2 \subseteq \dots \in \mathcal{L}_A$. Then $A \cap \bigcup_{n \in \mathbb{N}} B_n = \bigcup_{n \in \mathbb{N}} (A \cap B_n) \in \lambda(\mathcal{P})$, so $\bigcup_{n \in \mathbb{N}} B_n \in \mathcal{L}_A$.

This proves the claim.

CLAIM 2. For any $A \in \lambda(\mathcal{P})$ and any $B \in \mathcal{P}$, we have $A \cap B \in \lambda(\mathcal{P})$.

This follows from Claim 1: the family \mathcal{L}_B is a λ -system, and $\mathcal{P} \subseteq \mathcal{L}_B$ by the definition of a π -system, so $\mathcal{L}_B \supseteq \lambda(\mathcal{P}) \ni A$.

Let $A, B \in \lambda(\mathcal{P})$. The family \mathcal{L}_A is a λ -system (by Claim 1) containing \mathcal{P} (by Claim 2), so $\mathcal{L}_A \supseteq \lambda(\mathcal{P}) \ni B$. Hence, $A \cap B \in \lambda(\mathcal{P})$. □

Now we can complete the proof of the π - λ theorem.

PROOF OF π - λ THEOREM (THEOREM 5.9). Let \mathcal{P} be a π -system, \mathcal{L} a λ -system, and suppose $\mathcal{P} \subseteq \mathcal{L}$. On the one hand, by Lemma 5.13, the λ -system $\lambda(\mathcal{P})$ generated by \mathcal{P} is a σ -algebra, so $\sigma(\mathcal{P}) \subseteq \lambda(\mathcal{P})$. On the other hand, \mathcal{L} is a λ -system containing \mathcal{P} , so $\lambda(\mathcal{P}) \subseteq \mathcal{L}$. Combining these two observations completes the proof. □

COROLLARY 5.14: UNIQUENESS OF LEBESGUE-STIELTJES MEASURES

Suppose μ and ν are locally finite Borel measures on \mathbb{R} with the same distribution function $F_\mu = F_\nu = F$. Then $\mu = \nu$.

PROOF. Let \mathcal{P} be the π -system $\mathcal{P} = \{(a, b] : a, b \in \mathbb{R}\}$ of half-open intervals. Define

$$\mathcal{L} = \{E \in \text{Borel}(\mathbb{R}) : \mu(E \cap (-N, N]) = \nu(E \cap (-N, N]) \text{ for every } N \in \mathbb{N}\}.$$

CLAIM 1. \mathcal{L} is a λ -system

For every $N \in \mathbb{N}$,

$$\mu((-N, N]) = (\mu((-N, 0]) + \mu((0, N])) = F(N) - F(-N).$$

The same holds for ν , so $\mathbb{R} \in \mathcal{L}$.

Suppose $E \in \mathcal{L}$, and let $N \in \mathbb{N}$. By additivity of μ and ν , we have

$$\begin{aligned}\mu(E^c \cap (-N, N]) &= \mu((-N, N]) - \mu(E \cap (-N, N]) \\ &= \nu((-N, N]) - \nu(E \cap (-N, N]) \\ &= \nu(E^c \cap (-N, N]),\end{aligned}$$

so $E^c \in \mathcal{L}$.

Finally, \mathcal{L} is closed under countable disjoint unions as a consequence of countable additivity of the measures μ and ν .

CLAIM 2. $\mathcal{P} \subseteq \mathcal{L}$

The sets $(-N, N]$ belong to \mathcal{P} , which is a π -system, so it suffices to prove $\mu(P) = \nu(P)$ for every $P \in \mathcal{P}$. Let $P = (a, b] \in \mathcal{P}$. If $b \leq a$, then $P = \emptyset$, so $\mu(P) = \nu(P) = 0$. Suppose $a < b$. If $a \leq b$, then $\mu(P) = F(b) - F(a) = \nu(P)$.

By the π - λ theorem, $\sigma(\mathcal{P}) \subseteq \mathcal{L}$. But \mathcal{P} generates the Borel σ -algebra (we essentially showed this in the proof of Proposition 2.11), so $\mathcal{L} = \text{Borel}(\mathbb{R})$. Hence, applying continuity from below, we have

$$\mu(E) = \lim_{N \rightarrow \infty} \mu(E \cap (-N, N]) = \lim_{N \rightarrow \infty} \nu(E \cap (-N, N]) = \nu(E)$$

for every $E \in \text{Borel}(\mathbb{R})$. □

EXAMPLE 5.15

The locally finite condition cannot be dropped from Corollary 5.14. As an example, define a measure $\mu : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ by

$$\mu(B) = \#(B \cap \mathbb{Q}),$$

and let $\nu = \infty \cdot \lambda$, where λ is the Lebesgue measure. (We will construct λ later in this chapter, but for now, take it as a given that the Lebesgue measure exists.) Every non-empty interval in \mathbb{R} contains infinitely many rational points, so

$$F_\mu(x) = \begin{cases} \infty, & \text{if } x > 0; \\ 0, & \text{if } x = 0; \\ -\infty, & \text{if } x < 0. \end{cases}$$

Similarly, every non-empty interval in \mathbb{R} has positive Lebesgue measure, so multiplying by ∞ , the measure ν has the same distribution function $F_\nu = F_\mu$. However, μ and ν are not the same measure, since, for instance, $\mu(\mathbb{R} \setminus \mathbb{Q}) = 0$, while $\nu(\mathbb{R} \setminus \mathbb{Q}) = \infty$, and $\mu(\{0\}) = 1$, while $\nu(\{0\}) = 0$.

2. Half-Open Intervals

Now we begin the construction of Lebesgue–Stieltjes measures. Let us define some basic objects that we will work with for the construction.

DEFINITION 5.16

A *left-open, right-closed interval* is a set of the form

- \mathbb{R} ,
- \emptyset ,
- $(a, b]$ with $a, b \in \mathbb{R}$, $a < b$,
- $(-\infty, b]$ with $b \in \mathbb{R}$, or
- (a, ∞) with $a \in \mathbb{R}$.

The intersection of two left-open, right-closed intervals is a left-open, right-closed intervals, and the complement of a left-open, right-closed interval is either a left-open, right-closed interval or a disjoint union of two left-open, right-closed intervals. Therefore, the family of left-open, right-closed intervals forms a *semi-algebra* on \mathbb{R} . We recall the definition below.

DEFINITION 5.17

Let X be a set. A family $\mathcal{S} \subseteq \mathcal{P}(X)$ of subsets of X is a *semi-algebra* if

- $\emptyset, X \in \mathcal{S}$;
- if $A, B \in \mathcal{S}$, then $A \cap B \in \mathcal{S}$;
- if $A \in \mathcal{S}$, then $X \setminus A = \bigsqcup_{i=1}^n C_i$ for some $C_1, \dots, C_n \in \mathcal{S}$.

In Exercise 2.1, you showed the following fact:

PROPOSITION 5.18

Let \mathcal{S} be a semi-algebra on a set X . Then

$$\mathcal{A} = \left\{ \bigsqcup_{i=1}^n S_i : n \in \mathbb{N}, S_1, \dots, S_n \in \mathcal{S} \right\}$$

is an algebra.

NOTATION. We will denote the algebra generated by the semi-algebra of left-open, right-closed intervals by

$$\mathcal{A}_{int} = \left\{ \bigsqcup_{i=1}^n I_i : n \in \mathbb{N}, I_i \text{ is a left-open, right-closed interval} \right\}.$$

Note that the σ -algebra generated by \mathcal{A}_{int} is the Borel σ -algebra.

3. Premeasures and Outer Measures

We will begin the construction of the Lebesgue–Stieltjes measure associated to a distribution function F by assigning a measure to each element of \mathcal{A}_{int} .

DEFINITION 5.19

Let X be a set and $\mathcal{A} \subseteq \mathcal{P}(X)$ an algebra. A *premeasure* is a function $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ such that

- $\mu_0(\emptyset) = 0$;
- if $(A_n)_{n \in \mathbb{N}}$ is a sequence of pairwise disjoint elements of \mathcal{A} and $A = \bigsqcup_{n \in \mathbb{N}} A_n \in \mathcal{A}$, then $\mu_0(A) = \sum_{n=1}^{\infty} \mu_0(A_n)$.

Note that if \mathcal{A} is a σ -algebra, then a premeasure on \mathcal{A} is the same thing as a measure. More generally, if $\mu : \mathcal{B} \rightarrow [0, \infty]$ is a measure on a measurable space (X, \mathcal{B}) and $\mathcal{A} \subseteq \mathcal{B}$ is an algebra, then $\mu_0 = \mu|_{\mathcal{A}}$ defines a premeasure on \mathcal{A} .

The next proposition shows that we can associate to an increasing right-continuous function F a premeasure on the algebra \mathcal{A}_{int} generated by left-open, right-closed intervals. We will see afterwards how to extend this premeasure to a measure on the Borel subsets of \mathbb{R} .

PROPOSITION 5.20

Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be increasing and right-continuous. Define a function $\mu_{F,0} : \mathcal{A}_{int} \rightarrow [0, \infty]$ by

$$\mu_{F,0} \left(\bigsqcup_{i=1}^n (a_i, b_i] \right) = \sum_{i=1}^n (F(b_i) - F(a_i)).$$

Then $\mu_{F,0}$ is a premeasure on \mathcal{A}_{int} .

PROOF. We will first show that $\mu_{F,0}$ is a well-defined function on \mathcal{A}_{int} and then prove that it is a premeasure.

STEP 1. $\mu_{F,0}$ is well-defined

Every element of \mathcal{A}_{int} can always be written uniquely in the form

$$\bigsqcup_{i=1}^n (a_i, b_i]$$

with $-\infty \leq a_1 < b_1 < a_2 < b_2 < \dots < a_n < b_n \leq \infty$. Indeed, after writing the intervals in increasing order, if $a_{i+1} = b_i$ for some i , then the intervals $(a_i, b_i]$ and $(a_{i+1}, b_{i+1}]$ can be merged into the single interval $(a_i, b_{i+1}]$. This process of merging leaves the expression for $\mu_{F,0}$ unchanged, since if $b_i = a_{i+1}$, we have a telescoping phenomenon

$$(F(b_i) - F(a_i)) + (F(b_{i+1}) - F(a_{i+1})) = F(b_{i+1}) - F(a_i).$$

Thus, the formula for $\mu_{F,0}$ gives the same value for every possible expression of $A \in \mathcal{A}_{int}$ as a disjoint union of left-open, right-closed intervals.

STEP 2. If $(a, b] = \bigsqcup_{i=1}^{\infty} (a_i, b_i]$, then $\mu_{F,0}((a, b]) \leq \sum_{i=1}^{\infty} \mu_{F,0}((a_i, b_i])$.

Let $\delta > 0$ and let $\varepsilon_i > 0$ for $i \in \mathbb{N}$. Then $[a + \delta, b]$ is a closed interval covered by the union of open intervals $\bigcup_{i=1}^{\infty} (a_i, b_i + \varepsilon_i)$. By the Heine–Borel theorem (compactness of closed intervals in \mathbb{R}), there is a finite subcover i_1, \dots, i_n such that $[a + \delta, b] \subseteq \bigcup_{j=1}^n (a_{i_j}, b_{i_j} + \varepsilon_{i_j})$. Therefore, $(a + \delta, b] \subseteq \bigcup_{j=1}^n (a_{i_j}, b_{i_j} + \varepsilon_{i_j})$, so by Step 1,

$$\mu_{F,0}((a + \delta, b]) \leq \sum_{j=1}^n \mu_{F,0}((a_{i_j}, b_{i_j} + \varepsilon_{i_j}]) \leq \sum_{i=1}^{\infty} \mu_{F,0}((a_i, b_i + \varepsilon_i]).$$

Letting $\delta \rightarrow 0$,

$$\lim_{\delta \rightarrow 0^+} \mu_{F,0}((a + \delta, b]) = F(b) - \lim_{\delta \rightarrow 0^+} F(a + \delta) = F(b) - F(a) = \mu_{F,0}((a, b]),$$

since F is right-continuous. Similarly, given $\varepsilon > 0$, we can take ε_i sufficiently small so that $\sum_{i=1}^{\infty} \mu_{F,0}((a_i, b_i + \varepsilon_i)) \leq \sum_{i=1}^{\infty} \mu_{F,0}((a_i, b_i]) + \varepsilon$. Then letting $\varepsilon \rightarrow 0$ proves the desired inequality.

STEP 3. $\mu_{F,0}$ is countably additive.

Suppose $(A_n)_{n \in \mathbb{N}}$ is a sequence of pairwise disjoint elements of \mathcal{A}_{int} , and $A = \bigsqcup_{n \in \mathbb{N}} A_n \in \mathcal{A}_{int}$. Each of the sets A_n , belonging to the algebra \mathcal{A}_{int} , can be written in the form $A_n = \bigsqcup_{m=1}^{M_n} S_{n,m}$, where $S_{n,m}$ is a left-open, right-closed interval. We know $\mu_{F,0}(A_n) = \sum_{m=1}^{M_n} \mu_{F,0}(S_{n,m})$ by definition. Replacing $(A_n)_{n \in \mathbb{N}}$ by $(S_{n,m})_{n \in \mathbb{N}, 1 \leq m \leq M_n}$, we may assume from the start that A_n is a left-open, right-closed interval for each $n \in \mathbb{N}$.

Since $A \in \mathcal{A}$, we may also write $A = \bigsqcup_{m=1}^M S_m$ for some left-open, right-closed intervals S_m . Then

$$\begin{aligned} \mu_{F,0}(A) &= \sum_{m=1}^M \mu_{F,0}(S_m) && \text{(definition of } \mu_{F,0}) \\ &= \sum_{m=1}^M \mu_{F,0} \left(\bigsqcup_{n \in \mathbb{N}} (S_m \cap A_n) \right) && (A = \bigsqcup_{n \in \mathbb{N}} A_n) \\ &\leq \sum_{n,m} \mu_{F,0}(S_m \cap A_n) && \text{(Step 2)} \\ &= \sum_{n=1}^{\infty} \mu_{F,0} \left(\bigsqcup_{m=1}^M (S_m \cap A_n) \right) && \text{(definition of } \mu_{F,0}) \\ &= \sum_{n=1}^{\infty} \mu_{F,0}(A_n) && \text{(definition of } \mu_{F,0}) \end{aligned}$$

On the other hand, for $N \in \mathbb{N}$,

$$\sum_{n=1}^N \mu_{F,0}(A_n) = \mu_{F,0} \left(\bigsqcup_{n=1}^N A_n \right) \leq \mu_{F,0}(A),$$

so taking a limit as $N \rightarrow \infty$, $\sum_{n=1}^{\infty} \mu_{F,0}(A_n) \leq \mu_{F,0}(A)$.

□

The next stage in the construction is to extend the premeasure $\mu_{F,0}$ to an *outer measure* defined on all subsets of \mathbb{R} .

DEFINITION 5.21

Let X be a set. A function $\mu^* : \mathcal{P}(X) \rightarrow [0, \infty]$ is an *outer measure* if

- $\mu^*(\emptyset) = 0$;

- MONOTONE: if $A \subseteq B$, then $\mu^*(A) \leq \mu^*(B)$; and
- COUNTABLY SUBADDITIVE: for any sequence of sets $(A_n)_{n \in \mathbb{N}}$, one has $\mu^*(\bigcup_{n \in \mathbb{N}} A_n) \leq \sum_{n=1}^{\infty} \mu^*(A_n)$.

A premeasure can always be extended to an outer measure, as shown by the following proposition.

PROPOSITION 5.22

Let \mathcal{A} be an algebra on a set X , and suppose $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ is a premeasure. Then

$$\mu^*(E) = \inf \left\{ \sum_{n=1}^{\infty} \mu_0(A_n) : E \subseteq \bigcup_{n \in \mathbb{N}} A_n, A_n \in \mathcal{A} \right\}$$

defines an outer measure on X with $\mu^*|_{\mathcal{A}} = \mu_0$.

PROOF. Let us check the properties one at a time.

First, $\emptyset \in \mathcal{A}$, so $\mu^*(\emptyset) \leq \mu_0(\emptyset) = 0$.

Next, suppose $A \subseteq B$. Then any set containing B also contains A , so the expression defining $\mu^*(A)$ involves an infimum over a larger collection than the expression defining $\mu^*(B)$. Hence, $\mu^*(A) \leq \mu^*(B)$.

Now let us prove countable subadditivity. Let $(A_n)_{n \in \mathbb{N}}$ be a sequence of subsets of X . If

$$\sum_{n=1}^{\infty} \mu^*(A_n) = \infty,$$

there is nothing to show, so assume

$$\sum_{n=1}^{\infty} \mu^*(A_n) < \infty.$$

Let $\varepsilon > 0$. For each n , let $(A_{n,k})_{k \in \mathbb{N}}$ be a sequence of elements of \mathcal{A} such that $A_n \subseteq \bigcup_{k \in \mathbb{N}} A_{n,k}$ and

$$\sum_{k=1}^{\infty} \mu_0(A_{n,k}) < \mu^*(A_n) + \frac{\varepsilon}{2^n}.$$

Then $(A_{n,k})_{n,k \in \mathbb{N}}$ is a countable family of elements of the algebra \mathcal{A} , and $\bigcup_{n \in \mathbb{N}} A_n \subseteq \bigcup_{n,k \in \mathbb{N}} A_{n,k}$, so

$$\mu^* \left(\bigcup_{n \in \mathbb{N}} A_n \right) \leq \sum_{n,k} \mu_0(A_{n,k}) \leq \sum_{n=1}^{\infty} \left(\mu^*(A_n) + \frac{\varepsilon}{2^n} \right) = \sum_{n=1}^{\infty} \mu^*(A_n) + \varepsilon.$$

Letting $\varepsilon \rightarrow 0$ establishes countable subadditivity.

Finally, let us show $\mu^*|_{\mathcal{A}} = \mu_0$. Let $A \in \mathcal{A}$. Then by definition $\mu^*(A) \leq \mu_0(A)$. It remains to show $\mu^*(A) \geq \mu_0(A)$. Let $(A_n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{A} such that $A \subseteq \bigcup_{n \in \mathbb{N}} A_n$. Define a new sequence $(B_n)_{n \in \mathbb{N}}$ by $B_1 = A \cap A_1$ and $B_n = A \cap A_n \setminus (A_1 \cup \dots \cup A_{n-1})$. Since \mathcal{A} is an algebra, the sets B_n belong to \mathcal{A} . Moreover, $(B_n)_{n \in \mathbb{N}}$ is a sequence of pairwise disjoint sets whose union is $A \in \mathcal{A}$, so

$$\mu_0(A) = \sum_{n=1}^{\infty} \mu_0(B_n) = \sum_{n=1}^{\infty} (\mu_0(B_n) + \mu_0(A_n \setminus B_n)) \leq \sum_{n=1}^{\infty} \mu_0(A_n).$$

Taking an infimum over all such collections $(A_n)_{n \in \mathbb{N}}$ gives the desired inequality $\mu_0(A) \leq \mu^*(A)$. □

The outer measure μ_F^* obtained from the premeasure $\mu_{F,0}$ is not in general a measure on $\mathcal{P}(X)$. The problem is that, while μ_F^* is *subadditive*, it may fail to be additive. In order to obtain a measure, we restrict to the sets with better additive behavior.

DEFINITION 5.23

Let μ^* be an outer measure on a set X . A set $E \subseteq X$ is μ^* -measurable if for every $A \subseteq X$,

$$\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \setminus E). \quad (5.1)$$

REMARK. Outer measures are subadditive, so (5.1) is equivalent to the *a priori* weaker inequality

$$\mu^*(A) \geq \mu^*(A \cap E) + \mu^*(A \setminus E).$$

THEOREM 5.24: CARATHÉODORY'S THEOREM

Let μ^* be an outer measure on a set X . Let $\mathcal{M} \subseteq \mathcal{P}(X)$ be the family of μ^* -measurable sets. Then \mathcal{M} is a σ -algebra, and $\mu^*|_{\mathcal{M}}$ is a complete measure.

PROOF. We break the proof into several steps.

CLAIM 1. $X \in \mathcal{M}$.

Given $A \subseteq X$, we have $\mu^*(A \cap X) + \mu^*(A \setminus X) = \mu^*(A) + \mu^*(\emptyset) = \mu^*(A)$.

CLAIM 2. \mathcal{M} is closed under complementation.

Rewriting $A \setminus E = A \cap E^c$, the measurability condition (5.1) is symmetric in E and E^c .

CLAIM 3. \mathcal{M} is closed under finite unions.

Suppose $E, F \in \mathcal{M}$, and let $A \subseteq X$. We want to show

$$\mu^*(A) \geq \mu^*(A \cap (E \cup F)) + \mu^*(A \setminus (E \cup F)).$$

Writing $A \cap (E \cup F) = (A \cap E) \cup (A \cap F \cap E^c)$ and $A \setminus (E \cup F) = A \cap F^c \cap E^c$ and applying subadditivity of μ^* , we have

$$\mu^*(A \cap (E \cup F)) + \mu^*(A \setminus (E \cup F)) \leq \mu^*(A \cap E) + \underbrace{\mu^*(A \cap F \cap E^c)}_{\mu^*(A \cap E^c)} + \mu^*(A \cap F^c \cap E^c) = \mu^*(A).$$

Claims 1–3 show that \mathcal{M} is an algebra. The next claim upgrades \mathcal{M} to a σ -algebra and proves that $\mu^*|_{\mathcal{M}}$ is a measure.

CLAIM 4. \mathcal{M} is closed under countable disjoint unions, and $\mu^*|_{\mathcal{M}}$ is countably additive.

Suppose $(E_n)_{n \in \mathbb{N}}$ be a sequence of pairwise disjoint sets in \mathcal{M} , and let $E = \bigcup_{n \in \mathbb{N}} E_n$. Let $A \subseteq X$. As in Claim 3, we want to show

$$\mu^*(A) \geq \mu^*(A \cap E) + \mu^*(A \setminus E). \quad (5.2)$$

If $\mu^*(A) = \infty$, there is nothing to check, so assume $\mu^*(A) < \infty$. Let $F_N = \bigsqcup_{n=1}^N E_n$. By induction, we have

$$\mu^*(A \cap F_N) = \sum_{n=1}^N \mu^*(A \cap E_n).$$

Hence, by countable subadditivity of μ^* ,

$$\mu^*(A \cap E) \leq \sum_{n=1}^{\infty} \mu^*(A \cap E_n) = \lim_{N \rightarrow \infty} \mu^*(A \cap F_N).$$

For fixed $N \in \mathbb{N}$, $F_N \in \mathcal{M}$ by Claim 3, so

$$\mu^*(A) = \mu^*(A \cap F_N) + \mu^*(A \setminus F_N) \geq \mu^*(A \cap F_N) + \mu^*(A \setminus E).$$

Taking a limit as $N \rightarrow \infty$ gives (5.2).

Note that we actually proved the stronger inequality

$$\mu^*(A) \geq \sum_{n=1}^{\infty} \mu^*(A \cap E_n) + \mu^*(A \setminus E).$$

Taking $A = E$ establishes countable additivity of μ^* .

Finally, we check that $(X, \mathcal{M}, \mu^*|_{\mathcal{M}})$ is complete.

CLAIM 5. If $N \subseteq X$ and $\mu^*(N) = 0$, then $N \in \mathcal{M}$.

Let $A \subseteq X$. Then by monotonicity,

$$\mu^*(A \cap N) + \mu^*(A \setminus N) \leq \mu^*(N) + \mu^*(A) = \mu^*(A).$$

□

The last remaining piece to tie everything together is relating the σ -algebra \mathcal{M} in Carathéodory's theorem to the algebra on which the premeasure μ_0 was defined.

LEMMA 5.25

Let \mathcal{A} be an algebra on a set X . Let $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ be a premeasure, and let μ^* be the outer measure extending μ_0 as in Proposition 5.22. Then every element of \mathcal{A} is μ^* -measurable.

PROOF. Let $A \in \mathcal{A}$, and let $B \subseteq X$ be an arbitrary set. We want to show

$$\mu^*(B) \geq \mu^*(B \cap A) + \mu^*(B \setminus A). \quad (5.3)$$

Let $(A_n)_{n \in \mathbb{N}}$ be family of elements of \mathcal{A} such that $B \subseteq \bigcup_{n \in \mathbb{N}} A_n$. Then

$$\sum_{n=1}^{\infty} \mu_0(A_n) = \sum_{n=1}^{\infty} (\mu_0(A_n \cap A) + \mu_0(A_n \setminus A)) = \sum_{n=1}^{\infty} \mu_0(A_n \cap A) + \sum_{n=1}^{\infty} \mu_0(A_n \setminus A).$$

The union $\bigcup_{n \in \mathbb{N}} (A_n \cap A)$ contains $B \cap A$, and similarly, $\bigcup_{n \in \mathbb{N}} (A_n \setminus A)$ contains $B \setminus A$, so by the definition of the outer measure μ^* ,

$$\sum_{n=1}^{\infty} \mu_0(A_n) \geq \mu^*(B \cap A) + \mu^*(B \setminus A).$$

Taking an infimum over all such collections $(A_n)_{n \in \mathbb{N}}$ gives (5.3). □

Putting everything together, we get the following theorem.

THEOREM 5.26: HAHN–KOLMOGOROV EXTENSION THEOREM

Let $\mathcal{A} \subseteq \mathcal{P}(X)$ be an algebra on a set X , and let $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ be a premeasure. Then μ_0 extends to a complete measure $\mu : \mathcal{M} \rightarrow [0, \infty]$ defined on a σ -algebra $\mathcal{M} \supseteq \sigma(\mathcal{A})$. Moreover, if μ_0 is σ -finite, then the extension of μ_0 to $\sigma(\mathcal{A})$ is unique, and μ is the completion of this unique extension.

PROOF. By Proposition 5.22, μ_0 extends to an outer measure $\mu^* : \mathcal{P}(X) \rightarrow [0, \infty]$. Let \mathcal{M} be the σ -algebra of μ^* -measurable sets, and let $\mu = \mu^*|_{\mathcal{M}}$. Then μ is a complete measure by Carathéodory's theorem (Theorem 5.24), and by Lemma 5.25, $\mathcal{A} \subseteq \mathcal{M}$.

Uniqueness in the σ -finite case is a consequence of the π - λ theorem and follows on exactly the same lines as the proof of Corollary 5.14. \square

REMARK. When μ_0 is not σ -finite, it may have several different extensions to $\sigma(\mathcal{A})$. The outer measure construction is the maximal such extension in the sense that given any other extension $\nu : \sigma(\mathcal{A}) \rightarrow [0, \infty]$ of μ_0 , one has $\nu(E) \leq \mu(E)$ for every $E \in \sigma(\mathcal{A})$. We will return to this subject in the context of product measures later in the course, where it will sometimes be useful to work with a different extension of a premeasure than the one obtained by Carathéodory's theorem.

COROLLARY 5.27: EXISTENCE OF LEBESGUE–STIELTJES MEASURES

Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be increasing and right-continuous with $F(0) = 0$. There exists a σ -algebra \mathcal{M}_F containing the Borel subsets of \mathbb{R} and a complete measure $\mu_F : \mathcal{M}_F \rightarrow [0, \infty]$ with distribution function $F_{\mu_F} = F$.

PROOF. Let $\mu_{F,0}$ be the premeasure on \mathcal{A}_{int} given by Proposition 5.20. Then the extension μ_F given by the Hahn–Kolmogorov extension theorem (Theorem 5.26) is a complete measure with distribution function F . \square

4. Lebesgue Measure

DEFINITION 5.28

The *Lebesgue measure* on \mathbb{R} is the Lebesgue–Stieltjes measure associated to the distribution function $F(x) = x$.

PROPOSITION 5.29

Let λ be the Lebesgue measure on \mathbb{R} , and let \mathcal{M} be the σ -algebra of Lebesgue measurable sets.

- (1) TRANSLATION-INVARIANCE: $\lambda(E + t) = \lambda(E)$ for every $E \in \mathcal{M}$ and $t \in \mathbb{R}$;
- (2) REFLECTION-INVARIANCE: $\lambda(-E) = \lambda(E)$ for every $E \in \mathcal{M}$;
- (3) DILATION PROPERTY: $\lambda(tE) = |t|\lambda(E)$ for every $E \in \mathcal{M}$ and $t \in \mathbb{R}$;

PROOF. Compute the distribution function of the transformed measure and apply uniqueness of Lebesgue–Stieltjes measures (Corollary 5.14). We leave the details as an exercise. \square

Using the translation-invariance property of the Lebesgue measure, we can prove the existence of a non-measurable set.

THEOREM 5.30

There exists a Lebesgue non-measurable subset of \mathbb{R} .

PROOF. Define an equivalence relation on $[0, 1]$ by $x \sim y$ if $y - x \in \mathbb{Q}$. By the axiom of choice, let $E \subseteq [0, 1]$ be a set containing exactly one representative of each equivalence class. For each $t \in \mathbb{Q} \cap [0, 1]$, let $E_t = \{x + t \bmod 1 : x \in E\} \subseteq [0, 1]$.

CLAIM 1. The sets $(E_t)_{t \in \mathbb{Q} \cap [0, 1]}$ are pairwise disjoint.

For $t, s \in \mathbb{Q} \cap [0, 1]$ and $x, y \in E$, if $x + t \equiv y + s \pmod{1}$, then

$$y - x \equiv t - s \pmod{1},$$

so $x \sim y$. But E contains only one element from each equivalence class, so $x = y$ and $t = s$.

CLAIM 2. $\bigsqcup_{t \in \mathbb{Q} \cap [0, 1]} E_t = [0, 1]$

Let $x \in [0, 1]$. Then there exists $y \in E$ with $y \sim x$, since E has a representative of each equivalence class. Let $t = x - y \bmod 1 \in \mathbb{Q} \cap [0, 1]$. Then

$$y + t \equiv x \pmod{1}.$$

so $x \in E_t$.

Assume for contradiction that E is Lebesgue measurable.

CLAIM 3. For every $t \in \mathbb{Q} \cap [0, 1]$, E_t is Lebesgue measurable and $\lambda(E_t) = \lambda(E)$.

We can write

$$E_t = ((E + t) \cap [0, 1]) \sqcup ((E + t) \cap [1, 2]) - 1.$$

Therefore, by translation invariance of the Lebesgue measure,

$$\lambda(E_t) = \lambda(E + t) = \lambda(E).$$

Combining Claims 1–3 and using countable additivity of the Lebesgue measure,

$$1 = \lambda([0, 1]) = \sum_{t \in \mathbb{Q} \cap [0, 1]} \lambda(E_t) = \sum_{t \in \mathbb{Q} \cap [0, 1]} \lambda(E) = \infty \cdot \lambda(E).$$

There is no value of $\lambda(E)$ that can satisfy this equation. We have thus reached a contradiction, so E is non-measurable. \square

REMARK. The axiom of choice plays a crucial role in Theorem 5.30. Using the set-theoretic notion of an *inaccessible cardinal*, Robert Solovay constructed a model of set theory under the ZF axioms without choice in which every subset of \mathbb{R} is Lebesgue measurable [8].

5. Cantor Measure

Another interesting example of a Borel measure on the real line is the “uniform” measure on the middle-thirds Cantor set, which we will construct now. Recall that the Cantor set $C \subseteq [0, 1]$ is obtained by starting with the full interval $[0, 1]$ and iteratively removing the middle third of each remaining interval at each step. We can therefore write $[0, 1] \setminus C = \bigcup_{n=0}^{\infty} \bigcup_{k=0}^{2^n - 1} I_{n,k}$, where

$I_{n,0}, \dots, I_{n,2^{n-1}}$ is an enumeration of the removed intervals of length $3^{-(n+1)}$ in increasing order. How should we define the distribution function for a uniform measure on C ? Well, after step $n = 0$, where we remove the interval $(\frac{1}{3}, \frac{2}{3})$, we have half of the Cantor set to the left of this interval and the other half to the right, so the distribution function should take the value $\frac{1}{2}$ on the entirety of this interval. Arguing similarly, the distribution function should take the value $\frac{2k+1}{2^{n+1}}$ on the interval $I_{n,k}$ for each $n \geq 0$ and $k \in \{0, 1, \dots, 2^n - 1\}$. Since the union of the intervals $\bigcup_{n=0}^{\infty} \bigcup_{k=0}^{2^n-1} I_{n,k}$ are dense in $[0, 1]$, there is a unique way of interpolating between the values on the intervals $I_{n,k}$ in order to obtain a continuous function. We call this continuous function the *Cantor function* and its associated Lebesgue–Stieltjes measure the *Cantor measure*.




FIGURE 5.1. Iterative construction of the Cantor function

A more explicit description of the Cantor set is the collection of numbers in the interval $[0, 1]$ whose binary expansion consists entirely of the digits 0 and 2. From this description of the Cantor set, we can also obtain a formula for the Cantor function, namely

$$c(x) = \begin{cases} \sum_{j=1}^{\infty} \frac{a_j/2}{2^j}, & \text{if } x = \sum_{j=1}^{\infty} \frac{a_j}{3^j} \in C; \\ \sup_{y \leq x, y \in C} c(y), & \text{if } x \notin C. \end{cases}$$

The Cantor measure has a surprising combination of properties. Suppose you have a perfectly fair coin to flip, and you record the sequence of heads and tails as you flip the coin repeatedly. Recording heads as the digit 2 and tails as the digit 0, this sequence of coin flips produces a random element of the Cantor set in terms of its base 3 expansion. The distribution of this random element of the Cantor set is described by the Cantor measure.

The Cantor function is continuous, and so by Exercise 5.5, the Cantor measure is a continuous probability measure. This is despite the fact that all of the mass of the Cantor measure is concentrated on the Cantor set, which is a set of Lebesgue measure zero! This makes the Cantor measure an example of what is called a *singular measure*, and as a result, even though we have given a reasonable probabilistic method for constructing Cantor-distributed random variables, the Cantor measure does not have a probability density function. That is, there is no function $f : [0, 1] \rightarrow [0, \infty)$ for which we can express $\mu_C((a, b]) = \int_a^b f(x) dx$. We will reencounter singular measures and deal with them systematically later in the course.

6. Regularity of Lebesgue–Stieltjes Measures

Built into the definition of Lebesgue–Stieltjes measures is the fact that they are locally finite, but it is not at all obvious that Lebesgue–Stieltjes measures should have additional regularity properties. However, from the outer measure construction, we can quickly deduce several useful and nontrivial properties about Lebesgue–Stieltjes measures.

PROPOSITION 5.31

Let μ be a Lebesgue–Stieltjes measure on \mathbb{R} , and let \mathcal{M}_μ be the σ -algebra of μ -measurable sets.

(1) Let $E \subseteq \mathbb{R}$. The following are equivalent:

- (i) $E \in \mathcal{M}_\mu$;
- (ii) for any $\varepsilon > 0$, there exists a closed set F and an open set G such that $F \subseteq E \subseteq G$ and $\mu(G \setminus F) < \varepsilon$;
- (iii) there exists an F_σ set A and a G_δ set B such that $A \subseteq E \subseteq B$ and $\mu(B \setminus A) = 0$.

(2) OUTER REGULARITY: If $E \in \mathcal{M}_\mu$, then

$$\mu(E) = \inf \{\mu(U) : U \supseteq E \text{ open}\}.$$

(3) INNER REGULARITY: If $E \in \mathcal{M}_\mu$, then

$$\mu(E) = \sup \{\mu(K) : K \subseteq E \text{ compact}\}.$$

PROOF. We will first prove outer regularity (2), then prove the measurability conditions (1), and end with inner regularity (3).

(2) Let $E \in \mathcal{M}_\mu$. By monotonicity of μ , it suffices to show $\mu(E) \geq \inf\{\mu(U) : U \supseteq E \text{ open}\}$. As usual, if $\mu(E) = \infty$, there is nothing to check, so assume $\mu(E) < \infty$, and let $\varepsilon > 0$. From the outer measure construction, we have

$$\mu(E) = \inf \left\{ \sum_{n=1}^{\infty} \mu((a_n, b_n]) : E \subseteq \bigcup_{n \in \mathbb{N}} (a_n, b_n] \right\}.$$

Hence, there exists a family of left-open, right-closed intervals $((a_n, b_n])_{n \in \mathbb{N}}$ such that $E \subseteq \bigcup_{n \in \mathbb{N}} (a_n, b_n]$ and $\sum_{n=1}^{\infty} \mu((a_n, b_n]) < \mu(E) + \frac{\varepsilon}{2}$. For each $n \in \mathbb{N}$, let $\delta_n > 0$ such that $\mu((a_n, b_n + \delta_n)) < \mu((a_n, b_n]) + 2^{-(n+1)}\varepsilon$. Such δ_n exists by continuity of μ from above. Then for the open set $U = \bigcup_{n \in \mathbb{N}} (a_n, b_n + \delta_n)$, we have

$$\mu(U) \leq \sum_{n=1}^{\infty} \mu((a_n, b_n + \delta_n)) < \sum_{n=1}^{\infty} (\mu((a_n, b_n]) + 2^{-(n+1)}\varepsilon) < \mu(E) + \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \mu(E) + \varepsilon.$$

But $\varepsilon > 0$ was arbitrary, so we are done.

(1) We will prove the chain of implications (i) \implies (ii) \implies (iii) \implies (i).

(i) \implies (ii). Suppose $E \in \mathcal{M}_\mu$, and let $\varepsilon > 0$. Let $E_n = E \cap (n, n+1]$ for $n \in \mathbb{Z}$. By (2), there exists an open set $G_n \supseteq E_n$ such that $\mu(G_n) < \mu(E_n) + 2^{-|n|}\frac{\varepsilon}{6}$. Let $G = \bigcup_{n \in \mathbb{Z}} G_n$. Then G is open, $E \subseteq G$, and

$$\mu(G \setminus E) \leq \sum_{n=-\infty}^{\infty} \mu(G_n \setminus E_n) < \frac{\varepsilon}{2}.$$

Applying the same argument to E^c , we find an open set $U \subseteq \mathbb{R}$ such that $E^c \subseteq U$ and $\mu(U \setminus E^c) < \frac{\varepsilon}{2}$. Let $F = U^c$. Then F is closed, $F \subseteq E$, and $\mu(E \setminus F) = \mu(U \setminus E^c) < \frac{\varepsilon}{2}$. Therefore, $\mu(G \setminus F) \leq \mu(G \setminus E) + \mu(E \setminus F) < \varepsilon$.

(ii) \implies (iii). For each $n \in \mathbb{N}$, choose $F_n \subseteq E \subseteq G_n$ such that $\mu(G_n \setminus F_n) < \frac{1}{n}$ by (ii). Let $A = \bigcup_{n \in \mathbb{N}} F_n$ and $B = \bigcap_{n \in \mathbb{N}} G_n$. Then A is an F_σ set, B is a G_δ set, $A \subseteq E \subseteq B$, and $\mu(B \setminus A) = 0$.

(iii) \implies (i). Suppose (iii) holds. Then we may write $E = A \cup N$, where A is an F_σ set and $N = E \setminus A \subseteq B \setminus A$. Since μ is complete and $\text{Borel}(\mathbb{R}) \subseteq \mathcal{M}_\mu$, we have $E \in \mathcal{M}_\mu$.

(3) Let $E \in \mathcal{M}_\mu$, and let $\varepsilon > 0$. Then by (1), there exists a closed set $F \subseteq E$ such that $\mu(E \setminus F) < \varepsilon$. Let $F_n = F \cap [-n, n]$ for $n \in \mathbb{N}$. Then F_n is compact, $F_1 \subseteq F_2 \subseteq \dots$, and $F = \bigcup_{n \in \mathbb{N}} F_n$. Therefore, by continuity of μ from below, $\mu(F_n) \rightarrow \mu(F)$ as $n \rightarrow \infty$. Hence, $\sup\{\mu(K) : K \subseteq E \text{ compact}\} \geq \mu(F) \geq \mu(E) - \varepsilon$. \square

Chapter Notes

The construction of Lebesgue–Stieltjes measures presented in this chapter is along the same lines as in the book of Folland [2, Sections 1.4 and 1.5]. A similar approach is taken in [9, Section 3.3] and [10, Sections 1.7.1–1.7.3].

Exercises

5.1 Confirm that the examples given in Example 5.7 are π -systems or λ -systems as claimed.

5.2 Let X be a set. Suppose \mathcal{P} is a π -system containing X . Let \mathcal{F} be a family of functions from X to \mathbb{R} with the following three properties

- (1) for every $E \in \mathcal{P}$, one has $\mathbb{1}_E \in \mathcal{F}$;
- (2) the space of functions \mathcal{F} is a real vector space: if $f, g \in \mathcal{F}$ and $c \in \mathbb{R}$, then $cf + g \in \mathcal{F}$;
- (3) if $0 \leq f_1 \leq f_2 \leq \dots$ is an increasing sequence in \mathcal{F} and $f = \lim_{n \rightarrow \infty} f_n$ is bounded, then $f \in \mathcal{F}$.

Prove that \mathcal{F} contains every bounded $\sigma(\mathcal{P})$ -measurable function.

5.3 A family of sets $\mathcal{C} \subseteq \mathcal{P}(X)$ is called a *monotone class* if it is closed under countable monotone unions and intersections. That is,

- if $(A_n)_{n \in \mathbb{N}}$ is a sequence in \mathcal{C} and $A_1 \subseteq A_2 \subseteq \dots$, then $\bigcup_{n \in \mathbb{N}} A_n \in \mathcal{C}$, and
- if $(B_n)_{n \in \mathbb{N}}$ is a sequence in \mathcal{C} and $B_1 \supseteq B_2 \supseteq \dots$, then $\bigcap_{n \in \mathbb{N}} B_n \in \mathcal{C}$.

The goal of this problem is to prove the monotone class theorem.

- (a) Show that a family of sets is a σ -algebra if and only if it is both an algebra and a monotone class.
- (b) Show that the intersection of a family of monotone classes is a monotone class.
- (c) Let \mathcal{C} be a monotone class. Show that $\mathcal{C}' = \{E \subseteq X : X \setminus E \in \mathcal{C}\}$ is also a monotone class.
- (d) Let \mathcal{C} be a monotone class. Show that $\mathcal{C}_E = \{F \subseteq X : E \cup F \in \mathcal{C}\}$ is also a monotone class for every $E \subseteq X$.

Let $\mathcal{A} \subseteq \mathcal{P}(X)$ be an algebra on X , and let $\mathcal{C}(\mathcal{A})$ be the monotone class generated by \mathcal{A} . (This monotone class is well-defined by part (b).)

- (e) Use part (c) to show that $\mathcal{C}(\mathcal{A})$ is closed under complementation.
- (f) Use part (d) to show that $\mathcal{C}(\mathcal{A})$ is closed under finite unions.
- (g) Deduce the monotone class theorem: $\mathcal{C}(\mathcal{A}) = \sigma(\mathcal{A})$.

5.4 Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $X : \Omega \rightarrow \mathbb{R}$ be a measurable function (called a *random variable* in probability theory). Prove that the cumulative distribution function $F_X : \mathbb{R} \rightarrow [0, 1]$ defined by $F(x) = \mathbb{P}(X \leq x)$ determines the measure induced by X , $\mathbb{P}_X : \text{Borel}(\mathbb{R}) \rightarrow [0, 1]$ defined by $\mathbb{P}_X(A) = \mathbb{P}(X^{-1}(A))$.

5.5 Let μ be a locally finite Borel measure on \mathbb{R} with distribution function F . For $x \in \mathbb{R}$, define $F(x^-) = \lim_{y \rightarrow x^-} F(y)$. The limit exists since F is an increasing function.

- (a) Show that $\mu(\{x\}) = F(x) - F(x^-)$.
- (b) Deduce that μ is a continuous measure if and only if F is a continuous function.

5.6 Let $\lambda : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ be the Lebesgue measure on the Borel subsets of \mathbb{R} . Let μ be another Borel measure on \mathbb{R} such that

- (1) for all $A \in \text{Borel}(\mathbb{R})$ and $x \in \mathbb{R}$, $\mu(A + x) = \mu(A)$, and

(2) $0 < \mu((0, 1]) < \infty$.

Show that there is a number $\alpha > 0$ such that $\mu = \alpha\lambda$.

5.7 Let λ be the Lebesgue measure on \mathbb{R} , and let E be a Lebesgue measurable set. Prove: if $\lambda(E) > 0$, then E contains a non-measurable set.

5.8 Let λ be the Lebesgue measure on \mathbb{R} .

- For a set $A \subseteq \mathbb{R}$, let $A - A$ be the set of differences $A - A = \{x - y : x, y \in A\}$. Show $t \in A - A$ if and only if $A \cap (A + t) \neq \emptyset$.
- Let A be a Lebesgue-measurable set with $\lambda(A) > 0$. Show that $A - A$ contains an interval around 0.
- Suppose $(H, +) \subseteq (\mathbb{R}, +)$ is a proper subgroup that is Lebesgue measurable. Prove $\lambda(H) = 0$.

5.9 Let μ be a Lebesgue–Stieltjes measure on \mathbb{R} with $\mu(\mathbb{R}) < \infty$. Let μ^* be the associated outer measure $\mu^*(E) = \inf \left\{ \sum_{n=1}^{\infty} \mu((a_n, b_n]) : E \subseteq \bigcup_{n \in \mathbb{N}} (a_n, b_n] \right\}$. Define the *inner measure* of a set $E \subseteq \mathbb{R}$ by $\mu_*(E) = \mu(\mathbb{R}) - \mu^*(\mathbb{R} \setminus E)$.

- Show that $\mu_*(E) = \sup \{\mu(K) : K \subseteq E \text{ compact}\}$.
- Prove that a set $E \subseteq \mathbb{R}$ is μ -measurable if and only if $\mu^*(E) = \mu_*(E)$.

CHAPTER 6

Borel Measures on Locally Compact Hausdorff Spaces

In the previous chapter, we constructed locally finite Borel measures on the real line. This is already sufficient for many purposes in probability theory, where the structure of the underlying measure space is often insignificant and the main object of study is (real-valued) random variables. However, in other contexts, the underlying structure of the measure space may play a prominent role (for example, if one is interested in measures on manifolds), and for this, we need additional tools to construct measures on more general topological spaces. The goal of this section is to construct Borel measures on locally compact Hausdorff spaces that are “compatible with the topology” in a sense that will be made precise below.

1. Locally Compact Hausdorff Spaces

DEFINITION 6.1

A topological space X is

- **Hausdorff** if every pair of points can be separated by open sets: if $x, y \in X$ and $x \neq y$, then there are open sets $U \ni x$ and $V \ni y$ such that $U \cap V = \emptyset$;
- **locally compact** if for every point has a compact neighborhood: for $x \in X$, there is an open set U and a compact set K such that $x \in U \subseteq K$.

If X is both locally compact and Hausdorff, we say X is a *locally compact Hausdorff space* or an *LCH space* for short.

EXAMPLE 6.2

Examples of locally compact Hausdorff spaces include:

- the unit interval $[0, 1]$
- the middle-thirds Cantor set
- Euclidean space \mathbb{R}^d for $d \in \mathbb{N}$
- topological manifolds
- discrete spaces

Non-examples include:

- the rational numbers \mathbb{Q} (not locally compact)
- infinite-dimensional real or complex vector spaces (not locally compact)
- an infinite set with the co-finite topology (not Hausdorff)

DEFINITION 6.3

Let X be an LCH space and $f : X \rightarrow \mathbb{C}$ a continuous function. The *support of f* is the set

$$\text{supp}(f) = \overline{\{f \neq 0\}}.$$

We say that f is *compactly supported* if $\text{supp}(f)$ is a compact subset of X .

NOTATION. We denote the space of compactly supported continuous functions on a topological space X by $C_c(X)$.

2. Radon Measures and the Riesz Representation Theorem

DEFINITION 6.4

Let X be an LCH space. A Borel measure μ on X is a *Radon measure* if μ is

- LOCALLY FINITE: $\mu(K) < \infty$ for every compact $K \subseteq X$;
- OUTER REGULAR: for every Borel set $E \subseteq X$,

$$\mu(E) = \inf\{\mu(U) : U \text{ is open and } E \subseteq U\};$$

and

- INNER REGULAR ON OPEN SETS: for every open set $G \subseteq X$,

$$\mu(G) = \sup\{\mu(K) : K \text{ is compact and } K \subseteq G\}.$$

Let X be an LCH space, and suppose μ is a Radon measure on X . Given $f \in C_c(X)$, we have

$$\int_X |f| d\mu \leq \sup_{x \in X} |f(x)| \cdot \mu(\text{supp}(f)).$$

The quantity $\sup_{x \in X} |f(x)|$ is actually a maximum and is finite by the extreme value theorem, while $\mu(\text{supp}(f)) < \infty$ since μ is locally finite. Hence, $C_c(X) \subseteq L^1(\mu)$. Integration against the measure μ thus induces a *positive linear functional* on $C_c(X)$.

DEFINITION 6.5

A *linear functional* on $C_c(X)$ is a linear map $\varphi : C_c(X) \rightarrow \mathbb{C}$. We say that a linear functional $\varphi : C_c(X) \rightarrow \mathbb{C}$ is *positive* if $\varphi(f) \geq 0$ for every $f \in C_c(X)$ with $f \geq 0$.

It turns out that all positive linear functionals on $C_c(X)$ arise via integration against a measure.

THEOREM 6.6: RIESZ REPRESENTATION THEOREM

Let X be a locally compact Hausdorff space. Given a positive linear functional $\varphi : C_c(X) \rightarrow \mathbb{C}$, there exists a unique Radon measure μ such that

$$\varphi(f) = \int_X f d\mu \tag{6.1}$$

for every $f \in C_c(X)$.

3. Topological Lemmas

LEMMA 6.7

Let X be an LCH space. Suppose $K \subseteq X$ is compact, $U \subseteq X$ is open, and $K \subseteq U$. Then there exists an open set $V \subseteq X$ such that \overline{V} is compact and $K \subseteq V \subseteq \overline{V} \subseteq U$.

PROOF. We first handle the case $K = \{x\}$. Since X is locally compact, there is an open neighborhood $W \subseteq X$ such that $x \in W$ and \overline{W} is compact. If $\overline{W} \subseteq U$, then we are done. Suppose $\overline{W} \not\subseteq U$. Then $L = \overline{W} \setminus U$ is a compact set, and $x \notin L$. For each point $y \in L$, let V_y and O_y be open sets such that $x \in V_y$, $y \in O_y$, and $V_y \cap O_y = \emptyset$. (The sets V_y and O_y

exist, since X is Hausdorff.) By compactness of L , there is a finite collection $y_1, \dots, y_n \in L$ such that $L \subseteq \bigcup_{j=1}^n O_{y_j}$. Let $V = W \cap \bigcap_{j=1}^n V_{y_j}$. Then V is open, $x \in V$, and $\overline{V} \subseteq \overline{W}$ is compact. Moreover, if $z \in \overline{V}$, then $z \in \overline{V}_{y_j}$ for every $j \in \{1, \dots, n\}$. Hence, $z \notin O_{y_j}$, so $z \notin L$. Therefore, $\overline{V} \subseteq \overline{W} \setminus L \subseteq U$.

Suppose now that K is an arbitrary compact set. By the above, we may find open sets V_x , $x \in K$, such that $x \in V_x \subseteq \overline{V}_x \subseteq U$. By compactness, there is a finite subcover $x_1, \dots, x_n \in K$ such that $K \subseteq \bigcup_{j=1}^n V_{x_j}$. We then take $V = \bigcup_{j=1}^n V_{x_j}$. \square

COROLLARY 6.8

Let X be an LCH space. Suppose $K \subseteq X$ is compact, $U_1, \dots, U_N \subseteq X$ are open, and $K \subseteq \bigcup_{n=1}^N U_n$. Then there exists open sets $V_n \subseteq X$ such that $\overline{V}_n \subseteq U_n$ is compact and $K \subseteq \bigcup_{n=1}^N V_n$.

PROOF. We do a proof by induction on N . The base case ($N = 1$) is Lemma 6.7. Suppose the statement holds for some $N \in \mathbb{N}$, and let K be a compact set and U_1, \dots, U_{N+1} an open cover of K . Let $K_1 = K \setminus U_{N+1}$. Then K_1 is a compact set covered by U_1, \dots, U_N , so by the inductive hypothesis, there exist open sets V_1, \dots, V_N such that $\overline{V}_n \subseteq U_n$ is compact for $n \leq N$ and $K_1 \subseteq \bigcup_{n=1}^N V_n$. Now let $K_2 = K \setminus (\bigcup_{n=1}^N V_n)$. Then K_2 is a compact subset of U_{N+1} , so by Lemma 6.7 there exists an open set V_{N+1} such that $\overline{V}_{N+1} \subseteq U_{N+1}$ is compact and $K_2 \subseteq V_{N+1}$. Then $K \subseteq \bigcup_{n=1}^{N+1} V_n$, so the corollary holds by induction. \square

LEMMA 6.9: URYSOHN'S LEMMA FOR LCH SPACES

Let X be an LCH space. Given a compact set $K \subseteq X$ and an open set $U \subseteq X$ with $K \subseteq U$, there exists a compactly supported continuous function $f : X \rightarrow [0, 1]$ such that $f = 1$ on K and $\text{supp}(f) \subseteq U$.

PROOF. Let K be a compact subset of X and $U \subseteq X$ an open set with $K \subseteq U$. By Lemma 6.7, let V be an open set such that \overline{V} is compact and $K \subseteq V \subseteq \overline{V} \subseteq U$. We construct a function f supported on \overline{V} in terms of its sub-level sets.

Let $K(1) = K$, $V(0) = V$. Put $V(1) = \emptyset$ and $K(0) = \overline{V}$.

CLAIM 1. There are families of open sets $V(r)$ and compact sets $K(r)$ indexed by dyadic rationals $r \in [0, 1]$ such that

- for every dyadic rational $r \in [0, 1]$, $V(r) \subseteq K(r)$, and
- for dyadic rationals $r, s \in [0, 1]$, if $r > s$, then $K(r) \subseteq V(s)$.

We will prove the claim by induction on the denominators of dyadic rationals. By Lemma 6.7, let $V(1/2)$ be an open set such that $K(1/2) = \overline{V(1/2)}$ is compact and $K = K(1) \subseteq V(1/2) \subseteq K(1/2) \subseteq V(0) = V$.

Suppose we have constructed sets $V(r)$ and $K(r)$ with the desired properties for dyadic rationals $r \in (0, 1)$ with denominators 2^n for $n < N$. Let $r \in (0, 1)$ by a dyadic rational with denominator 2^N , say $r = \frac{2j-1}{2^N}$, $j \in \{1, \dots, 2^{N-1}\}$. By the induction hypothesis, we have a compact set $K\left(\frac{j}{2^{N-1}}\right)$ and an open set $V\left(\frac{j-1}{2^{N-1}}\right)$ such that $K\left(\frac{j}{2^{N-1}}\right) \subseteq V\left(\frac{j-1}{2^{N-1}}\right)$. Applying Lemma 6.7, we then obtain an open set $V(r)$ such that $K(r) = \overline{U(r)}$ is compact, and $K\left(\frac{j}{2^{N-1}}\right) \subseteq V(r) \subseteq K(r) \subseteq V\left(\frac{j-1}{2^{N-1}}\right)$. The claim thus holds by induction.

Define $f(x) = 0$ if $x \notin V(0)$ and $f(x) = \sup\{r \geq 0 : x \in V(r)\}$ otherwise. By construction $K = K(1) \subseteq V(r)$ for every dyadic rational $r \in [0, 1]$, so $f = 1$ on K . Moreover, $\text{supp}(f) \subseteq K(0) \subseteq U$. It remains to show that f is continuous.

CLAIM 2. For $a \in \mathbb{R}$,

$$\{f > a\} = \begin{cases} \emptyset, & \text{if } a \geq 1; \\ \bigcup_{r > a} V(r), & \text{if } 0 \leq a < 1; \\ X, & \text{if } a < 0. \end{cases}$$

The function f takes values between 0 and 1, so the cases $a < 0$ and $a \geq 1$ are immediate. Let $0 \leq a < 1$. Suppose $x \in X$ and $f(x) > a$. Then by the definition of f , there exists $r > a$ such that $x \in V(r)$. Hence, $x \in \bigcup_{r > a} V(r)$. Conversely, if $x \in \bigcup_{r > a} V(r)$, then $f(x) \geq r > a$.

CLAIM 3. For $b \in \mathbb{R}$,

$$\{f < b\} = \begin{cases} X, & \text{if } b > 1; \\ \bigcup_{r < b} (X \setminus K(r)), & \text{if } 0 < b \leq 1; \\ \emptyset, & \text{if } b \leq 0. \end{cases}$$

As in the previous claim, since f takes values in the interval $[0, 1]$, the cases $b > 1$ and $b \leq 0$ are immediate. Let $0 < b \leq 1$. Suppose $f(x) < b$. Taking $s \in (f(x), b)$, we have $x \notin V(s)$. Let $r \in (s, b)$. Then since $K(r) \subseteq V(s)$, we conclude $x \notin K(r)$. Hence, $x \in \bigcup_{r < b} (X \setminus K(r))$. Conversely, if $x \notin K(r)$ for some $r < b$, then $x \notin V(s)$ for $s \geq r$, so $f(x) \leq r < b$.

Combining Claims 2 and 3, for any $a, b \in \mathbb{R}$, the set $\{a < f < b\}$ is an intersection of two open sets and therefore open. Thus, f is continuous. \square

NOTATION. For a compact set $K \subseteq X$ and a function $f : X \rightarrow [0, 1]$, we write $K \prec f$ if $f = 1$ on K . Given an open set $U \subseteq X$ and a function $f : X \rightarrow [0, 1]$, we write $f \prec U$ if $\text{supp}(f) \subseteq U$. With this notation, the conclusion of Urysohn's lemma reads $K \prec f \prec U$.

COROLLARY 6.10: PARTITION OF UNITY

Let X be an LCH space. Let $K \subseteq X$ be a compact set and $U_1, \dots, U_N \subseteq X$ an open cover of K . Then there exist compactly supported continuous functions $h_n : X \rightarrow [0, 1]$ such that $h_n \prec U_n$ for each $n \in \{1, \dots, N\}$ and $\sum_{n=1}^N h_n = 1$ on K .

PROOF. First apply Corollary 6.8 to obtain open sets V_1, \dots, V_N such that $\overline{V}_n \subseteq U_n$ is compact and $K \subseteq \bigcup_{n=1}^N V_n$. Then by Urysohn's lemma, let $f_n \in C_c(X)$ with $\overline{V}_n \prec f_n \prec U_n$. Define $h_1 = f_1, h_2 = (1 - f_1)f_2, \dots, h_N = (1 - f_1) \dots (1 - f_{N-1})f_N$. For each $n \in \{1, \dots, N\}$, $h_n \leq f_n$, so h_n has compact support, and $h_n \prec U_n$. Moreover, it can be checked by induction on N that

$$\sum_{n=1}^N h_n = 1 - \prod_{n=1}^N (1 - f_n).$$

For $x \in K$, at least one of the functions $f_n(x)$ is equal to 1, so $\sum_{n=1}^N h_n(x) = 1$. \square

LEMMA 6.11

Let X be a locally compact Hausdorff space, and suppose μ is a Radon measure on X . Then for any open set $U \subseteq X$,

$$\mu(U) = \sup \left\{ \int_X f \, d\mu : f \in C_c(X), 0 \leq f \prec U \right\}.$$

PROOF. Clearly $\mu(U) \geq \int_X f \, d\mu$ for any $f \in C_c(x)$ with $0 \leq f \leq \mathbb{1}_U$. Let us prove the reverse inequality. Let $c < \mu(U)$ be arbitrary. Then by inner regularity of μ on open sets, there exists a compact set $K \subseteq U$ such that $\mu(K) > c$. Then by Urysohn's lemma, there is a continuous function $f \in C_c(X)$ such that $K \prec f \prec U$. By monotonicity of the integral, we then have

$$\int_X f \, d\mu \geq \mu(K) > c.$$

\square

4. Proof of the Riesz Representation Theorem

PROOF OF RIESZ REPRESENTATION THEOREM. We will carry out the proof in several steps.

STEP 1. Uniqueness.

Suppose μ and ν are two Radon measures satisfying (6.1). By Lemma 6.11, μ and ν must agree on all open subsets of X . But then by outer regularity, μ and ν agree on all Borel sets.

STEP 2. Defining an Outer Measure.

Motivated by Lemma 6.11, we define

$$m(U) = \sup \{ \varphi(f) : f \in C_c(X), 0 \leq f \prec U \}$$

for open subsets $U \subseteq X$, and let

$$\mu^*(E) = \inf \{ m(U) : U \text{ is open and } E \subseteq U \}$$

for $E \subseteq X$.

We must check that μ^* is an outer measure. That $\mu^*(\emptyset) = 0$ and μ^* is monotone are both easy consequences of the definition of μ^* . Suppose $(E_n)_{n \in \mathbb{N}}$ is a countable family of subsets of X . We want to show

$$\mu^* \left(\bigcup_{n \in \mathbb{N}} E_n \right) \leq \sum_{n=1}^{\infty} \mu^*(E_n).$$

If the sum diverges, there is nothing to show, so assume the sum is finite. Let $\varepsilon > 0$ be arbitrary. Take $(U_n)_{n \in \mathbb{N}}$ open sets such that $E_n \subseteq U_n$ and $\mu^*(E_n) > m(U_n) - 2^{-n}\varepsilon$. Let $f \in C_c(X)$ with $0 \leq f \prec \bigcup_{n \in \mathbb{N}} U_n$. Since the sets $(U_n)_{n \in \mathbb{N}}$ are open and $\text{supp}(f)$ is a compact set, there exists $N \in \mathbb{N}$ such that $\text{supp}(f) \subseteq \bigcup_{n=1}^N U_n$. By partition of unity (Corollary 6.10), let $h_1, \dots, h_N \in C_c(X)$ such that $h_n \prec U_n$ and $\sum_{n=1}^N h_n = 1$ on $\text{supp}(f)$. Letting $f_n = f \cdot h_n$, we have $f = \sum_{n=1}^N f_n$. Therefore,

$$\varphi(f) = \sum_{n=1}^N \varphi(f_n) \leq \sum_{n=1}^N m(U_n) \leq \sum_{n=1}^{\infty} m(U_n) \leq \sum_{n=1}^{\infty} \mu^*(E_n) + \varepsilon.$$

Taking a supremum over all such f , we conclude

$$\mu^* \left(\bigcup_{n \in \mathbb{N}} E_n \right) \leq m \left(\bigcup_{n \in \mathbb{N}} U_n \right) = \sup \left\{ \varphi(f) : f \in C_c(X), 0 \leq f \prec \bigcup_{n \in \mathbb{N}} U_n \right\} \leq \sum_{n=1}^{\infty} \mu^*(E_n) + \varepsilon.$$

Since ε was arbitrary, this proves that μ^* is countably subadditive and therefore an outer measure.

STEP 3. Borel Sets are μ^* -Measurable.

By Carathéodory's theorem (Theorem 5.24), the family of μ^* -measurable sets is a σ -algebra, so it suffices to check that every open set is μ^* -measurable. Let $U \subseteq X$ be an open set. We want to show

$$\mu^*(E) \geq \mu^*(E \cap U) + \mu^*(E \setminus U) \quad (6.2)$$

for every $E \subseteq X$ with $\mu^*(E) < \infty$. Let $E \subseteq X$ be any set with $\mu^*(E) < \infty$. Let $\varepsilon > 0$ be arbitrary, and choose an open set $V \subseteq X$ with $E \subseteq V$ and $\mu^*(E) > m(V) - \varepsilon$. The set $V \cap U$ is open, so choose a function $f_1 \in C_c(X)$ with $0 \leq f_1 \prec V \cap U$ such that $\varphi(f_1) > m(V \cap U) - \varepsilon$. Let $K = \text{supp}(f_1)$, and then choose a function $f_2 \in C_c(X)$ with $0 \leq f_2 \prec V \setminus K$ such that $\varphi(f_2) > m(V \setminus K) - \varepsilon$. Then

$$\mu^*(E) > m(V) - \varepsilon \geq \varphi(f_1 + f_2) - \varepsilon > m(V \cap U) + m(V \setminus K) - 3\varepsilon \geq \mu^*(E \cap U) + \mu^*(E \setminus U) - 3\varepsilon.$$

Taking $\varepsilon \rightarrow 0$ gives (6.2).

We can now define a Borel measure μ by $\mu = \mu^*|_{\text{Borel}(X)}$.

STEP 4. Regularity of μ .

The measure μ is outer regular by construction. We will check that it is also inner regular on open sets. Let $U \subseteq X$ be open, and let $c < \mu(U)$. By the definition of μ , there exists a function $f \in C_c(X)$ with $0 \leq f \prec U$ such that $\varphi(f) > c$. Let $K = \text{supp}(f)$. We claim $\mu(K) \geq \varphi(f) > c$. From the definition of μ , it suffices to show: if $V \supseteq K$ is open, then there exists a function $g \in C_c(X)$ with $0 \leq g \prec V$ with $\varphi(g) > c$. But this is immediate upon taking $g = f$.

STEP 5. Local Finiteness.

Let $K \subseteq X$ be compact. We want to show $\mu(K) < \infty$. It suffices to find an open set $U \supseteq K$ with $m(U) < \infty$. Since X is locally compact, there exists an open set $U \supseteq K$ such that \overline{U} is compact (see Lemma 6.7). Let $f \in C_c(X)$ with $\overline{U} \prec f$. Suppose $g \in C_c(X)$ with $0 \leq g \prec U$. Then $f - g \geq 0$, so $\varphi(g) \leq \varphi(f)$. Thus,

$$m(U) = \sup\{\varphi(g) : g \in C_c(x), 0 \leq g \leq 1, \text{supp}(g) \subseteq U\} \leq \varphi(f) < \infty.$$

STEP 6. Proving the Identity (6.1).

Let $f \in C_c(X)$, and let $K = \text{supp}(f)$. By decomposing f into real and imaginary parts, then positive and negative parts and scaling, we may assume $0 \leq f \leq 1$. Given $N \in \mathbb{N}$, we decompose $K = \bigsqcup_{n=0}^N K_n$, where $K_0 = \{x \in K : f(x) = 0\}$ and $K_n = \{x \in K : f(x) \in (\frac{n-1}{N}, \frac{n}{N}]\}$ for $n \geq 1$. For $n \in \{1, \dots, N\}$, let

$$f_n(x) = \begin{cases} 0, & \text{if } x \in K_m, m < n \\ f(x) - \frac{n-1}{N}, & \text{if } x \in K_n \\ \frac{1}{N}, & \text{if } x \in K_m, m > n. \end{cases}$$

Then $f_n \in C_c(X)$ and $f = \sum_{n=1}^N f_n$. Moreover, $\bigsqcup_{m>n} \overline{K}_m \prec N f_n \prec \bigsqcup_{m \geq n} K_m$. We can therefore estimate $\varphi(f)$ and $\int_X f \, d\mu$ as follows:

$$\frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m>n} K_m\right) \leq \varphi(f) \leq \frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m \geq n} K_m\right).$$

and

$$\frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m>n} K_m\right) \leq \int_X f \, d\mu \leq \frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m \geq n} K_m\right).$$

All that remains is to check that the two sides of the inequality become arbitrarily close as $N \rightarrow \infty$. Observe:

$$\frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m \geq n} K_m\right) - \frac{1}{N} \sum_{n=1}^N \mu\left(\bigsqcup_{m>n} K_m\right) = \frac{1}{N} \sum_{n=1}^N \mu(K_n) = \frac{1}{N} \mu(K),$$

and $\mu(K) < \infty$ by Step 5. Therefore, taking $N \rightarrow \infty$ and applying the squeeze theorem, we conclude

$$\int_X f \, d\mu = \varphi(f).$$

□

EXAMPLE 6.12

Let $R : C_c(\mathbb{R}) \rightarrow \mathbb{C}$ be the functional induced by Riemann integration. That is, if $f : \mathbb{R} \rightarrow \mathbb{C}$ with compact support, say $\text{supp}(f) \subseteq [a, b]$, then $R(f) = \int_a^b f(x) \, dx$. The measure representing the functional R is the Lebesgue measure on \mathbb{R} .

Exercises

6.1 Let X be an LCH space, and let $\mu : \text{Borel}(X) \rightarrow [0, \infty]$ be a Borel measure on X . Prove the following are equivalent:

- (i) $\mu(K) < \infty$ for all $K \subseteq X$ compact;
- (ii) for every $x \in X$, there exists an open set $U \subseteq X$ with $x \in U$ such that $\mu(U) < \infty$.

6.2 (a) Show that the Dirac functional $\delta_0 : C([0, 1]) \rightarrow \mathbb{C}$ defined by $\delta_0(f) = f(0)$ is not of the form

$$\delta_0(f) = \int_0^1 f(t)g(t) \, dt$$

for a function $g \in C([0, 1])$.

(b) Define a positive linear functional $\psi : C([0, 1]) \rightarrow \mathbb{C}$ by

$$\psi(f) = \frac{f(0) + f(1)}{2} + \int_0^1 tf(t) \, dt.$$

Determine the measure μ corresponding to ψ , and calculate $\mu([0, 1])$.

6.3 In this exercise, we will construct a Haar measure on the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. For this, recall that one can identify functions $f : \mathbb{T} \rightarrow \mathbb{C}$ with 1-periodic functions $F : \mathbb{R} \rightarrow \mathbb{C}$ (i.e. we require $F(x + 1) = F(x)$). Furthermore, f is continuous (measurable) if and only if F is continuous (measurable). We define a measure m on \mathbb{T} by requiring that

$$\int_{\mathbb{T}} f \, dm = \int_0^1 F(x) \, dx$$

where f and F are continuous functions corresponding to each other. Justify that m is well-defined and show that m is a Haar measure on \mathbb{T} .

6.4 This exercise shows that the locally finite assumption cannot be removed in Theorem 9.5. We consider the (Alexandrov) one-point compactification of \mathbb{N} , i.e. the topological space (X, τ) with $X = \mathbb{N} \cup \{\infty\}$ and $\tau = \{U \subseteq X : U \subseteq \mathbb{N} \text{ or } U^c \text{ is finite}\}$.

- (a) Show that (X, τ) is a compact Hausdorff space in which every subset is σ -compact. Check that the Borel σ -algebra is $\text{Borel}(X) = \mathcal{P}(X)$.
- (b) Let $\mu : \mathcal{P}(X) \rightarrow [0, \infty]$ be the counting measure. Prove that μ is inner regular but not outer regular.

CHAPTER 7

Products of Measure Spaces

DEFINITION 7.1

Let (X, \mathcal{B}) and (Y, \mathcal{C}) be measurable spaces. A *measurable rectangle* in $X \times Y$ is a set of the form $B \times C$ such that $B \in \mathcal{B}$ and $C \in \mathcal{C}$. The *product σ -algebra* $\mathcal{B} \otimes \mathcal{C}$ on $X \times Y$ is the σ -algebra generated by measurable rectangles.

REMARK. The product can also be defined in a category-theoretic way. Let $\pi_X : X \times Y$ be the projection onto the first coordinate, $\pi_X(x, y) = x$, and let $\pi_Y : X \times Y$ be the projection onto the second coordinate, $\pi_Y(x, y) = y$. The maps π_X and π_Y are easily checked to be measurable maps defined on $(X \times Y, \mathcal{B} \otimes \mathcal{C})$. The product measurable space $(X \times Y, \mathcal{B} \otimes \mathcal{C})$ satisfies the following universal property (see Figure 7.1): for any measurable space (Z, \mathcal{D}) and any measurable functions $f : Z \rightarrow X$ and $g : Z \rightarrow Y$, there is a unique measurable function $h : Z \rightarrow X \times Y$ such that $\pi_X \circ h = f$ and $\pi_Y \circ h = g$. This universal property characterizes the product space $(X \times Y, \mathcal{B} \otimes \mathcal{C})$ uniquely up to isomorphism.

$$\begin{array}{ccccc} & & Z & & \\ & \swarrow f & \downarrow h & \searrow g & \\ X & \xleftarrow{\pi_X} & X \times Y & \xrightarrow{\pi_Y} & Y \end{array}$$

FIGURE 7.1. Universal property of product spaces.

DEFINITION 7.2

Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be measure spaces. A measure $\rho : \mathcal{B} \otimes \mathcal{C} \rightarrow [0, \infty]$ is a *product measure* of μ and ν if $\rho(B \times C) = \mu(B)\nu(C)$ for every $B \in \mathcal{B}$ and $C \in \mathcal{C}$.

THEOREM 7.3

Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be measure spaces. There exists a product measure $\rho : \mathcal{B} \otimes \mathcal{C} \rightarrow [0, \infty]$. Moreover, if μ and ν are σ -finite, then there is a unique product measure.

The uniqueness part of Theorem 7.3 follows by the π - λ theorem, so we present its proof first.

PROOF OF UNIQUENESS OF PRODUCT MEASURE FOR σ -FINITE SPACES. Suppose (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) are σ -finite measure spaces, and suppose $\rho_1, \rho_2 : \mathcal{B} \otimes \mathcal{C} \rightarrow [0, \infty]$ are product measures of μ and ν . We want to show that $\rho_1 = \rho_2$. We will use the π - λ theorem.

The family \mathcal{R} of measurable rectangles is a π -system (see Example 5.7 and Exercise 5.1).

Since μ and ν are σ -finite measures, we may write $X = \bigcup_{n \in \mathbb{N}} X_n$ with $X_1 \subseteq X_2 \subseteq \dots$ such that $\mu(X_n) < \infty$ and $Y = \bigcup_{n \in \mathbb{N}} Y_n$ with $Y_1 \subseteq Y_2 \subseteq \dots$ such that $\nu(Y_n) < \infty$. Define a family

$$\mathcal{L} = \{E \in \mathcal{B} \otimes \mathcal{C} : \rho_1(E \cap (X_n \times Y_n)) = \rho_2(E \cap (X_n \times Y_n)) \text{ for every } n \in \mathbb{N}\}.$$

By the same argument as in the proof of Corollary 5.14, \mathcal{L} is a λ -system on $X \times Y$.

Moreover, $\mathcal{R} \subseteq \mathcal{L}$. Indeed, if $E = B \times C$ is a measurable rectangle, then $E \cap (X_n \times Y_n) = (B \cap X_n) \times (C \cap Y_n)$ is also a measurable rectangle for every $n \in \mathbb{N}$, so

$$\rho_1(E \cap (X_n \times Y_n)) = \mu(B \cap X_n)\nu(C \cap Y_n) = \rho_2(E \cap (X_n \times Y_n))$$

by the definition of a product measure.

Thus, by the π - λ theorem, $\mathcal{L} \supseteq \sigma(\mathcal{R}) = \mathcal{B} \otimes \mathcal{C}$. Applying continuity from below of the measures ρ_1 and ρ_2 , given an arbitrary measurable set $E \in \mathcal{B} \otimes \mathcal{C}$, we have

$$\rho_1(E) = \lim_{n \rightarrow \infty} \rho_1(E \cap (X_n \times Y_n)) = \lim_{n \rightarrow \infty} \rho_2(E \cap (X_n \times Y_n)) = \rho_2(E).$$

That is, $\rho_1 = \rho_2$. □

The preceding proof shows that it makes sense to talk about *the* product measure of a pair of σ -finite measures.

For the existence part of Theorem 7.3, several different constructions of product measures are possible, and in the case of non- σ -finite spaces, different constructions may produce different measures.

1. Cross-Sectional Product Measures and the Fubini–Tonelli Theorem

DEFINITION 7.4

Let X and Y be sets, and let $(x, y) \in X \times Y$.

- for a set $E \subseteq X \times Y$, the *x-section* E_x and the *y-section* E^y of E are defined by

$$E_x = \{v \in Y : (x, v) \in E\} \quad \text{and} \quad E^y = \{u \in X : (u, y) \in E\}.$$

- for a function f defined on $X \times Y$, the *x-section* f_x and the *y-section* f^y of f are defined by

$$f_x(v) = f(x, v) \quad \text{and} \quad f^y(u) = f(u, y).$$

REMARK. If $E \subseteq X \times Y$, then we have the identities $(\mathbb{1}_E)_x = \mathbb{1}_{E_x}$ and $(\mathbb{1}_E)^y = \mathbb{1}_{E^y}$.

PROPOSITION 7.5

Let (X, \mathcal{B}) and (Y, \mathcal{C}) be measurable spaces.

- (1) If $E \in \mathcal{B} \otimes \mathcal{C}$, then $E_x \in \mathcal{C}$ for every $x \in X$ and $E^y \in \mathcal{B}$ for every $y \in Y$.
- (2) If f is a $(\mathcal{B} \otimes \mathcal{C})$ -measurable function on $X \times Y$, then f_x is \mathcal{C} -measurable for every $x \in X$ and f^y is \mathcal{B} -measurable for every $y \in Y$.

PROOF. (1) Consider the family

$$\mathcal{F} = \{E \subseteq X \times Y : E_x \in \mathcal{C} \text{ for every } x \in X \text{ and } E^y \in \mathcal{B} \text{ for every } y \in Y\}.$$

Then \mathcal{F} contains all measurable rectangles, since

$$(B \times C)_x = \begin{cases} C, & \text{if } x \in B; \\ \emptyset, & \text{if } x \notin B, \end{cases} \quad \text{and} \quad (B \times C)^y = \begin{cases} B, & \text{if } y \in C; \\ \emptyset, & \text{if } y \notin C. \end{cases} \quad (7.1)$$

Moreover, since taking cross-sections is compatible with (countable) unions and complements, \mathcal{F} is a σ -algebra. Hence, $\mathcal{B} \otimes \mathcal{C} \subseteq \mathcal{F}$, which proves (1).

(2) This follows from (1) by noting that pre-images are compatible with cross-sections in the sense that $(f_x)^{-1}(E) = (f^{-1}(E))_x$ and $(f^y)^{-1}(E) = (f^{-1}(E))^y$. \square

THEOREM 7.6

Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be measure spaces.

(1) If ν is s-finite, then the map $x \mapsto \nu(E_x)$ is measurable, and $\rho : \mathcal{B} \otimes \mathcal{C} \rightarrow [0, \infty]$ defined by

$$\rho(E) = \int_X \nu(E_x) d\mu(x)$$

is a product measure of μ and ν .

(2) If μ and ν are both s-finite, then

$$\int_X \nu(E_x) d\mu(x) = \int_Y \mu(E^y) d\nu(y)$$

for every $E \in \mathcal{B} \otimes \mathcal{C}$.

PROOF. (1) Suppose ν is s-finite.

CLAIM 1. The map $x \mapsto \nu(E_x)$ is measurable.

We can write $\nu = \sum_{n=1}^{\infty} \nu_n$ for some finite measures $\nu_n : \mathcal{C} \rightarrow [0, \infty)$. Since a countable sum of measurable functions is measurable, it suffices to prove measurability under the stronger hypothesis that ν is finite.

Consider the family of sets

$$\mathcal{L} = \{E \subseteq X \times Y : x \mapsto \nu(E_x) \text{ is measurable}\}.$$

Using (7.1), we see that \mathcal{L} contains the π -system of measurable rectangles. By the π - λ theorem, it therefore suffices to prove that \mathcal{L} is a λ -system.

The set $X \times Y$ is a measurable rectangle so belongs to \mathcal{L} .

Suppose $E \in \mathcal{L}$. Noting that $((X \times Y) \setminus E)_x = Y \setminus E_x$, we have that

$$\nu(((X \times Y) \setminus E)_x) = \nu(Y) - \nu(E_x)$$

is measurable, so $(X \times Y) \setminus E \in \mathcal{L}$.

Finally, if $(E_n)_{n \in \mathbb{N}}$ is a sequence of pairwise disjoint elements of \mathcal{L} and $E = \bigsqcup_{n \in \mathbb{N}} E_n$, then

$$\nu(E_x) = \sum_{n=1}^{\infty} \nu((E_n)_x)$$

is measurable, so $E \in \mathcal{L}$.

Measurability of $x \mapsto \nu(E_x)$ means that ρ is a well-defined function.

CLAIM 2. ρ is a measure on $(X \times Y, \mathcal{B} \otimes \mathcal{C})$.

Since $\emptyset_x = \emptyset$ for every $x \in X$, we have

$$\rho(\emptyset) = \int_X \nu(\emptyset) \, d\mu = \int_X 0 \, d\mu = 0 \cdot \mu(X) = 0.$$

Suppose $(E_n)_{n \in \mathbb{N}}$ is a sequence of pairwise disjoint measurable subsets of $X \times Y$. Then

$$\begin{aligned} \rho\left(\bigsqcup_{n \in \mathbb{N}} E_n\right) &= \int_X \nu\left(\bigsqcup_{n \in \mathbb{N}} (E_n)_x\right) \, d\mu(x) \\ &= \int_X \sum_{n=1}^{\infty} \nu((E_n)_x) \, d\mu(x) \\ &\stackrel{(*)}{=} \sum_{n=1}^{\infty} \int_X \nu((E_n)_x) \, d\mu(x) \\ &= \sum_{n=1}^{\infty} \rho(E_n). \end{aligned}$$

In step (*), we used Theorem 3.12.

CLAIM 3. ρ is a product measure of μ and ν .

Let $B \in \mathcal{B}$ and $C \in \mathcal{C}$. As noted previously (see (7.1)), $(B \otimes C)_x = C$ if $x \in B$ and $(B \otimes C)_x = \emptyset$ if $x \notin B$. Hence, the function $x \mapsto \nu((B \otimes C)_x)$ is a simple function, and integrating with respect to μ gives

$$\rho(B \times C) = \int_X \nu((B \otimes C)_x) \, d\mu(x) = \nu(C) \cdot \mu(B) + \nu(\emptyset) \cdot \mu(X \setminus B) = \mu(B)\nu(C).$$

(2) Now suppose μ and ν are s -finite. Let

$$\rho_1(E) = \int_X \nu(E_x) \, d\mu(x) \quad \text{and} \quad \rho_2(E) = \int_Y \mu(E^y) \, d\nu(y).$$

CLAIM 3. $\rho_1 = \rho_2$

Write $\mu = \sum_{n=1}^{\infty} \mu_n$ and $\nu = \sum_{n=1}^{\infty} \nu_n$ for some finite measures μ_n, ν_n . Then by Theorem 3.12,

$$\rho_1(E) = \sum_{m,n} \int_X \nu_n(E_x) \, d\mu_m(x) \quad \text{and} \quad \rho_2(E) = \sum_{m,n} \int_Y \mu_m(E^y) \, d\nu_n(y). \quad (7.2)$$

For each $m, n \in \mathbb{N}$, the measures

$$\rho_{1,m,n}(E) = \int_X \nu_n(E_x) \, d\mu_m(x) \quad \text{and} \quad \rho_{2,m,n}(E) = \int_Y \mu_m(E^y) \, d\nu_n(y)$$

are product measures of μ_m and ν_n (by Claims 1 and 2). But the product of (σ) -finite measures is unique, so $\rho_{1,m,n} = \rho_{2,m,n}$. Hence, by (7.2), $\rho_1 = \rho_2$.

□

DEFINITION 7.7

Given s-finite measure spaces (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) , we call the product measure obtained by Theorem 7.6 the *cross-sectional product measure* and denote it by $\mu \overset{cs}{\otimes} \nu$.

Theorem 7.6 extends to a result about integration of measurable functions on products of s-finite measure spaces.

THEOREM 7.8: FUBINI–TONELLI THEOREM

Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be s-finite measure spaces.

(1) (Tonelli) Let $f : X \times Y \rightarrow [0, \infty]$ be a $(\mathcal{B} \otimes \mathcal{C})$ -measurable function. Then $x \mapsto \int_Y f_x d\nu$ and $y \mapsto \int_X f^y d\mu$ are measurable functions, and

$$\int_{X \times Y} f d(\mu \overset{cs}{\otimes} \nu) = \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x) = \int_Y \left(\int_X f(x, y) d\mu(x) \right) d\nu(y). \quad (7.3)$$

(2) (Fubini) Suppose $f \in L^1(\mu \overset{cs}{\otimes} \nu)$. Then $f_x \in L^1(\nu)$ for μ -a.e. $x \in X$, $f_y \in L^1(\mu)$ for ν -a.e. $y \in Y$, the almost-everywhere defined functions $x \mapsto \int_Y f_x d\nu$ and $y \mapsto \int_X f^y d\mu$ belong to $L^1(\mu)$ and $L^1(\nu)$ respectively, and (7.3) holds.

PROOF. (1) If $f = \mathbb{1}_E$ for some $E \in \mathcal{B} \otimes \mathcal{C}$, then (1) holds by Theorem 7.6. Hence, (1) holds for simple functions. For general f , let $(f_n)_{n \in \mathbb{N}}$ be a sequence of simple functions such that $0 \leq f_1 \leq f_2 \leq \dots$ and $f_n \rightarrow f$ pointwise as in Proposition 3.7. Then $(f_n)_x$ increases to f_x and $(f_n)^y$ increases to f^y , so (7.3) holds by repeated application of the monotone convergence theorem.

(2) Let $f \in L^1(\mu \overset{cs}{\otimes} \nu)$. By (1),

$$\int_{X \times Y} |f| d(\mu \overset{cs}{\otimes} \nu) = \int_X \left(\int_Y |f_x| d\nu \right) d\mu(x) = \int_Y \left(\int_X |f^y| d\mu \right) d\nu(y).$$

This integral is finite, so $\int_Y |f_x| d\nu < \infty$ for μ -a.e. $x \in X$ and $\int_X |f^y| d\mu < \infty$ for ν -a.e. $y \in Y$ by Proposition 3.20. That is, $f_x \in L^1(\nu)$ for μ -a.e. $x \in X$, and $f^y \in L^1(\mu)$ for ν -a.e. $y \in Y$. Moreover, by the triangle inequality for integrals,

$$\int_X \left| \int_Y f_x d\nu \right| d\mu(x) \leq \int_X \left(\int_Y |f_x| d\nu \right) d\mu(x) < \infty$$

and similarly for the iterated integral in the other order.

The identity (7.3) holds for the positive and negative parts of the real and imaginary parts of f by (1), and these can be recombined to conclude (7.3) for the function f itself. \square

EXAMPLE 7.9

Let λ be the Lebesgue measure on \mathbb{R} . For an integrable function f , the *Fourier transform* of f is the function $\widehat{f} : \mathbb{R} \rightarrow \mathbb{R}$ defined by $\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i \xi x} dx$. Given integrable functions f and g , we define the *convolution* $(f * g)(x) = \int_{\mathbb{R}} f(x-y) g(y) dy$. Then $f * g$ is well-defined almost everywhere, integrable, and $\widehat{(f * g)}(\xi) = \widehat{f}(\xi) \widehat{g}(\xi)$.

To see this, consider the function $\Phi(x, y) = f(x - y)g(y)$. Since f and g are measurable, Φ is also measurable. Moreover, by Tonelli's theorem,

$$\int_{\mathbb{R}^2} |\Phi(x, y)| d(x, y) = \int_{\mathbb{R}} |g(y)| \left(\int_{\mathbb{R}} |f(x - y)| dx \right) dy = \left(\int_{\mathbb{R}} |f| d\lambda \right) \left(\int_{\mathbb{R}} |g| d\lambda \right) < \infty.$$

Therefore, by Fubini's theorem, $f * g$ is almost everywhere well-defined, integrable, and satisfies

$$\begin{aligned} \widehat{(f * g)}(\xi) &= \int_{\mathbb{R}} (f * g)(x) e^{-2\pi i \xi x} dx \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} f(x - y)g(y) e^{-2\pi i \xi x} dy dx \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} f(x - y)g(y) e^{-2\pi i \xi x} dx dy \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} f(t)g(y) e^{-2\pi i \xi(t+y)} dt dy \\ &= \int_{\mathbb{R}} f(t) e^{-2\pi i \xi t} dt \int_{\mathbb{R}} g(y) e^{-2\pi i \xi y} dy \\ &= \widehat{f}(\xi) \widehat{g}(\xi). \end{aligned}$$

As in the example above, the utility of Fubini's theorem is often interchanging the order of an iterated integral, and the product measure acts simply as an auxiliary object to justify this swap. In practice, this means that we do not need to be particularly concerned by the fact that there may be more than one product measure. The validity of the Fubini–Tonelli theorem for s -finite (and not necessarily σ -finite) measures has found applications in the theory of Markov processes [3].

2. The Maximal Product Measure

Let (X, \mathcal{B}) and (Y, \mathcal{C}) be measurable spaces. The intersection of measurable rectangles $B_1 \times C_1$ and $B_2 \times C_2$ is again a measurable rectangle: $(B_1 \times C_1) \cap (B_2 \times C_2) = (B_1 \cap B_2) \times (C_1 \cap C_2)$. The complement of a measurable rectangle is a disjoint union of three measurable rectangles: $(B \times C)^c = (B^c \times C) \sqcup (B \times C^c) \sqcup (B^c \times C^c)$. Thus, the family of measurable rectangles is a semi-algebra on $X \times Y$. We can therefore build a product measure using an outer measure construction similar to what appeared in Section 3.

We may define an algebra

$$\mathcal{A} = \left\{ \bigsqcup_{i=1}^n (B_i \times C_i) : n \in \mathbb{N}, B_1 \times C_1, \dots, B_n \times C_n \text{ pairwise disjoint measurable rectangles} \right\}.$$

Given measures $\mu : \mathcal{B} \rightarrow [0, \infty]$ and $\nu : \mathcal{C} \rightarrow [0, \infty]$ on X and Y respectively, we define a premeasure ρ_0 on \mathcal{A} by

$$\rho_0 \left(\bigsqcup_{i=1}^n (B_i \times C_i) \right) = \sum_{i=1}^n \mu(B_i) \nu(C_i).$$

We can then extend ρ_0 to an outer measure

$$\begin{aligned} \rho^*(E) &= \inf \left\{ \sum_{n=1}^{\infty} \rho_0(A_n) : E \subseteq \bigcup_{n \in \mathbb{N}} A_n, A_n \in \mathcal{A} \right\} \\ &= \inf \left\{ \sum_{n=1}^{\infty} \mu(B_n) \nu(C_n) : E \subseteq \bigcup_{n \in \mathbb{N}} (B_n \times C_n), B_n \in \mathcal{B}, C_n \in \mathcal{C} \right\} \end{aligned}$$

by Proposition 5.22. By Lemma 5.25, elements of \mathcal{A} are ρ^* -measurable, and so $\rho = \rho^*|_{\mathcal{B} \otimes \mathcal{C}}$ defines a product measure by Theorem 5.24. When it is ambiguous (i.e., when dealing with non- σ -finite spaces), we will denote this product measure by $\mu \stackrel{\text{max}}{\otimes} \nu$ and refer to it as the *maximal product measure* of μ and ν . The reason for this terminology is the following theorem.

THEOREM 7.10

Let (X, \mathcal{B}, μ) and (Y, \mathcal{C}, ν) be measure spaces. Then for every $E \in \mathcal{B} \otimes \mathcal{C}$,

$$(\mu \stackrel{\text{max}}{\otimes} \nu)(E) = \sup \{ \rho(E) : \rho \text{ is a product measure of } \mu \text{ and } \nu \}.$$

Moreover, if $(\mu \stackrel{\text{max}}{\otimes} \nu)(E) < \infty$, then $\rho(E) = (\mu \stackrel{\text{max}}{\otimes} \nu)(E)$ for every product measure ρ .

EXAMPLE 7.11

Let $X = [0, 1]$, $\mathcal{B} = \text{Borel}([0, 1])$, and let μ be the Lebesgue measure on $[0, 1]$. Let $Y = [0, 1]$ also but with the σ -algebra $\mathcal{C} = \mathcal{P}([0, 1])$ and counting measure ν . Note that, since Y is uncountable, ν is not s -finite. However, μ is (s -)finite, so can define a cross-sectional product measure $\rho : \mathcal{B} \otimes \mathcal{C} \rightarrow [0, \infty]$ by

$$\rho(E) = \int_Y \mu(E^y) d\nu(y) = \sum_{y \in Y} \mu(E^y).$$

To see that this is different than the product measure $\mu \stackrel{\text{max}}{\otimes} \nu$, consider the diagonal $\Delta = \{(t, t) : t \in [0, 1]\}$.

Using the cross-sectional product measure, we have

$$\rho(\Delta) = \sum_{y \in Y} \mu(\{y\}) = 0.$$

Now let us compute the measure $(\mu \stackrel{\text{max}}{\otimes} \nu)(\Delta)$. Let $(B_n \times C_n)_{n \in \mathbb{N}}$ be a family of measurable rectangles such that $\Delta \subseteq \bigcup_{n \in \mathbb{N}} (B_n \times C_n)$. Let $S = \{n \in \mathbb{N} : \mu(B_n) = 0\}$, and let $E = \bigcup_{n \in S} B_n$. Then $F = [0, 1] \setminus E$ has $\mu(F) = 1$, and $\Delta_F = \{(t, t) : t \in F\}$ satisfies $\Delta_F \subseteq \bigcup_{n \notin S} (B_n \times C_n)$. Since $\mu(F) = 1$, F is uncountable. But $F \subseteq \bigcup_{n \notin S} C_n$, so C_{n_0} is uncountable (in particular, infinite) for some $n_0 \notin S$. Therefore,

$$\sum_{n=1}^{\infty} \mu(B_n) \nu(C_n) \geq \underbrace{\mu(B_{n_0})}_{>0} \underbrace{\nu(C_{n_0})}_{\infty} = \infty.$$

This proves $(\mu \stackrel{\text{max}}{\otimes} \nu)(\Delta) = \infty$.

PROOF OF THEOREM 7.10. Let ρ be a product measure of μ and ν and let $E \in \mathcal{B} \otimes \mathcal{C}$. Suppose $(B_n \times C_n)_{n \in \mathbb{N}}$ is a family of measurable rectangles such that $E \subseteq \bigcup_{n \in \mathbb{N}} (B_n \times C_n)$. Then by countable subadditivity of ρ and the fact that ρ is a product measure, we have

$$\rho(E) \leq \sum_{n=1}^{\infty} \rho(B_n \times C_n) = \sum_{n=1}^{\infty} \mu(B_n) \nu(C_n).$$

Taking an infimum over such families, we conclude $\rho(E) \leq (\mu \stackrel{\text{max}}{\otimes} \nu)(E)$. This proves the first part of the theorem.

Suppose $E \in \mathcal{B} \otimes \mathcal{C}$ and $(\mu \stackrel{\max}{\otimes} \nu)(E) < \infty$. Let $\varepsilon > 0$. There exists a family $(B_n \times C_n)_{n \in \mathbb{N}}$ of measurable rectangles such that $E \subseteq \bigcup_{n \in \mathbb{N}} (B_n \times C_n)$ and

$$\sum_{n=1}^{\infty} \mu(B_n) \nu(C_n) \leq (\mu \stackrel{\max}{\otimes} \nu)(E) + \varepsilon.$$

Let $A = \bigcup_{n \in \mathbb{N}} (B_n \times C_n)$. By countable subadditivity and monotonicity,

$$(\mu \stackrel{\max}{\otimes} \nu)(E) \leq (\mu \stackrel{\max}{\otimes} \nu)(A) \leq \sum_{n=1}^{\infty} \mu(B_n) \nu(C_n) \leq (\mu \stackrel{\max}{\otimes} \nu)(E) + \varepsilon.$$

In particular, $(\mu \stackrel{\max}{\otimes} \nu)(A \setminus E) < \varepsilon$. By the first part of the theorem, it follows that $\rho(A \setminus E) < \varepsilon$.

Define a sequence $(A_n)_{n \in \mathbb{N}}$ in the algebra \mathcal{A} generated by measurable rectangles by $A_1 = B_1 \times C_1$ and $A_n = (B_n \times C_n) \setminus (A_1 \cup \dots \cup A_{n-1})$ for $n \geq 2$. Then the sets $(A_n)_{n \in \mathbb{N}}$ are pairwise disjoint and $\bigcup_{n \in \mathbb{N}} A_n = A$. Moreover, since the rectangles satisfy $\mu(B_n) \nu(C_n) < \infty$ for every $n \in \mathbb{N}$, additivity of the arbitrary product measure ρ implies that the value $\rho(A_n)$ is the same for every product measure. Therefore,

$$\rho(A) = \sum_{n=1}^{\infty} \rho(A_n) = \sum_{n=1}^{\infty} (\mu \stackrel{\max}{\otimes} \nu)(A_n) = (\mu \stackrel{\max}{\otimes} \nu)(A).$$

Thus,

$$\rho(E) = \rho(A) - \rho(A \setminus E) > (\mu \stackrel{\max}{\otimes} \nu)(A) - \varepsilon.$$

Combining with the first part of the theorem, we conclude $\rho(E) = (\mu \stackrel{\max}{\otimes} \nu)(E)$. □

Exercises

7.1 The goal of this exercise is to show that integration computes the “area under the curve”. Let (X, \mathcal{B}, μ) be an s-finite measure space. Given a measurable function $f : X \rightarrow [0, \infty]$, we may define the region “under” f as the set

$$R_f = \{(x, t) \in X \times \mathbb{R} : 0 \leq t \leq f(x)\}.$$

Let λ be the Lebesgue measure on \mathbb{R} . Prove:

(a) If $f : X \rightarrow [0, \infty]$ is measurable, then $R_f \in \mathcal{B} \otimes \text{Borel}(\mathbb{R})$, and

$$\int_X f \, d\mu = (\mu \stackrel{\text{cs}}{\otimes} \lambda)(R_f).$$

(b) If $f : X \rightarrow \mathbb{R}$ is integrable, then

$$\int_X f \, d\mu = (\mu \stackrel{\text{cs}}{\otimes} \lambda)(R_{f^+}) - (\mu \stackrel{\text{cs}}{\otimes} \lambda)(R_{f^-}).$$

7.2 Let λ be the Lebesgue measure on \mathbb{R} . Suppose $A, B \subseteq \mathbb{R}$ are Lebesgue-measurable sets such that $\lambda(A), \lambda(B) > 0$. Prove that there exists $t \in \mathbb{R}$ such that $\lambda(A \cap (B - t)) > 0$.

7.3 Let $X = [0, 1]$ with the standard topology, and let $Y = [0, 1]$ with the discrete topology. Define a functional $\varphi : C_c(X \times Y) \rightarrow \mathbb{C}$ by $\varphi(f) = \sum_{y \in [0, 1]} \int_0^1 f(x, y) \, dx$, where the inner integral is the Riemann integral. Let μ be the Radon measure representing φ . Is μ a product measure of $\lambda|_{\text{Borel}([0,1])}$ and the counting measure?

Part 3

Additional Topics in Measure Theory

CHAPTER 8

L^p Spaces

1. Topological Vector Spaces

DEFINITION 8.1

A *topological vector space* is a pair (V, τ) such that V is a (real or complex) vector space and τ is a topology on V such that

- the addition map $V \times V \rightarrow V$, $(u, v) \mapsto u + v$, is continuous, and
- the map $\mathbb{K} \times V \rightarrow V$, $(c, v) \mapsto cv$ is continuous, where \mathbb{K} is the field of scalars.

EXAMPLE 8.2

The Euclidean space $V = \mathbb{R}^d$ with the standard topology is a topological vector space for $d \in \mathbb{N}$.

A typical means of defining a topology on a vector space is with a norm. Recall (Definition 3.18) that a *norm* on a vector space V is a function $\|\cdot\| : V \rightarrow [0, \infty)$ such that

- TRIANGLE INEQUALITY: $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in V$;
- ABSOLUTE HOMOGENEITY: $\|cv\| = |c| \|v\|$ for all $v \in V$ and all scalars c ; and
- POSITIVE DEFINITENESS: if $v \in V$ and $\|v\| = 0$, then $v = 0$.

The triangle inequality and positive definiteness imply that the function $d : V \times V \rightarrow [0, \infty)$ defined by $d(u, v) = \|u - v\|$ is a metric on V , so we can endow V with this metric space topology.

DEFINITION 8.3

A normed vector space $(V, \|\cdot\|)$ is a *Banach space* if it is complete (as a metric space).

EXAMPLE 8.4

The standard Euclidean norm on \mathbb{R}^d , i.e. $\|(x_1, \dots, x_d)\| = \sqrt{x_1^2 + \dots + x_d^2}$, makes \mathbb{R}^d into a Banach space.

One way of studying normed spaces (or more general topological vector spaces) is in terms of linear functionals. We have already seen the utility of understanding linear functionals on a normed space in the Riesz representation theorem, which described the positive linear functionals on the normed space $C_c(X)$ as Radon measures on the underlying LCH space X .

DEFINITION 8.5

Let \mathbb{K} be either \mathbb{R} or \mathbb{C} . Let $(V, \|\cdot\|_V)$ be a normed vector space over \mathbb{K} . A *linear functional* is a linear map $\varphi : V \rightarrow \mathbb{K}$. The *dual space* V^* is the vector space of continuous linear

functionals on V with norm

$$\|\varphi\|_{V^*} = \sup_{\|v\|_V \leq 1} |\varphi(v)|.$$

PROPOSITION 8.6

Let V be a normed vector space. Then the dual V^* is a Banach space.

PROOF. We need to prove two things: (1) $\|\cdot\|_{V^*}$ defines a norm on V^* and (2) $(V^*, \|\cdot\|_{V^*})$ is complete.

CLAIM 1. $\|\cdot\|_{V^*}$ is a norm on V^* .

Let $\varphi, \psi \in V^*$. Then for $v \in V$ with $\|v\|_V \leq 1$, we have

$$|(\varphi + \psi)(v)| \leq |\varphi(v)| + |\psi(v)| \leq \|\varphi\|_{V^*} + \|\psi\|_{V^*},$$

so $\|\cdot\|_{V^*}$ satisfies the triangle inequality.

Suppose $\varphi \in V^*$ and $c \in \mathbb{K}$. Then for $v \in V$, we have $|(c\varphi)(v)| = |c\varphi(v)| = |c|\varphi(v)|$. Taking a supremum over $v \in V$ with $\|v\|_V \leq 1$ gives absolute homogeneity of $\|\cdot\|_{V^*}$.

Finally, if $\|\varphi\|_{V^*} = 0$, then $\varphi(v) = 0$ for every $v \in V$ with $\|v\|_V \leq 1$. Hence, for arbitrary $v \in V \setminus \{0\}$, we have $\varphi(v) = \|v\|_V \varphi\left(\frac{v}{\|v\|_V}\right) = 0$, so $\varphi = 0$. Thus, $\|\cdot\|_{V^*}$ is positive definite.

CLAIM 2. $(V^*, \|\cdot\|_{V^*})$ is complete.

Suppose $(\varphi_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in V^* . Then for $v \in V$, the sequence $(\varphi_n(v))_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathbb{K} , since $|\varphi_n(v) - \varphi_m(v)| \leq \|\varphi_n - \varphi_m\|_{V^*} \|v\|_V$. Since the field \mathbb{K} is complete, we may define $\varphi(v) = \lim_{n \rightarrow \infty} \varphi_n(v)$ for every $v \in V$. We must check that $\varphi \in V^*$ and $\varphi_n \rightarrow \varphi$ in V^* .

First, $\|\varphi\|_{V^*} \leq \lim_{n \rightarrow \infty} \|\varphi_n\|_{V^*} < \infty$, so $\varphi \in V^*$ (see Exercise 8.2). Next, given $v \in V$ with $\|v\|_V \leq 1$, we have

$$|\varphi(v) - \varphi_n(v)| = \left| \lim_{m \rightarrow \infty} \varphi_m(v) - \varphi_n(v) \right| \leq \sup_{m \geq n} |\varphi_m(v) - \varphi_n(v)| \leq \sup_{m \geq n} \|\varphi_m - \varphi_n\|_{V^*}.$$

Thus,

$$\limsup_{n \rightarrow \infty} \|\varphi - \varphi_n\|_{V^*} \leq \lim_{n \rightarrow \infty} \sup_{m \geq n} \|\varphi_m - \varphi_n\|_{V^*} = 0,$$

so $\varphi_n \rightarrow \varphi$. □

2. L^p Norms

For the remainder of this chapter, we will focus on vector spaces of functions associated with a measure space.

DEFINITION 8.7

Let (X, \mathcal{B}, μ) be a measure space.

- For $1 \leq p < \infty$, the *L^p norm* of a measurable function $f : X \rightarrow \mathbb{C}$ is the quantity

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{1/p}.$$

- The *$(\mu\text{-})$ essential supremum* of a nonnegative measurable function $f : X \rightarrow [0, \infty]$ is

$$\text{ess sup}(f) = \inf \{c \geq 0 : f \leq c \text{ } \mu\text{-a.e.}\} = \inf \{c \geq 0 : \mu(\{f > c\}) = 0\}.$$

- The *L^∞ norm* of a measurable function $f : X \rightarrow \mathbb{C}$ is $\|f\|_\infty = \text{ess sup}(|f|)$.
- For $1 \leq p \leq \infty$, the *L^p space* $L^p(\mu)$ is the space

$$L^p(\mu) = \left\{ [f]_\mu : \|f\|_p < \infty \right\},$$

where $[f]_\mu$ is the equivalence class $[f]_\mu = \{g : X \rightarrow \mathbb{C} \text{ measurable} : g = f \text{ } \mu\text{-a.e.}\}$.

REMARK. If $f, g : X \rightarrow \mathbb{C}$ are measurable functions and $f = g \mu\text{-a.e.}$, then $\int_X |f|^p d\mu = \int_X |g|^p d\mu$, so the value of the L^p norm only depends on the equivalence class and not the choice of representative.

While for technical reasons, we view $L^p(\mu)$ as a space of equivalence classes of functions, it is often more convenient to work with actual functions (i.e., representatives of the equivalence classes). As long as we only perform “countable operations” on functions, we can safely pass between the two different points of view, since the family of null sets is a σ -ideal. For this reason, it is standard practice in mathematics to write expressions like “ $f \in L^p(\mu)$,” and we will also engage in this abuse of notation.

LEMMA 8.8

Let (X, \mathcal{B}, μ) be a measure space, and let $f : X \rightarrow \mathbb{C}$ be a measurable function. Then $|f| \leq \|f\|_\infty \mu\text{-a.e.}$

PROOF. Let $M = \|f\|_\infty$. If $M = \infty$, there is nothing to show, so assume $M < \infty$. By the definition of the essential supremum, we have $\mu(\{|f| > M + \frac{1}{n}\}) = 0$ for each $n \in \mathbb{N}$. Taking a union over $n \in \mathbb{N}$ and applying continuity from below of the measure μ , we conclude $\mu(\{|f| > M\}) = 0$. That is, $|f| \leq M$ a.e. \square

One of the main goals of this chapter is to prove that $\|\cdot\|_p$ defines a norm for every $p \in [1, \infty]$ and the vector space $L^p(\mu)$ is a Banach space.

3. Convexity and the Inequalities of Jensen and Minkowski

DEFINITION 8.9

Let V be a real vector space.

- A set $C \subseteq V$ is *convex* if it contains the entire line segment between every pair of points in C . That is, for every $x, y \in C$ and every $t \in [0, 1]$, one has $tx + (1-t)y \in C$.
- Let C be a convex set. A function $\varphi : C \rightarrow \mathbb{R}$ is *convex* if $\varphi(tx + (1-t)y) \leq t\varphi(x) + (1-t)\varphi(y)$ for all $x, y \in C$ and $t \in [0, 1]$.

By induction, one can show that convex sets are closed under convex combinations: if $x_1, \dots, x_n \in C$ and $\lambda_1, \dots, \lambda_n \geq 0$ are coefficients with $\sum_{j=1}^n \lambda_j = 1$, then $\sum_{j=1}^n \lambda_j x_j \in C$. Similarly, if f is a convex function, then its behavior on convex combinations is governed by the inequality

$\varphi\left(\sum_{j=1}^n \lambda_j x_j\right) \leq \sum_{j=1}^n \lambda_j \varphi(x_j)$. This is the discrete version of a fundamental inequality for convex functions known as Jensen's inequality.

THEOREM 8.10: JENSEN'S INEQUALITY

Let (X, \mathcal{B}, μ) be a probability space. Let $f : X \rightarrow I$ be an integrable function taking values in an interval $I \subseteq \mathbb{R}$, and suppose $\varphi : I \rightarrow \mathbb{R}$ is convex. Then

$$\varphi\left(\int_X f \, d\mu\right) \leq \int_X \varphi \circ f \, d\mu.$$

PROOF. Let $t = \int_X f \, d\mu \in I$. We want to show $\int_X \varphi \circ f \, d\mu \geq \varphi(t)$.

If $t \in \partial I$, then $f = t$ a.e., so $\varphi \circ f = \varphi(t)$ a.e. and $\int_X \varphi \circ f \, d\mu = \varphi(t)$.

Suppose $t \in \text{int}(I)$. By Exercise 8.3, let $m \in \mathbb{R}$ such that

$$\varphi(s) \geq m(s - t) + \varphi(t) \quad (\forall s \in I).$$

Then

$$\int_X \varphi \circ f \, d\mu \geq \int_X (m(f - t) + \varphi(t)) \, d\mu = m \underbrace{\int_X f \, d\mu}_{t} - mt + \varphi(t) = \varphi(t).$$

□

The central inequality for the L^p norm is Minkowski's inequality, which shows that $\|\cdot\|_p$ is indeed a norm.

THEOREM 8.11: MINKOWSKI'S INEQUALITY

Let (X, \mathcal{B}, μ) be a measure space, and let $p \in [1, \infty]$. Let $f, g : X \rightarrow \mathbb{C}$ be measurable functions. Then

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

Before proving Minkowski's inequality, let us confirm that it makes $(L^p(\mu), \|\cdot\|_p)$ into a normed vector space.

COROLLARY 8.12

Let (X, \mathcal{B}, μ) be a measure space, and let $p \in [1, \infty]$. Then $L^p(\mu)$ is a vector space, and $\|\cdot\|_p : L^p(\mu) \rightarrow [0, \infty)$ is a norm.

PROOF. Let $f, g \in L^p(\mu)$. (Technically, we should say $[f]_\mu, [g]_\mu \in L^p(\mu)$, but this can become notationally cumbersome and is not standard practice in mathematics; see the remark following Definition 8.7.) By Minkowski's inequality, $\|f + g\|_p \leq \|f\|_p + \|g\|_p < \infty$, so $f + g \in L^p(\mu)$. Moreover, if $p < \infty$, then given $f \in L^p(\mu)$ and $c \in \mathbb{C}$, we have

$$\|cf\|_p = \left(\int_X |cf|^p \, d\mu \right)^{1/p} = \left(|c|^p \int_X |f|^p \, d\mu \right)^{1/p} = |c| \|f\|_p < \infty, \quad (8.1)$$

so $cf \in L^p(\mu)$. If $p = \infty$, then we can similarly check that $\|cf\|_\infty = |c| \|f\|_\infty$. This proves that $L^p(\mu)$ is a vector space. Minkowski's inequality is the triangle inequality for the L^p norm, and (8.1) proves absolute homogeneity. Finally, $\|\cdot\|_p$ is positive definite by Proposition 3.23 in the case $p < \infty$ and by Lemma 8.8 when $p = \infty$. □

REMARK. The final line of the proof of Corollary 8.12 reveals why we want to view L^p as a space of equivalence classes rather than a space of functions: on the space of functions, $\|\cdot\|_p$ is only a seminorm, but working with equivalence classes of functions turns $\|\cdot\|_p$ into a proper norm.

PROOF OF MINKOWSKI'S INEQUALITY. We split the proof into two cases: $p < \infty$ or $p = \infty$.

CASE 1. $p < \infty$

Let $B = \{w : X \rightarrow \mathbb{C} \text{ measurable} : \|w\|_p \leq 1\}$. We will show that B is a convex set. Let $u, v \in B$ and $t \in [0, 1]$. Then

$$\begin{aligned} \int_X |tu + (1-t)v|^p d\mu &\leq \int_X (t|u| + (1-t)|v|)^p d\mu && \text{(triangle inequality)} \\ &\leq \int_X (t|u|^p + (1-t)|v|^p) d\mu && (x \mapsto x^p \text{ is convex}) \\ &= t\|u\|_p^p + (1-t)\|v\|_p^p \leq 1. \end{aligned}$$

Now let $f, g : X \rightarrow \mathbb{C}$ be measurable functions. If $\|f\|_p = \infty$ or $\|g\|_p = \infty$, there is nothing to prove. Moreover, if $\|f\|_p = 0$, then $f = 0$ a.e., so $f + g = g$ a.e., and there is nothing to prove. Similarly, there is nothing to show in the case that $\|g\|_p = 0$. Thus, we may assume $0 < \|f\|_p, \|g\|_p < \infty$. Then $u = \frac{f}{\|f\|_p}, v = \frac{g}{\|g\|_p} \in B$. Hence, letting $t = \frac{\|f\|_p}{\|f\|_p + \|g\|_p}$, we have

$$\frac{\|f + g\|_p}{\|f\|_p + \|g\|_p} = \|tu + (1-t)v\|_p \leq 1.$$

CASE 2. $p = \infty$

By Lemma 8.8, the sets $N_1 = \{|f| > \|f\|_\infty\}$ and $N_2 = \{|g| > \|g\|_\infty\}$ are μ -null sets. Therefore, $N = N_1 \cup N_2$ is a null set. Suppose $x \in X \setminus N$. Then

$$|f(x) + g(x)| \leq |f(x)| + |g(x)| \leq \|f\|_\infty + \|g\|_\infty.$$

Therefore, $\{|f + g| > \|f\|_\infty + \|g\|_\infty\} \subseteq N$ is a null set, so $\|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty$.

□

4. Riesz–Fischer Theorem

Our next goal is prove that $(L^p(\mu), \|\cdot\|_p)$ is a Banach space.

THEOREM 8.13

Let (X, \mathcal{B}, μ) be a measure space, and let $1 \leq p < \infty$. Suppose $(f_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $L^p(\mu)$. Then there is a subsequence $(f_{n_k})_{k \in \mathbb{N}}$ that converges a.e. to a function $f \in L^p(\mu)$ and $\|f_n - f\|_p \rightarrow 0$.

PROOF. The strategy of the proof is to choose a subsequence along which f_{n_k} is “quickly Cauchy” and then to show that “quickly Cauchy” sequences converge a.e. and in L^p .

CLAIM 1. There exists a subsequence $n_1 < n_2 < \dots$ such that

$$\|f_{n_{k+1}} - f_{n_k}\|_p < 2^{-k} \quad (8.2)$$

for every $k \in \mathbb{N}$.

We construct $(n_k)_{k \in \mathbb{N}}$ by induction. Choose $n_1 \in \mathbb{N}$ such that

$$\sup_{m \geq n_1} \|f_{n_1} - f_m\|_p < \frac{1}{2}.$$

Given n_1, \dots, n_k , choose $n_{k+1} > n_k$ such that

$$\sup_{m \geq n_{k+1}} \|f_{n_{k+1}} - f_m\|_p < 2^{-(k+1)}.$$

Then by induction, we have constructed a sequence $n_1 < n_2 < \dots$ satisfying (8.2).

Let $g_k = f_{n_{k+1}} - f_{n_k}$ for $k \in \mathbb{N}$, and let $G_k = \sum_{j=1}^k |g_j|$. By Minkowski's inequality, $\|G_k\|_p \leq \sum_{j=1}^k \|g_j\|_p < 1$. Letting $G = \sum_{j=1}^{\infty} |g_j|$, we have by the monotone convergence theorem that

$$\int_X G^p d\mu = \lim_{k \rightarrow \infty} \underbrace{\int_X G_k^p d\mu}_{\|G_k\|_p^p} \leq 1.$$

In particular, $G < \infty$ a.e., so the series $g(x) = \sum_{j=1}^{\infty} g_j(x)$ converges absolutely for a.e. $x \in X$. Let $f = g + f_{n_1}$.

CLAIM 2. $f_{n_k} \rightarrow f$ a.e as $k \rightarrow \infty$

Let $S_k = \sum_{j=1}^{k-1} g_j$. Then $S_k \rightarrow g$ a.e. (by the definition of g), so $f_{n_1} + S_k \rightarrow f$ a.e. as $k \rightarrow \infty$. But expanding g_j , the sum S_k is telescoping and we have

$$f_{n_1} + S_k = f_{n_1} + (f_{n_2} - f_{n_1}) + \dots + (f_{n_k} - f_{n_{k-1}}) = f_{n_k}.$$

CLAIM 3. $\|f_n - f\|_p \rightarrow 0$.

Note that $|f_{n_k} - f| = |\sum_{j=k+1}^{\infty} g_j| \leq \sum_{j=k+1}^{\infty} |g_j|$, so by Minkowski's inequality, for $n \geq n_k$,

$$\|f_n - f\|_p \leq \|f_n - f_{n_k}\|_p + \|f_{n_k} - f\|_p \leq 2^{-k} + \sum_{j=k+1}^{\infty} 2^{-j} = 2 \cdot 2^{-k}$$

Therefore, $\|f_n - f\|_p \rightarrow 0$ as $n \rightarrow \infty$.

CLAIM 4. $f \in L^p(\mu)$

By Minkowski's inequality, $\|f\|_p \leq \|f_n\|_p + \|f_n - f\|_p$. The term $\|f_n\|_p$ is finite, and $\|f_n - f\|_p < 1$ for all large enough n .

□

THEOREM 8.14

Let (X, \mathcal{B}, μ) be a measure space. Suppose $(f_n)_{n \in \mathbb{N}}$ is a Cauchy sequence $L^\infty(\mu)$. Then there exists a measurable set $X_0 \subseteq X$ such that $\mu(X \setminus X_0) = 0$ and $(f_n)_{n \in \mathbb{N}}$ is uniformly bounded and uniformly Cauchy on X_0 . In particular, $(f_n)_{n \in \mathbb{N}}$ converges a.e. and in L^∞ to a function $f \in L^\infty(\mu)$.

PROOF. Put

$$X_0 = \underbrace{\bigcap_{n \in \mathbb{N}} \{|f_n| \leq \|f_n\|_\infty\}}_{\text{uniformly bounded}} \cap \underbrace{\bigcap_{n, m \in \mathbb{N}} \{|f_n - f_m| \leq \|f_n - f_m\|_\infty\}}_{\text{uniformly Cauchy}}.$$

Since $(f_n)_{n \in \mathbb{N}}$ is uniformly Cauchy on X_0 , we may define $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ as the (uniform) limit of f_n on X_0 . The values of f outside of X_0 will not play any role, so let us set $f(x) = 0$ for $x \in X \setminus X_0$. Then f is measurable and uniformly bounded, so $f \in L^\infty(\mu)$. Moreover, $f_n \rightarrow f$ uniformly outside of the null set $X \setminus X_0$, so $\|f_n - f\|_\infty \leq \sup_{x \in X_0} |f_n(x) - f(x)| \rightarrow 0$. \square

Combining Theorem 8.13 (for $p < \infty$) and Theorem 8.14 (for $p = \infty$), we obtain the Riesz–Fischer theorem.

THEOREM 8.15: RIESZ–FISCHER THEOREM

Let (X, \mathcal{B}, μ) be a measure space, and let $1 \leq p \leq \infty$. Then $(L^p(\mu), \|\cdot\|_p)$ is a Banach space.

5. Hölder's Inequality

LEMMA 8.16: WEIGHTED ARITHMETIC MEAN–GEOMETRIC MEAN INEQUALITY

Let $n \in \mathbb{N}$, $x_1, \dots, x_n \geq 0$, and $\lambda_1, \dots, \lambda_n \geq 0$ such that $\sum_{j=1}^n \lambda_j = 1$. Then

$$\prod_{j=1}^n x_j^{\lambda_j} \leq \sum_{j=1}^n \lambda_j x_j.$$

PROOF. The function $x \mapsto \log x$ is strictly increasing, so it suffices to prove the inequality after taking the logarithm of both sides. But $x \mapsto -\log x$ is also a convex function, so by Jensen's inequality,

$$-\log \left(\sum_{j=1}^n \lambda_j x_j \right) \leq -\sum_{j=1}^n \lambda_j \log x_j = -\log \left(\prod_{j=1}^n x_j^{\lambda_j} \right).$$

\square

THEOREM 8.17: HÖLDER'S INEQUALITY

Let (X, \mathcal{B}, μ) be a measure space. Let $f, g : X \rightarrow \mathbb{C}$ be measurable functions. Let $p, q \in [1, \infty]$ such that $\frac{1}{p} + \frac{1}{q} = 1$. Then

$$\|fg\|_1 \leq \|f\|_p \|g\|_q.$$

More generally, if $n \in \mathbb{N}$, $f_1, \dots, f_n : X \rightarrow \mathbb{C}$ are measurable functions, and $p_1, \dots, p_n \in [1, \infty]$ such that $\sum_{j=1}^n \frac{1}{p_j} = 1$, then

$$\left\| \prod_{j=1}^n f_j \right\|_1 \leq \prod_{j=1}^n \|f_j\|_{p_j}.$$

PROOF. Let $n \in \mathbb{N}$, let $f_1, \dots, f_n : X \rightarrow \mathbb{C}$ be measurable functions, and suppose $p_1, \dots, p_n \in [1, \infty]$ such that $\sum_{j=1}^n \frac{1}{p_j} = 1$. If $\|f_j\|_{p_j} = 0$ for some j , then $f_j = 0$ a.e., so $\prod_{j=1}^n f_j = 0$ a.e., and there is nothing to prove. We may therefore assume $\|f_j\|_{p_j} > 0$ for all j . Now if $\|f_j\|_{p_j} = \infty$ for some j , then $\prod_{j=1}^n \|f_j\|_{p_j} = \infty$, and again there is nothing to prove, so we may assume $0 < \|f_j\|_{p_j} < \infty$.

For each j , let $u_j = \frac{f_j}{\|f_j\|_{p_j}}$ so that $\|f_j\|_{p_j} = 1$. Let $J = \{1 \leq j \leq n : p_j = \infty\}$, and note that $|u_j| \leq 1$ a.e. for $j \in J$. Moreover, $\sum_{j \notin J} \frac{1}{p_j} = 1$, so by Lemma 8.16,

$$\frac{\left\| \prod_{j=1}^n f_j \right\|_1}{\prod_{j=1}^n \|f_j\|_{p_j}} \leq \int_X \prod_{j=1}^n |u_j| \, d\mu = \int_X \prod_{j \notin J} (|u_j|^{p_j})^{1/p_j} \, d\mu \leq \sum_{j \notin J} \frac{1}{p_j} \underbrace{\int_X |u_j|^{p_j} \, d\mu}_{\|u_j\|_{p_j}^{p_j}=1} = 1.$$

□

DEFINITION 8.18

The exponents $p \in [1, \infty]$ and $q \in [1, \infty]$ are called *conjugate* if $\frac{1}{p} + \frac{1}{q} = 1$.

The only self-conjugate exponent is $p = 2$, and this provides the space $L^2(\mu)$ with the additional structure of an inner product space. Namely, defining $\langle \cdot, \cdot \rangle : L^2(\mu) \times L^2(\mu) \rightarrow \mathbb{C}$ by $\langle f, g \rangle = \int_X f\bar{g} \, d\mu$, we have the following properties:

- CONJUGATE SYMMETRY: $\langle f, g \rangle = \overline{\langle g, f \rangle}$;
- LINEARITY IN THE FIRST ARGUMENT: $\langle cf + g, h \rangle = c\langle f, h \rangle + \langle g, h \rangle$;
- POSITIVE DEFINITENESS: $f \neq 0 \implies \langle f, f \rangle > 0$.

The norm induced by this inner product is the L^2 norm; that is, $\|f\|_2 = \langle f, f \rangle$. The special case of Hölder's inequality for L^2 functions is the *Cauchy–Schwarz inequality*: $|\langle f, g \rangle| \leq \|f\|_2 \|g\|_2$. Complete inner product spaces (such as $L^2(\mu)$) are called *Hilbert spaces* and have many advantages over general normed spaces. For example, one can discuss orthogonality of elements of $L^2(\mu)$ and work with *orthonormal bases* for the space. Also, Hilbert spaces are self-dual in the sense that every linear functional on a Hilbert space can be represented as an inner product with a fixed element of the Hilbert space. We will not discuss Hilbert spaces in more detail in this course, but they play an important role in functional analysis and have many applications in physics.

6. Convolutions and Young's Inequality

DEFINITION 8.19

Let $f, g : \mathbb{R} \rightarrow \mathbb{C}$ be Lebesgue-measurable functions. The *convolution* of f and g is defined by

$$(f * g)(x) = \int_{\mathbb{R}} f(x - y)g(y) dy$$

when $y \mapsto f(x - y)g(y)$ is integrable.

We saw in the section on Fubini's theorem that the convolution of integrable functions is defined a.e. and integrable (see Example 7.9). Young's inequality provides a substantial generalization for convolutions of functions from other L^p spaces.

THEOREM 8.20: YOUNG'S INEQUALITY

Suppose $1 \leq p, q, r \leq \infty$ and $\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{r}$. Let $f \in L^p(\mathbb{R})$ and $g \in L^q(\mathbb{R})$. Then $f * g$ is defined a.e. and

$$\|f * g\|_r \leq \|f\|_p \|g\|_q.$$

PROOF. When $r = \infty$ (i.e., p and q are conjugate), then $f * g$ is defined everywhere and is bounded by $\|f\|_p \|g\|_q$ by Hölder's inequality.

For $r < \infty$, defining new constants s, t by $\frac{1}{s} = 1 - \frac{1}{q}$ and $\frac{1}{t} = 1 - \frac{1}{p}$, we have the identity

$$ab = (a^p b^q)^{1/r} (a^p)^{1/s} (b^q)^{1/t} \quad (8.3)$$

for $a, b \geq 0$. Moreover, $\frac{1}{r} + \frac{1}{s} + \frac{1}{t} = 1$, so the generalized Hölder inequality applies. We leave the details of the computations as Exercise 8.8. □

Exercises

8.1 Show that a normed vector space is a topological vector space.

8.2 Let $(V, \|\cdot\|_V)$ be a normed space, and let $\varphi : V \rightarrow \mathbb{K}$ be a linear functional. Show that the following are equivalent:

- (i) φ is continuous;
- (ii) φ is continuous at $0 \in V$;
- (iii) φ is bounded, i.e. $\|\varphi\|_{V^*} = \sup_{\|v\|_V \leq 1} |\varphi(v)| < \infty$.

8.3 Let $I \subseteq \mathbb{R}$ be an interval, and let $\varphi : I \rightarrow \mathbb{R}$ be a convex function. Suppose $t \in \text{int}(I)$. Show that there is a number $m \in \mathbb{R}$ such that for every $s \in I$,

$$\varphi(s) \geq m(s - t) + \varphi(t).$$

8.4 Let (X, \mathcal{B}, μ) be a measure space. Suppose $f : X \rightarrow \mathbb{C}$ is a measurable function, and $f \in L^{p_0}(\mu)$ for some $p_0 \in [1, \infty)$ and $f \in L^\infty(\mu)$. Show that $\|f\|_\infty = \lim_{p \rightarrow \infty} \|f\|_p$.

8.5 Let $p, q \in (1, \infty)$ be conjugate exponents, and let $f \in L^p(\mu)$. Show that

$$\|f\|_p = \sup_{\|g\|_q \leq 1} \left| \int_X f g \, d\mu \right|.$$

8.6

(a) Let (X, \mathcal{B}, μ) be a probability space. Let $f : X \rightarrow \mathbb{C}$ be a measurable function. Show that for any $1 \leq p < q \leq \infty$, one has

$$\|f\|_p \leq \|f\|_q.$$

(b) We denote by $\ell^p(\mathbb{N})$ the L^p space associated to \mathbb{N} with the counting measure. That is, $\ell^p(\mathbb{N})$ is the space of sequences $\mathbf{a} = (a_n)_{n \in \mathbb{N}}$ with $\sum_{n=1}^{\infty} |a_n|^p < \infty$, and the norm on the space is given by $\|\mathbf{a}\|_p = (\sum_{n=1}^{\infty} |a_n|^p)^{1/p}$. Show that for any $1 \leq p < q \leq \infty$, one has

$$\|\mathbf{a}\|_p \geq \|\mathbf{a}\|_q$$

8.7 Suppose $p, q \in (1, \infty)$ are conjugate. Show that $\ell^p(\mathbb{N})^* \cong \ell^q(\mathbb{N})$.

8.8 Complete the proof of Young's inequality.

8.9 The goal of this problem is to show that almost everywhere convergence is a “non-topological” notion of convergence.

(a) Let X be a topological space. Recall that a sequence $(x_n)_{n \in \mathbb{N}}$ in X converges to $x \in X$ if and only if for every open neighborhood U of x , there exists $N \in \mathbb{N}$ such that $x_n \in U$ for $n \geq N$. Show that the following are equivalent:

- (i) $(x_n)_{n \in \mathbb{N}}$ converges to x
- (ii) every subsequence $(x_{n_k})_{k \in \mathbb{N}}$ has a further subsequence $(x_{n_{k_l}})_{l \in \mathbb{N}}$ that converges to x

(b) Let (X, \mathcal{B}, μ) be a measure space, and suppose $(f_n)_{n \in \mathbb{N}}$ is a sequence of integrable functions that converges in $L^1(\mu)$ to an integrable function f . Show that every subsequence of $(f_n)_{n \in \mathbb{N}}$ has a further subsequence that converges μ -almost everywhere to f .

(c) Define the *typewriter sequence* $f_n : [0, 1] \rightarrow \{0, 1\}$ by

$$\begin{aligned} f_1 &= 1, \\ f_2 &= \mathbb{1}_{[0, \frac{1}{2})}, f_3 = \mathbb{1}_{[\frac{1}{2}, 1)} \\ f_4 &= \mathbb{1}_{[0, \frac{1}{4})}, f_5 = \mathbb{1}_{[\frac{1}{4}, \frac{1}{2})}, f_6 = \mathbb{1}_{[\frac{1}{2}, \frac{3}{4})}, f_7 = \mathbb{1}_{[\frac{3}{4}, 1)}, \\ f_8 &= \mathbb{1}_{[0, \frac{1}{8})}, f_9 = \mathbb{1}_{[\frac{1}{8}, \frac{1}{4})}, f_{10} = \mathbb{1}_{[\frac{1}{4}, \frac{3}{8})}, f_{11} = \mathbb{1}_{[\frac{3}{8}, \frac{1}{2})}, f_{12} = \mathbb{1}_{[\frac{1}{2}, \frac{5}{8})}, f_{13} = \mathbb{1}_{[\frac{5}{8}, \frac{3}{4})}, f_{14} = \mathbb{1}_{[\frac{3}{4}, \frac{7}{8})}, f_{15} = \mathbb{1}_{[\frac{7}{8}, 1)} \end{aligned}$$

The general term of the sequence is

$$f_n = \mathbb{1}_{\left[\frac{n-2^k}{2^k}, \frac{n-2^k+1}{2^k}\right)}$$

for $k = \lfloor \log_2(n) \rfloor$. Let λ be the Lebesgue measure on $[0, 1]$. Show that $f_n \rightarrow 0$ in $L^1(\lambda)$, but $(f_n(x))_{n \in \mathbb{N}}$ does not converge for any $x \in [0, 1]$.

(d) Conclude that there is no topology on the space of Lebesgue-measurable functions $f : [0, 1] \rightarrow \mathbb{R}$ such that almost everywhere convergence with respect to Lebesgue measure agrees with convergence in the topology.

CHAPTER 9

Regularity and Littlewood's Principles

The British mathematician J. E. Littlewood laid out three principles for real analysis [5, Section 4.1]:

There are three principles, roughly expressible in the following terms: Every (measurable) set is nearly a finite sum of intervals; every function (of class L^p) is nearly continuous; every convergent sequence of functions is nearly uniformly convergent.

Littlewood's principles were formulated in the context of the Lebesgue measure on \mathbb{R} , but they are in fact a useful guide to measure theory on very general spaces. To make sense of “intervals” or of functions being “nearly continuous” requires a topology, so we will restrict to LCH spaces with Radon measures to address the first and second principle. However, the third principle can be substantiated in fully general measure spaces.

1. The First Principle: Approximation of Measurable Sets

Littlewood's first principle can be recast as a statement about regularity of measures. Let us first interpret it for the Lebesgue measure on \mathbb{R} and then try to generalize. By Proposition 5.31, if $E \subseteq \mathbb{R}$ is a Lebesgue-measurable set, then for every $\varepsilon > 0$, there exists an open set $U \subseteq \mathbb{R}$ such that $E \subseteq U$ and $\lambda(U \setminus E) < \varepsilon$. Approximating open sets by finite unions of intervals, we deduce the following statement of Littlewood's first principle (in the slightly more general context of Lebesgue–Stieltjes measures).

PROPOSITION 9.1

Let μ be a Lebesgue–Stieltjes measure on \mathbb{R} , and suppose $E \subseteq \mathbb{R}$ is a μ -measurable set with $\mu(E) < \infty$. Then for every $\varepsilon > 0$, there is a finite family of disjoint open intervals $(a_1, b_1), \dots, (a_n, b_n)$ such that

$$\mu \left(E \triangle \bigsqcup_{j=1}^n (a_j, b_j) \right) < \varepsilon.$$

PROOF. Suppose E is μ -measurable with $\mu(E) < \infty$. Let $\varepsilon > 0$. By outer regularity of μ (see Proposition 5.31), let $U \subseteq \mathbb{R}$ be an open set such that $E \subseteq U$ and $\mu(U \setminus E) < \frac{\varepsilon}{2}$. Now by Exercise 1.1, we may write U as a countable union of disjoint open intervals, say $U = \bigsqcup_{j=1}^{\infty} (a_j, b_j)$. Then applying continuity of μ from below, there exists $n \in \mathbb{N}$ such that $\mu \left(\bigsqcup_{j=1}^n (a_j, b_j) \right) > \mu(U) - \frac{\varepsilon}{2}$. Thus,

$$\mu \left(E \triangle \bigsqcup_{j=1}^n (a_j, b_j) \right) \leq \mu(U \setminus E) + \mu \left(U \setminus \bigsqcup_{j=1}^n (a_j, b_j) \right) < \varepsilon.$$

□

One application of Proposition 9.1 that we have already seen in Exercise 5.8 is Steinhaus's theorem.

THEOREM 9.2: STEINHAUS'S THEOREM

Let λ be the Lebesgue measure on \mathbb{R} . Suppose $E \subseteq \mathbb{R}$ is a Lebesgue-measurable set and $\lambda(E) > 0$. Then $E - E = \{x - y : x, y \in E\}$ contains an open interval around 0.

Another application is the Riemann–Lebesgue lemma from Fourier analysis.

THEOREM 9.3: RIEMANN–LEBESGUE LEMMA

Let $f \in L^1(\mathbb{R})$, and define $\widehat{f}(\xi) = \int_{\mathbb{R}} f(x)e^{-2\pi i \xi x} dx$ for $\xi \in \mathbb{R}$. Then $\lim_{|\xi| \rightarrow \infty} \widehat{f}(\xi) = 0$.

PROOF. First suppose $f = \mathbb{1}_I$, where $I = (a, b)$ is an open interval. Then for $\xi \neq 0$,

$$\widehat{f}(\xi) = \int_a^b e^{-2\pi i \xi x} dx = \frac{e^{-2\pi i \xi b} - e^{-2\pi i \xi a}}{-2\pi i \xi}.$$

In particular, $|\widehat{f}(\xi)| \leq \frac{1}{\pi|\xi|}$, so $\widehat{f}(\xi) \rightarrow 0$ as $|\xi| \rightarrow \infty$. By linearity, the conclusion also holds for finite (disjoint) unions of open intervals.

Now suppose $f = \mathbb{1}_E$ for an arbitrary measurable set $E \subseteq \mathbb{R}$ with $\lambda(E) < \infty$. Let $\varepsilon > 0$. By Proposition 9.1, there is a set J that is a finite union of open intervals such that $\lambda(E \Delta J) < \varepsilon$. Hence,

$$|\widehat{f}(\xi)| = \left| \int_{\mathbb{R}} \mathbb{1}_E(x)e^{-2\pi i \xi x} dx \right| \leq \underbrace{\left| \int_{\mathbb{R}} \mathbb{1}_J(x)e^{-2\pi i \xi x} dx \right|}_{\rightarrow 0 \text{ as } |\xi| \rightarrow 0} + \underbrace{\int_{\mathbb{R}} |\mathbb{1}_E(x) - \mathbb{1}_J(x)| dx}_{\lambda(E \Delta J) < \varepsilon}.$$

Therefore, by linearity, the conclusion holds for integrable simple functions.

Finally, suppose $f \in L^1(\mathbb{R})$ is an arbitrary integrable function. Let $\varepsilon > 0$. Then there is a simple function $s \in L^1(\mathbb{R})$ such that $\|f - s\|_1 < \varepsilon$ (see Exercise 3.9). We then have

$$|\widehat{f}(\xi)| \leq \underbrace{|\widehat{s}(\xi)|}_{\rightarrow 0 \text{ as } |\xi| \rightarrow \infty} + \underbrace{|\widehat{(f - s)}(\xi)|}_{\leq \|f - s\|_1 < \varepsilon}.$$

□

Littlewood's first principle has a natural generalization to second countable LCH spaces, which can be used to prove versions of Steinhaus's theorem and the Riemann–Lebesgue lemma in more general topological groups. (We will work with second countable spaces in order to have sets playing the role of intervals.) As preparation for stating a generalized form of Proposition 9.1, let us revisit regularity properties of Radon measures in the context of second countable spaces.

PROPOSITION 9.4

Let X be a locally compact Hausdorff space, and let μ be a Radon measure on X . Then μ is inner regular on all σ -finite sets. That is, if $E \subseteq X$ is a Borel set and $E = \bigcup_{n \in \mathbb{N}} E_n$, where each set E_n is a Borel set with $\mu(E_n) < \infty$, then

$$\mu(E) = \sup\{\mu(K) : K \text{ is compact and } K \subseteq E\}. \quad (9.1)$$

PROOF. We will first handle the case that $\mu(E) < \infty$. Let $\varepsilon > 0$. By outer regularity of μ , there is an open set $U \supseteq E$ such that $\mu(U) < \mu(E) + \frac{\varepsilon}{2}$. Applying outer regularity again, we may find

an open set $V \supseteq U \setminus E$ such that $\mu(V) < \frac{\varepsilon}{2}$. Since μ is inner regular on open sets by assumption, let $K \subseteq U$ be a compact set with $\mu(K) > \mu(U) - \frac{\varepsilon}{2}$. Then $K \setminus V$ is a compact subset of E , and

$$\mu(K \setminus V) \geq \mu(K) - \mu(V) > \mu(U) - \varepsilon \geq \mu(E) - \varepsilon.$$

But ε was arbitrary, so $\mu(E) \leq \sup\{\mu(K) : K \subseteq E \text{ compact}\}$.

Now we handle the general σ -finite case. Write $E = \bigcup_{n \in \mathbb{N}} E_n$ with $E_n \subseteq X$ Borel and $\mu(E_n) < \infty$ for each $n \in \mathbb{N}$. By the finite case above, there exist compact subsets $K_n \subseteq E_n$ with $\mu(E_n \setminus K_n) < 2^{-n}$. For each $N \in \mathbb{N}$, we then have that $\bigcup_{n=1}^N K_n$ is a compact subset of E , and

$$\mu\left(\bigcup_{n=1}^N K_n\right) \geq \mu\left(\bigcup_{n=1}^N E_n\right) - \sum_{n=1}^N \mu(E_n \setminus K_n) > \mu\left(\bigcup_{n=1}^N E_n\right) - 1.$$

Moreover, $\sup_{N \in \mathbb{N}} \mu\left(\bigcup_{n=1}^N E_n\right) = \mu(E) = \infty$, so $\sup_{N \in \mathbb{N}} \mu\left(\bigcup_{n=1}^N K_n\right) = \infty$ as desired. \square

In second countable LCH spaces, locally finite measures are σ -finite, so Proposition 9.4 says that Radon measures are inner regular on all Borel subsets of second countable LCH spaces. (Recall that the definition of a Radon measure only requires inner regularity on the open subsets of X .) When a measure is both inner and outer regular on all Borel sets, we say that it is a *regular* Borel measure. In Proposition 5.31, we showed that every locally finite measure on \mathbb{R} is regular. This extends to second countable LCH spaces.

THEOREM 9.5

Let X be a second countable LCH space or, more generally, an LCH space in which every open set is σ -compact. Then every locally finite Borel measure on X is regular.

PROOF. Let X be an LCH space in which every open set is σ -compact, and let μ be a locally finite Borel measure on X . Since X is σ -compact by assumption, the measure μ is σ -finite, so it suffices by Proposition 9.4 to show that μ is a Radon measure.

Because μ is locally finite, compactly supported continuous functions are μ -integrable, so we may define a positive linear function $I_\mu : C_c(X) \rightarrow \mathbb{C}$ by $I_\mu(f) = \int_X f \, d\mu$. By the Riesz representation theorem, there is a unique Radon measure ν such that $I_\mu(f) = \int_X f \, d\nu$ for every $f \in C_c(X)$. Our goal is thus to show $\mu = \nu$ so that μ is Radon.

CLAIM 1. If U is open, then $\mu(U) = \nu(U)$.

Let $U \subseteq X$ be open. We may write $U = \bigcup_{n \in \mathbb{N}} K_n$ for some compact sets K_n . Let $f_1 \in C_c(X)$ with $K_1 \prec f_1 \prec U$. Then construct inductively $f_n \in C_c(X)$ such that $\bigcup_{j=1}^{n-1} \text{supp}(f_j) \cup K_n \prec f_n \prec U$. Then $0 \leq f_1 \leq f_2 \leq \dots$, and $f_n \rightarrow \mathbb{1}_U$ pointwise. Thus, by the monotone convergence theorem,

$$\mu(U) = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\nu = \nu(U).$$

Write $X = \bigcup_{n \in \mathbb{N}} K_n$ with K_n compact. Then there are open sets V_n such that \overline{V}_n is compact and $K_n \subseteq V_n$ by Lemma 6.7. Let $X_n = \bigcup_{j=1}^n V_j$ so that $X_1 \subseteq X_2 \subseteq \dots$ is an increasing family of open sets with compact closure such that $\bigcup_{n \in \mathbb{N}} X_n = X$.

CLAIM 2. The family $\mathcal{L} = \{E \in \text{Borel}(X) : \mu(E \cap X_n) = \nu(E \cap X_n) \text{ for every } n \in \mathbb{N}\}$ is a λ -system.

The proof is the same as Claim 1 in Corollary 5.14.

The family of open subsets of a topological space is a π -system, so combining Claims 1 and 2 with the π - λ theorem, we conclude that $\mu(E \cap X_n) = \nu(E \cap X_n)$ for every Borel set $E \subseteq X$ and every $n \in \mathbb{N}$. Applying continuity from below, it follows that $\mu = \nu$. \square

COROLLARY 9.6

Let X be a second countable LCH space, and let $\mu : \text{Borel}(X) \rightarrow [0, \infty]$ be a locally finite measure on X . Let \mathcal{U} be a countable base for the topology on X . Suppose $E \in \text{Borel}(X)$ and $\mu(E) < \infty$. Then for any $\varepsilon > 0$, there is a finite collection of basic open sets $U_1, \dots, U_n \in \mathcal{U}$ such that

$$\mu \left(E \triangle \bigcup_{j=1}^n U_j \right) < \varepsilon.$$

PROOF. By Theorem 9.5, the measure μ is Radon. In particular, μ is outer regular, so we may follow the argument in the proof of Proposition 9.1. \square

2. The Second Principle: L^p Functions are Nearly Continuous

There are (at least) two different ways in which L^p functions are “nearly” continuous. One is in terms of the L^p norm.

PROPOSITION 9.7

Let X be an LCH space, and let μ be a Radon measure on X . Then for every $p \in [1, \infty)$, $C_c(X)$ is a dense subspace of $L^p(\mu)$.

PROOF. Let $p \in [1, \infty)$.

STEP 1. $C_c(X) \subseteq L^p(\mu)$.

Let $f \in C_c(X)$, and let $K = \text{supp}(f) \subseteq X$. Then

$$\int_X |f|^p d\mu \leq \max_{x \in K} |f(x)|^p \cdot \mu(K) < \infty$$

by the extreme value theorem and local finiteness of μ , so $f \in L^p(\mu)$.

We will show $C_c(X)$ is dense in $L^p(\mu)$ by approximating successively by more convenient families of functions. Let $\mathcal{S} = \{s : X \rightarrow \mathbb{C} : s \text{ is simple and } \mu(\{s \neq 0\}) < \infty\}$.

STEP 2. \mathcal{S} is dense in $L^p(\mu)$.

Note that a simple function $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j}$ written in standard form has L^p norm satisfying $\|s\|_p^p = \sum_{j=1}^n |c_j|^p \mu(E_j)$. Therefore, $s \in L^p(\mu)$ if and only if $\mu(E_j) < \infty$ for every j such that $c_j \neq 0$. In other words, \mathcal{S} is exactly the collection of simple functions that belong to $L^p(\mu)$.

Let $f \in L^p(\mu)$ be arbitrary. Let $(s_n)_{n \in \mathbb{N}}$ be a sequence of simple functions such that $0 \leq |s_1| \leq |s_2| \leq \dots \leq |f|$ and $s_n \rightarrow f$ a.e. (Such a sequence exists by applying Proposition 3.7 to the positive and negative parts of the real and imaginary parts of f .) Then $|s_n - f|^p \rightarrow 0$ a.e. and $|s_n - f|^p \leq 2^p |f|^p$, so by the dominated convergence theorem, $s_n \rightarrow f$ in $L^p(\mu)$.

STEP 3. $C_c(X)$ is dense in \mathcal{S} (with respect to $\|\cdot\|_p$).

Given a simple function $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j} \in \mathcal{S}$, it suffices by Minkowski's inequality to approximate each of the functions $\mathbb{1}_{E_j}$. Let $E \in \text{Borel}(X)$ with $\mu(E) < \infty$, and let $\varepsilon > 0$. By Proposition 9.4, the measure μ is inner regular on E , and μ is outer regular on all Borel sets, so we may find a compact set K and an open set U such that $K \subseteq E \subseteq U$ and $\mu(U \setminus K) < \varepsilon$. By Urysohn's lemma, let $f \in C_c(X)$ with $K \prec f \prec U$. Then since $f = \mathbb{1}_E = 1$ on K and $f = \mathbb{1}_E = 0$ on $X \setminus U$, we have

$$\int_X |f - \mathbb{1}_E|^p d\mu \leq \mu(U \setminus K) < \varepsilon.$$

□

Another sense in which measurable functions are nearly continuous is provided by Lusin's theorem.

THEOREM 9.8: LUSIN'S THEOREM, VERSION I

Let X be an LCH space, and let Y be a second countable topological space. Let μ be a regular Borel measure on X . Suppose $f : X \rightarrow Y$ is a Borel measurable function. Then for any $\varepsilon > 0$, there is a closed set $E \subseteq X$ such that $\mu(X \setminus E) < \varepsilon$ and $f|_E$ is continuous.

PROOF. Let $\mathcal{U} = \{U_n : n \in \mathbb{N}\}$ be a countable base for the topology on Y . For $n \in \mathbb{N}$, let $B_n = f^{-1}(U_n) \in \text{Borel}(X)$. Since μ is regular, we may find $F_n \subseteq B_n \subseteq G_n$ such that F_n is closed, G_n is open, and $\mu(G_n \setminus F_n) < 2^{-n}\varepsilon$. Let $E = X \setminus \bigcup_{n \in \mathbb{N}} (G_n \setminus F_n)$. Then $\mu(X \setminus E) \leq \sum_{n=1}^{\infty} \mu(G_n \setminus F_n) < \varepsilon$.

CLAIM 1. E is closed.

For each $n \in \mathbb{N}$, the set $G_n \setminus F_n$ is open, so $\bigcup_{n \in \mathbb{N}} (G_n \setminus F_n)$ is open. Therefore, E is the complement of an open set, so E is closed.

CLAIM 2. $f|_E$ is continuous.

Let $g = f|_E : E \rightarrow Y$. Then for each $n \in \mathbb{N}$,

$$g^{-1}(U_n) = B_n \cap E = G_n \cap E$$

is open in E , so g is continuous.

□

It is important to note that Lusin's theorem does NOT say that the set of points of continuity of f has large measure. For example, $1_{\mathbb{Q}} : \mathbb{R} \rightarrow \mathbb{R}$ is a measurable function that is nowhere continuous. However, we can give another interpretation of Lusin's theorem, saying that measurable functions (under some condition) agree with a continuous function outside of a set of small measure.

THEOREM 9.9: LUSIN'S THEOREM, VERSION II

Let X be an LCH space, and let μ be a Radon measure on X . Suppose $f : X \rightarrow \mathbb{C}$ is measurable and $\mu(\{f \neq 0\}) < \infty$. Then for any $\varepsilon > 0$, there exists $g \in C_c(X)$ such that $\mu(\{f \neq g\}) < \varepsilon$.

One of the ingredients for the proof is the Tietze extension theorem (which we will not prove).

THEOREM 9.10: TIETZE EXTENSION THEOREM

Let X be an LCH space and let $K \subseteq X$ be compact and $U \subseteq X$ open such that $K \subseteq U$. If $f : K \rightarrow \mathbb{C}$ is a continuous function, then there exists $g \in C_c(X)$ such that $g = f$ on K and $\text{supp}(g) \subseteq U$.

Note that if $f = 1$ on K , then we recover Urysohn's lemma from the Tietze extension theorem.

PROOF OF LUSIN'S THEOREM, VERSION II. Let $E = \{f \neq 0\}$. Since E has finite measure, μ is regular on E by Proposition 9.4. Therefore, there exists a compact set $K \subseteq E$ and an open set $U \supseteq E$ such that $\mu(U \setminus K) < \frac{\varepsilon}{2}$. Applying version I of Lusin's theorem on the space K , there exists a closed set $C \subseteq K$ with $\mu(K \setminus C) < \frac{\varepsilon}{2}$ such that $f|_C$ is continuous. Then by Tietze's extension theorem, there exists $g \in C_c(X)$ such that $g = f$ on C and $\text{supp}(g) \subseteq U$. Hence, $\{f \neq g\} \subseteq U \setminus C$, so $\mu(\{f \neq g\}) < \varepsilon$. □

3. The Third Principle: Convergent Sequences of Functions are Nearly Uniformly Convergent

Now we turn to the third principle, expressed by Egorov's theorem.

THEOREM 9.11: EGOROV'S THEOREM

Let (X, \mathcal{B}, μ) be a finite measure space, and let Y be a separable metric space. Suppose $f_n : X \rightarrow Y$ is a sequence of measurable functions that converges a.e. to a measurable function $f : X \rightarrow Y$. Then for any $\varepsilon > 0$, there is a set $E \in \mathcal{B}$ such that $\mu(E) < \varepsilon$ and $f_n \rightarrow f$ uniformly on $X \setminus E$.

PROOF. For $k, n \in \mathbb{N}$, let $E_{k,n} = \bigcup_{m \geq n} \{d(f_m, f) \geq \frac{1}{k}\}$.

CLAIM 1. For each $k, n \in \mathbb{N}$, we have $E_{k,n} \in \mathcal{B}$.

It suffices to check that $d(f_m, f)$ is a measurable function for each $m \in \mathbb{N}$. Let $S \subseteq Y$ be a countable dense subset. Then for $y, z \in Y$, we have $d(y, z) = \inf_{s \in S} (d(y, s) + d(z, s))$. For each $s \in S$, let $D_s : Y \rightarrow [0, \infty)$ be the function $D_s(y) = d(y, s)$. The function D_s is continuous, hence Borel measurable. Thus,

$$d(f_m, f) = \inf_{s \in S} (D_s \circ f_m + D_s \circ f)$$

is measurable by Proposition 2.11.

CLAIM 2. For any $k \in \mathbb{N}$, $\mu(\bigcap_{n \in \mathbb{N}} E_{k,n}) = 0$.

The set $\bigcap_{n \in \mathbb{N}} E_{k,n}$ is the set of points $x \in X$ such that $d(f_m, f) \geq \frac{1}{k}$ for infinitely many $m \in \mathbb{N}$. Hence, $\bigcap_{n \in \mathbb{N}} E_{k,n} \subseteq \{x \in X : f_n(x) \not\rightarrow f(x)\}$ is a null set.

Since μ is a finite measure, we may apply continuity from above and let $n_k \in \mathbb{N}$ such that $\mu(E_{k,n_k}) < 2^{-k}\varepsilon$. Let $E = \bigcup_{k \in \mathbb{N}} E_{k,n_k}$. Then $\mu(E) \leq \sum_{k=1}^{\infty} \mu(E_{k,n_k}) < \varepsilon$.

CLAIM 3. $f_n \rightarrow f$ uniformly on $X \setminus E$.

By definition, if $x \notin E_{k,n_k}$, then $d(f_m(x), f(x)) < \frac{1}{k}$ for all $m \geq n_k$. Therefore,

$$\sup_{x \in X \setminus E_{k,n_k}} d(f_m(x), f(x)) < \frac{1}{k}$$

for all $m \geq n_k$ and all $k \in \mathbb{N}$, so $f_n \rightarrow f$ uniformly on $X \setminus E = \bigcap_{k \in \mathbb{N}} (X \setminus E_{k,n_k})$.

□

COROLLARY 9.12: BOUNDED CONVERGENCE THEOREM

Let (X, \mathcal{B}, μ) be a finite measure space. Suppose $f_n : X \rightarrow \mathbb{C}$ is a sequence of measurable functions that converges almost everywhere to a measurable function $f : X \rightarrow \mathbb{C}$. If there exists $M < \infty$ such that $|f_n| \leq M$ a.e., then $f_n \rightarrow f$ in $L^1(\mu)$. In particular,

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

PROOF. This is a special case of the dominated convergence theorem: since $\mu(X) < \infty$, the function $g(x) = M$ is integrable. We will give a proof using Egorov's theorem (which importantly does not rely on the dominated convergence theorem).

Let $\varepsilon > 0$. By Egorov's theorem, let $E \in \mathcal{B}$ such that $\mu(E) < \varepsilon$ and $f_n \rightarrow f$ uniformly on $X \setminus E$. Then

$$\int_X |f_n - f| \, d\mu = \int_E |f_n - f| \, d\mu + \int_{X \setminus E} |f_n - f| \, d\mu.$$

The first term $\int_E |f_n - f| \, d\mu$ can be bounded using the triangle inequality by $2M\mu(E) < 2M\varepsilon$. The second term is bounded by

$$\sup_{x \in X \setminus E} |f_n(x) - f(x)| \cdot \underbrace{\mu(X \setminus E)}_{\leq \mu(X) < \infty} \xrightarrow{n \rightarrow \infty} 0.$$

Hence, $\limsup_{n \rightarrow \infty} \|f_n - f\|_1 \leq 2M\varepsilon$. But $\varepsilon > 0$ was arbitrary, so $f_n \rightarrow f$ in $L^1(\mu)$. □

Chapter Notes

Some authors prove Lusin's theorem as a consequence of Egorov's theorem. For example, a special case of version I of Lusin's theorem appears as Exercise 44 in [2, Section 2.4], with a hint

to use Egorov's theorem, and version II is proved using Egorov's theorem in [2, Theorem 7.10]. The proofs provided in these lecture notes are shorter and more direct than the deduction from Egorov's theorem.

Exercises

9.1 Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a Lebesgue-measurable function and $f(x + y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$.

- (a) Use Lusin's theorem to show that f is continuous at 0.
- (b) Conclude that $f(x) = xf(1)$ for every $x \in \mathbb{R}$.

9.2 Let X be an LCH space, and let μ be a Radon measure on X . Let $f : X \rightarrow \mathbb{R}$ be integrable. In this exercise, you will prove the Vitali–Carathéodory theorem: for any $\varepsilon > 0$, there exist $u, v : X \rightarrow \mathbb{R}$ such that u is upper semicontinuous and bounded above, v is lower semicontinuous and bounded below, $u \leq f \leq v$, and $\int_X (v - u) \, d\mu < \varepsilon$.

- (a) Justify that one can assume without loss of generality that $f \geq 0$.
- (b) Show that there are measurable sets $(E_n)_{n \in \mathbb{N}}$ and nonnegative real numbers $c_n \geq 0$ such that $f = \sum_{n=1}^{\infty} c_n \mathbb{1}_{E_n}$.
- (c) Find appropriate compact sets $(K_n)_{n \in \mathbb{N}}$ and open sets $(U_n)_{n \in \mathbb{N}}$ such that $u = \sum_{n=1}^N c_n \mathbb{1}_{K_n}$ and $v = \sum_{n=1}^{\infty} c_n \mathbb{1}_{U_n}$ have the desired properties for carefully chosen $N \in \mathbb{N}$.

9.3 Let (X, \mathcal{B}, μ) be a finite measure space. We say that a sequence of functions $(f_n)_{n \in \mathbb{N}}$ converges in measure to f if for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that if $n \geq N$, then $\mu(\{|f - f_n| > \varepsilon\}) < \varepsilon$. Show that the following are equivalent:

- (i) $f_n \rightarrow f$ in measure
- (ii) every subsequence $(f_{n_k})_{k \in \mathbb{N}}$ has a further subsequence $(f_{n_{k_l}})_{l \in \mathbb{N}}$ that converges to f a.e.

CHAPTER 10

Differentiation of Measures

1. Signed and Complex Measures

DEFINITION 10.1

Let (X, \mathcal{B}) be a measurable space. A *signed measure* is a countably additive function $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ with $\mu(\emptyset) = 0$. A *complex measure* is a countably additive function $\mu : \mathcal{B} \rightarrow \mathbb{C}$ with $\mu(\emptyset) = 0$.

REMARK. Since the value $\infty - \infty$ is undefined, a signed measure may only take at most one of the values ∞ or $-\infty$.

The main motivation for introducing the more general notion of signed and complex measures is that they arise quite naturally when integrating functions.

EXAMPLE 10.2

Let (X, \mathcal{B}, μ) be a measure space.

SIGNED MEASURE. Let $f : X \rightarrow [-\infty, \infty]$ be a measurable function, and suppose at least one of $\int_X f^+ d\mu$ or $\int_X f^- d\mu$ is finite, so that $\int_X f d\mu$ can be defined as $\int_X f^+ d\mu - \int_X f^- d\mu$. Then $\nu(E) = \int_E f d\mu$ is a signed measure.

COMPLEX MEASURE. If $f \in L^1(\mu)$, then $\nu(E) = \int_E f d\mu$ defines a complex measure.

Signed and complex measures retain the continuity properties of (positive) measures. Note, however, that we may lose monotonicity and countable subadditivity.

PROPOSITION 10.3

Let (X, \mathcal{B}) be a measurable space, and let μ be a signed or complex measure on (X, \mathcal{B}) .

- (1) **CONTINUITY FROM BELOW:** if $E_1 \subseteq E_2 \subseteq \dots \in \mathcal{B}$, then $\mu(\bigcup_{n \in \mathbb{N}} E_n) = \lim_{n \rightarrow \infty} \mu(E_n)$.
- (2) **CONTINUITY FROM ABOVE:** if $E_1 \supseteq E_2 \supseteq \dots \in \mathcal{B}$ and $|\mu(E_1)| < \infty$, then $\mu(\bigcap_{n \in \mathbb{N}} E_n) = \lim_{n \rightarrow \infty} \mu(E_n)$.

PROOF. The proof is the same as for positive measures (see Proposition 2.15). For completeness, we repeat the argument here to see that positivity of the measure is inconsequential.

(1) Let $E'_1 = E_1$ and $E'_n = E_n \setminus E_{n-1}$ for $n \geq 2$. For convenience, we will set $E_0 = \emptyset$ so that we also have $E'_1 = E_1 \setminus E_0$. Then

$$\mu\left(\bigcup_{n \in \mathbb{N}} E_n\right) = \mu\left(\bigsqcup_{n \in \mathbb{N}} E'_n\right) = \sum_{n \in \mathbb{N}} \mu(E'_n) \stackrel{(*)}{=} \sum_{n \in \mathbb{N}} (\mu(E_n) - \mu(E_{n-1})) \stackrel{(**)}{=} \lim_{n \rightarrow \infty} \mu(E_n).$$

The step (*) uses additivity of μ , and (**) comes from the telescoping of the sum.

(2) Define a new sequence $A_n = E_1 \setminus E_n$. Then $\emptyset = A_1 \subseteq A_2 \subseteq \dots$, so

$$\mu \left(\bigcup_{n \in \mathbb{N}} A_n \right) = \lim_{n \rightarrow \infty} \mu(A_n)$$

by (1). But $\bigcup_{n \in \mathbb{N}} A_n = E_1 \setminus \bigcap_{n \in \mathbb{N}} E_n$, so

$$\mu(E_1) - \mu \left(\bigcap_{n \in \mathbb{N}} E_n \right) = \mu \left(\bigcup_{n \in \mathbb{N}} A_n \right) = \lim_{n \rightarrow \infty} \mu(A_n) = \mu(E_1) - \lim_{n \rightarrow \infty} \mu(E_n),$$

whence we deduce that (2) holds, since $|\mu(E_1)| < \infty$. □

DEFINITION 10.4

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure. A set $E \in \mathcal{B}$ is

- *null* if for every $F \in \mathcal{B}$, if $F \subseteq E$, then $\mu(F) = 0$;
- *positive* if for every $F \in \mathcal{B}$, if $F \subseteq E$, then $\mu(F) \geq 0$;
- *negative* if for every $F \in \mathcal{B}$, if $F \subseteq E$, then $\mu(F) \leq 0$.

THEOREM 10.5: HAHN DECOMPOSITION THEOREM

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure. There exists a partition $X = P \sqcup N$ such that $P, N \in \mathcal{B}$, P is a positive subset, and N is a negative subset. Moreover, if $X = P' \sqcup N'$ is another such partition, then $P \Delta P' = N \Delta N'$ is a null set.

DEFINITION 10.6

The ordered pair (P, N) in the conclusion of the Hahn decomposition theorem is called a *Hahn decomposition* of μ .

Before proving the Hahn decomposition theorem, we discuss a useful corollary. This requires one more definition.

DEFINITION 10.7

Let μ and ν be positive, signed, or complex measures on a measurable space (X, \mathcal{B}) . We say that μ and ν are *mutually singular*, denoted $\mu \perp \nu$, if there is a measurable partition $X = E \sqcup F$ such that E is ν -null and F is μ -null.

THEOREM 10.8: JORDAN DECOMPOSITION THEOREM

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure. There is a unique decomposition $\mu = \mu^+ - \mu^-$ as a sum of mutually singular positive measures.

PROOF. Let (P, N) be a Hahn decomposition by Theorem 10.5. Define $\mu^+(E) = \mu(E \cap P)$ and $\mu^-(E) = -\mu(E \cap N)$. Then μ^+ and μ^- are mutually singular positive measures, and $\mu = \mu^+ - \mu^-$.

To see uniqueness, suppose μ_1 and μ_2 are mutually singular positive measures such that $\mu = \mu_1 - \mu_2$. Write $X = A \sqcup B$ such that A is μ_2 -null and B is μ_1 -null. We claim that (A, B) is another Hahn decomposition. Indeed, for $E \in \mathcal{B}$, $E \subseteq A$, we have $\mu_2(E) = 0$, so $\mu(E) = \mu_1(E) \geq 0$. That is, A is μ -positive. Similarly, for $E \in \mathcal{B}$, $E \subseteq B$, we have $\mu(E) = -\mu_2(E) \leq 0$, so B is μ -negative. Thus, by the Hahn decomposition theorem, $A \triangle P = B \triangle N$ is μ -null. For any $E \in \mathcal{B}$, we then have

$$\mu^+(E) = \mu(E \cap P) = \mu(E \cap A) = \mu_1(E) \quad \text{and} \quad \mu^-(E) = -\mu(E \cap N) = -\mu(E \cap B) = \mu_2(E).$$

□

DEFINITION 10.9

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure with Jordan decomposition $\mu = \mu^+ - \mu^-$. The measure μ^+ is called the *positive variation* of μ and μ^- the *negative variation* of μ . The sum $\mu^+ + \mu^-$ is called the *total variation* of μ and is denoted by $|\mu|$.

PROPOSITION 10.10

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure with total variation $|\mu|$. Then for any $E \in \mathcal{B}$,

$$\begin{aligned} |\mu|(E) &= \inf \{ \nu(E) : \nu \text{ is a measure and } |\nu(F)| \leq \nu(F) \text{ for all } F \in \mathcal{B} \} \\ &= \sup \left\{ \sum_{n=1}^{\infty} |\mu(E_n)| : E = \bigsqcup_{n \in \mathbb{N}} E_n \right\}. \end{aligned}$$

PROOF. Exercise 10.1. □

We now turn to proving the Hahn decomposition theorem. Let us start with a criterion for a pair (P, N) to be a Hahn decomposition.

LEMMA 10.11

Let \mathcal{A} be an algebra of subsets of a set X , and let $\mu : \mathcal{A} \rightarrow (-\infty, \infty]$ be a finitely additive function on \mathcal{A} . Let $X = P \sqcup N$ with $P, N \in \mathcal{A}$. The following are equivalent:

- (i) (P, N) is a Hahn decomposition;
- (ii) $\mu(N) = \inf_{A \in \mathcal{A}} \mu(A)$.

PROOF. (i) \implies (ii). Suppose (P, N) is a Hahn decomposition. Let $A \in \mathcal{A}$. Then

$$\mu(A) = \underbrace{\mu(A \cap P)}_{\geq 0} + \underbrace{\mu(A \cap N)}_{\leq 0} \geq \mu(A \cap N) + \underbrace{\mu(N \setminus A)}_{\leq 0} = \mu(N).$$

So (ii) holds.

(ii) \implies (i). Suppose (ii) holds. We need to check that N is negative and P is positive.

Suppose $A \in \mathcal{A}$ and $A \subseteq N$. Then $\mu(N) \leq \mu(N \setminus A) = \mu(N) - \mu(A)$, so $\mu(A) \leq 0$. Thus, N is μ -negative.

Now suppose $A \in \mathcal{A}$ and $A \subseteq P$. Then $\mu(N) \leq \mu(N \sqcup A) = \mu(N) + \mu(A)$, so $\mu(A) \geq 0$. That is, P is μ -positive. □

Now we can prove the Hahn decomposition theorem.

PROOF OF HAHN DECOMPOSITION THEOREM. We will first give a construction of a Hahn decomposition and then show μ -essential uniqueness.

STEP 1. Existence.

Since μ assumes at most one of the values ∞ or $-\infty$, let us assume that $\mu(E) > -\infty$ for every $E \in \mathcal{B}$ (if not, we may work with the measure $-\mu$ instead). Let $a = \inf\{\mu(E) : E \in \mathcal{B}\}$. By Lemma 10.11, it suffices to find a set $N \in \mathcal{B}$ such that $\mu(N) = a$.

Let $(E_n)_{n \in \mathbb{N}}$ be a sequence of measurable sets such that $\mu(E_n) \rightarrow a$. We are going to replace this sequence by a better-behaving sequence from which we can extract the set N as a limit. The $\mathcal{A}_n = \mathcal{A}(E_1, \dots, E_n)$ be the algebra generated by E_1, \dots, E_n . Since \mathcal{A}_n is a finite collection of sets, let $A_n \in \mathcal{A}_n$ such that $\mu(A_n) = \min\{\mu(A) : A \in \mathcal{A}_n\}$. By Lemma 10.11, the pair $(X \setminus A_n, A_n)$ is a Hahn decomposition for $\mu|_{\mathcal{A}_n}$.

By construction, the algebras $(\mathcal{A}_n)_{n \in \mathbb{N}}$ are nested: $\mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$. Therefore, $A_1, \dots, A_n \in \mathcal{A}_n$. Let $N_m = \bigcap_{k \geq m} A_k$ and $N_m^n = \bigcap_{k=m}^n A_k$. For each $m \in \mathbb{N}$, the sequence of sets $(N_m^n)_{n \geq m}$ decreases to N_m . Moreover, the set $N_m^n = A_m$ has finite measure, so by continuity from above, $\mu(N_m^n) \rightarrow \mu(N_m)$ as $n \rightarrow \infty$. For $n > m$,

$$\mu(N_m^{n-1}) = \mu(N_m^n) + \mu((X \setminus A_n) \cap N_m^{n-1}) \geq \mu(N_m^n),$$

where we have used that $N_m^{n-1} \in \mathcal{A}_n$ and $X \setminus A_n$ is μ -positive on the algebra \mathcal{A}_n . Thus,

$$\mu(E_m) \geq \mu(A_m) = \mu(N_m^n) \geq \mu(N_m^{n+1}) \geq \mu(N_m^{n+2}) \geq \dots,$$

so $a \leq \mu(N_m) \leq \mu(E_m)$. Therefore, $\lim_{m \rightarrow \infty} \mu(N_m) = a$. But the sequence $(N_m)_{m \in \mathbb{N}}$ is increasing, so letting $N = \bigcup_{m \in \mathbb{N}} N_m$, we have $\mu(N) = a$ by continuity from below.

STEP 2. Essential Uniqueness.

Suppose (P', N') is another Hahn decomposition. Then $N' \triangle N = P' \triangle P = (N \cap P') \sqcup (N' \cap P)$. The set $N \cap P'$ is both negative (since it is a subset of the negative set N) and positive (since it is a subset of the positive set P'). Therefore, $N \cap P'$ is a null set, and similarly for $N' \cap P$.

□

2. Radon–Nikodym Derivatives

DEFINITION 10.12

Let (X, \mathcal{B}, μ) be a (positive) measure space. Let ν be a signed or complex measure on (X, \mathcal{B}) . We say that ν is *absolutely continuous* with respect to μ , denoted $\nu \ll \mu$, if every μ -null set is ν -null. Two positive measures μ and ν are *equivalent*, written $\mu \approx \nu$, if $\mu \ll \nu$ and $\nu \ll \mu$.

We have already seen the prototypical example of an absolutely continuous measure.

EXAMPLE 10.13

Let (X, \mathcal{B}, μ) be a measure space, and let $f \in L^1(\mu)$. Then $\nu : \mathcal{B} \rightarrow \mathbb{C}$ defined by $\nu(E) = \int_E f \, d\mu$ is a complex measure, and $\nu \ll \mu$.

THEOREM 10.14: LEBESGUE DECOMPOSITION THEOREM

Let (X, \mathcal{B}) be a measurable space. Let $\mu, \nu : \mathcal{B} \rightarrow [0, \infty]$ be two positive measures on (X, \mathcal{B}) , and suppose ν is s -finite. Then there is a unique decomposition $\nu = \nu_a + \nu_s$ as a sum of positive measures such that $\nu_a \ll \mu$ and $\nu_s \perp \mu$.

PROOF. We prove existence and then uniqueness.

Write $\nu = \sum_{n=1}^{\infty} \nu_n$ with $\nu_n(X) < \infty$, and define $\tilde{\nu} = \sum_{n=1}^{\infty} 2^{-n} \frac{\nu_n}{\nu_n(X)+1}$. Then $\tilde{\nu}(X) < 1$, and $\tilde{\nu} \approx \nu$. Let (P, N) be a Hahn decomposition for the measure $\infty \cdot \mu - \tilde{\nu}$, and define $\nu_a(E) = \nu(E \cap P)$ and $\nu_s(E) = \nu(E \cap N)$.

CLAIM 1. $\nu_a \ll \mu$.

Suppose $\mu(E) = 0$. Then since $E \cap P$ is a positive set for $\infty \cdot \mu - \tilde{\nu}$, we have $0 \leq -\tilde{\nu}(E \cap P)$, so $\tilde{\nu}(E \cap P) = 0$. But $\nu \ll \tilde{\nu}$, so $\nu_a(E) = \nu(E \cap P) = 0$.

CLAIM 2. $\nu_s \perp \mu$.

By construction P is a ν_s -null set, so it suffices to show $N = X \setminus P$ is μ -null. But N is negative for $\infty \cdot \mu - \tilde{\nu}$, so in particular, $\infty \cdot \mu(N) - \tilde{\nu}(N) \leq 0$. That is, $\infty \cdot \mu(N) \leq \tilde{\nu}(N) < \infty$. Hence, $\mu(N) = 0$.

Now let us check uniqueness. Suppose $\nu = \alpha + \beta$ is another decomposition with $\alpha \ll \mu$ and $\beta \perp \mu$. Since $\nu_s \perp \mu$ and $\beta \perp \mu$, we may find sets $A, B \in \mathcal{B}$ such that $\mu(A) = \mu(B) = 0$ and $\nu_s(X \setminus A) = \beta(X \setminus B) = 0$. Then $C = A \cup B$ is a μ -null set, so $\nu_a(C) = \alpha(C) = 0$. Therefore, for $E \in \mathcal{B}$,

$$\nu_a(E) = \nu_a(\underbrace{E \cap C}_{\subseteq C}) + \nu_a(E \setminus C) = \nu_a(E \setminus C) + \nu_s(\underbrace{E \setminus C}_{\subseteq X \setminus A}) = \nu(E \setminus C)$$

and

$$\nu_s(E) = \nu_s(E \cap C) + \nu_s(\underbrace{E \setminus C}_{\subseteq X \setminus A}) = \nu_s(E \cap C) + \nu_a(\underbrace{E \cap C}_{\subseteq C}) = \nu(E \cap C).$$

Similarly, $\alpha(E) = \nu(E \setminus C)$ and $\beta(E) = \nu(E \cap C)$, so $\alpha = \nu_a$ and $\beta = \nu_s$. □

THEOREM 10.15: RADON–NIKODYM THEOREM

Let (X, \mathcal{B}, μ) be a σ -finite measure space. Let $\nu : \mathcal{B} \rightarrow [0, \infty]$ be a positive measure, and suppose $\nu \ll \mu$. Then there exists a measurable function $f : X \rightarrow [0, \infty]$ such that

$$\nu(E) = \int_E f \, d\mu$$

for every $E \in \mathcal{B}$.

Before proving the Radon–Nikodym theorem, we make several remarks and observations. First, the conclusion $\nu(E) = \int_E f \, d\mu$ for every $E \in \mathcal{B}$ is often abbreviated as $d\nu = f \, d\mu$, which is justified by the following observation:

PROPOSITION 10.16

Let (X, \mathcal{B}, μ) be a measurable space, and let $\nu : \mathcal{B} \rightarrow [0, \infty]$ be the measure defined by $\nu(E) = \int_E f \, d\mu$ for $E \in \mathcal{B}$. Then $\int_X g \, d\nu = \int_X g f \, d\mu$ for every measurable function $g : X \rightarrow [0, \infty]$ and every ν -integrable function $g : X \rightarrow \mathbb{C}$.

PROOF. If $g = \mathbb{1}_E$ for a measurable set $E \in \mathcal{B}$, then $\int_X g \, d\nu = \int_X g f \, d\mu$ by the definition of the measure ν . Thus, $\int_X g \, d\nu = \int_X g f \, d\mu$ for nonnegative simple functions by linearity, then for nonnegative measurable functions by the monotone convergence theorem, and then for integrable functions by another application of linearity. \square

Next, the conclusion of the Radon–Nikodym theorem holds also for signed and complex measures.

COROLLARY 10.17: RADON–NIKODYM THEOREM FOR SIGNED AND COMPLEX MEASURES

Let (X, \mathcal{B}, μ) be a σ -finite measure space.

- (1) If $\nu : \mathcal{B} \rightarrow [-\infty, \infty]$ is a signed measure and $\nu \ll \mu$, then there exists a measurable function $f : X \rightarrow [-\infty, \infty]$ such that at least one of the quantities $\int_X f^+ \, d\mu$ or $\int_X f^- \, d\mu$ is finite and $d\nu = f \, d\mu$.
- (2) If $\nu : \mathcal{B} \rightarrow \mathbb{C}$ is a complex measure and $\nu \ll \mu$, then there exists a μ -integrable function $f : X \rightarrow \mathbb{C}$ such that $d\nu = f \, d\mu$.

PROOF. (1) Let $\nu = \nu^+ - \nu^-$ be the Jordan decomposition of ν . Then $\nu^+, \nu^- \ll \mu$, so there exist measurable functions $f^+ : X \rightarrow [0, \infty]$ and $f^- : X \rightarrow [0, \infty]$ such that $\nu^+ = f^+ \, d\mu$ and $\nu^- = f^- \, d\mu$. Taking $f = f^+ - f^-$ completes the proof.

(2) Write $\nu = \nu_1 + i\nu_2$ as a combination of finite signed measures ν_1 and ν_2 and apply (1) to find μ -integrable functions $f_1, f_2 : X \rightarrow \mathbb{R}$ such that $d\nu_j = f_j \, d\mu$. Then let $f = f_1 + i f_2$. \square

REMARK. We have not defined integration against signed and complex measures, so the notation $d\nu = f \, d\mu$ in Corollary 10.17 is not immediately justified by Proposition 10.16. To fill in this gap, we may define, for a signed measure ν , the integral $\int_X g \, d\nu = \int_X g \, d\nu^+ - \int_X g \, d\nu^-$ whenever both quantities are finite. Then decomposing a complex measure $\nu = \nu_1 + i\nu_2$ as a combination of signed measures, we define $\int_X g \, d\nu = \int_X g \, d\nu_1 + i \int_X g \, d\nu_2$ when both quantities are finite.

Finally, the function f appearing in the conclusion of the Radon–Nikodym theorem is μ -essentially unique. That is, if g is another function satisfying $d\nu = g \, d\mu$, then $f = g$ μ -a.e. We call this μ -essentially unique function the Radon–Nikodym derivative. To be precise:

DEFINITION 10.18

Let (X, \mathcal{B}, μ) be a measure space, and let ν be a positive, signed, or complex measure. If there is a measurable function f such that $d\nu = f \, d\mu$, then f is called the *Radon–Nikodym derivative of ν with respect to μ* , denoted by $\frac{d\nu}{d\mu} = f$ (a.e.).

Let us now turn to the proof of the Radon–Nikodym theorem.

PROOF OF RADON–NIKODYM THEOREM. We first deal with the case that μ is finite.

CASE 1. μ is a finite measure.

We construct the function f through its level sets by considering the Hahn decomposition of the signed measure $\nu - r\mu$ for $r > 0$. In order to ensure measurability in the end, we will consider only rational values of r .

For $r \in \mathbb{Q}$, $r > 0$, let (P_r, N_r) be a Hahn decomposition of $\nu - r\mu$, and let $f_r = r \cdot 1_{P_r}$. Define $f : X \rightarrow [0, \infty]$ by $f = \sup_{r \in \mathbb{Q}, r > 0} f_r$. Then f is measurable, and the pair $(\{f > c\}, \{f \leq c\})$ is a Hahn decomposition of $\nu - c\mu$ for every $c \in [0, \infty)$. Let us prove $\nu(E) = \int_E f \, d\mu$ for every $E \in \mathcal{B}$.

Let $t > 1$, and let $F_k = \{t^k < f \leq t^{k+1}\}$ for $k \in \mathbb{Z}$. Then F_k is positive for $\nu - t^k \mu$ and negative for $\nu - t^{k+1} \mu$. Therefore,

$$t^{-1} \int_{E \cap F_k} f \, d\mu \leq t^k \mu(E \cap F_k) \leq \nu(E \cap F_k) \leq t^{k+1} \mu(E \cap F_k) \leq t \int_{E \cap F_k} f \, d\mu.$$

Let $F_\infty = \{f = \infty\}$. Then

$$\int_{E \cap F_\infty} f \, d\mu = \infty \cdot \mu(E \cap F_\infty).$$

Moreover, $\nu - c\mu$ is positive on F_∞ for every $c > 0$, so $\nu \geq \infty \cdot \mu$ on F_∞ . On the other hand, $\nu \ll \mu$, so $\nu \leq \infty \cdot \mu$. Hence, $\int_{E \cap F_\infty} f \, d\mu = \nu(E \cap F_\infty)$. Now, summing over $k \in \mathbb{Z} \cup \{\infty\}$,

$$t^{-1} \int_E f \, d\mu \leq \nu(E) \leq t \int_E f \, d\mu.$$

Letting $t \rightarrow 1$, we conclude $\nu(E) = \int_E f \, d\mu$ by the squeeze theorem.

Now we prove the general case.

CASE 2. μ is σ -finite.

Let $X = \bigcup_{n \in \mathbb{N}} X_n$ with $\mu(X_n) < \infty$. We may assume that the sets $(X_n)_{n \in \mathbb{N}}$ are pairwise disjoint. By Case 1, there are measurable functions $f_n : X_n \rightarrow [0, \infty]$ such that $\nu(E) = \int_E f_n \, d\mu$ for $E \in \mathcal{B}$, $E \subseteq X_n$. We then define $f(x) = f_n(x)$ if $x \in X_n$.

□

Using the Radon–Nikodym theorem, we can give an additional property that make the class of σ -finite measures quite natural.

PROPOSITION 10.19

Let (X, \mathcal{B}, μ) be a measure space. The following are equivalent:

- (i) μ is σ -finite;
- (ii) there exists a finite measure ν such that $\mu \approx \nu$;
- (iii) there exists a σ -finite measure ν such that $\mu \ll \nu$.

PROOF. Problem 10 in the exam study guide. □

The Lebesgue decomposition theorem and Radon–Nikodym theorem are often combined into a single theorem, the Lebesgue–Radon–Nikodym theorem, which can be stated as follows.

COROLLARY 10.20: LEBESGUE–RADON–NIKODYM THEOREM

Let (X, \mathcal{B}) be a measurable space. Let $\mu, \nu : \mathcal{B} \rightarrow [0, \infty]$ be measures on (X, \mathcal{B}) . Suppose μ is σ -finite and ν is s -finite. Then there is a unique singular measure $\nu_s \perp \mu$ and a μ -essentially unique measurable function $f : X \rightarrow [0, \infty]$ such that

$$\nu(E) = \int_E f \, d\mu + \nu_s(E)$$

for every $E \in \mathcal{B}$.

3. Complex Borel Measures on \mathbb{R} and their Derivatives

Suppose μ is a Lebesgue–Stieltjes measure on \mathbb{R} . We say that μ is *absolutely continuous* (respectively, *singular*) if it is absolutely continuous (resp., singular) with respect to the Lebesgue measure λ . Given the distribution function F_μ , how can we determine the Lebesgue decomposition $\mu = \mu_a + \mu_s$ into absolutely continuous and singular measures? And if $\mu \ll \lambda$, how does the Radon–Nikodym derivative $\frac{d\mu}{d\lambda}$ relate to F_μ ?

Before answering these questions, we slightly expand the objects under consideration. If $\mu : \text{Borel}(\mathbb{R}) \rightarrow \mathbb{R}$ is a finite signed measure, then we can write $\mu = \mu^+ - \mu^-$ and the distribution function for μ will be a difference of increasing right-continuous functions $F_\mu = F_{\mu^+} - F_{\mu^-}$. How can such functions be characterized? Inspired by the second expression for the total variation measure in Proposition 10.10, we make the following definition:

DEFINITION 10.21

Let $f : \mathbb{R} \rightarrow \mathbb{C}$. The *total variation* of f is the function

$$T_f(x) = \sup \left\{ \sum_{j=0}^{n-1} |f(x_{j+1}) - f(x_j)| : n \in \mathbb{N}, -\infty < x_0 < x_1 < \dots < x_n \leq x \right\}$$

We say that f is of *bounded variation* if $\text{Var}(f) = \sup_{x \in \mathbb{R}} T_f(x) < \infty$.

NOTATION. We denote the class of functions of bounded variation by $\text{BV}(\mathbb{R})$.

We make a few observations. If $f \in \text{BV}(\mathbb{R})$ is real-valued, then $T_f(b) - T_f(a) \geq |f(b) - f(a)|$, so $T_f \pm f$ is an increasing function. We can thus write $f = \frac{1}{2}(T_f + f) - \frac{1}{2}(T_f - f)$ as a difference of increasing bounded functions. On the other hand, if f_1 and f_2 are bounded and increasing, then $T_{f_1 - f_2} \leq f_1 + f_2$, so $T_{f_1 - f_2}$ is of bounded variation. Thus, we have the following characterization of when a function can be expressed as a difference of increasing bounded functions.

PROPOSITION 10.22

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ can be expressed as $f = f_1 - f_2$ with f_1, f_2 bounded and increasing if and only if $f \in \text{BV}(\mathbb{R})$.

We can combine this observation with the Hahn decomposition theorem and the classification of positive Borel measures on \mathbb{R} to characterize complex Borel measures on \mathbb{R} . Define the class of “normalized” functions of bounded variation by

$$\text{NBV}(\mathbb{R}) = \left\{ F \in \text{BV}(\mathbb{R}) : F \text{ is right continuous and } \lim_{x \rightarrow -\infty} F(x) = 0 \right\}.$$

THEOREM 10.23

Let $F \in \text{NBV}(\mathbb{R})$. Then there is a unique complex Borel measure $\mu_F : \text{Borel}(\mathbb{R}) \rightarrow \mathbb{C}$ such that $F(x) = \mu((-\infty, x])$ for every $x \in \mathbb{R}$. Moreover, the total variation of μ_F is the measure $|\mu_F| = \mu_{T_F}$.

REMARK. When we defined Lebesgue–Stieltjes measures, we used distribution functions normalized to have $F(0) = 0$. The reason for making such a choice was that the distribution function for a Lebesgue–Stieltjes measure may be unbounded; in particular, if μ is a Lebesgue–Stieltjes measure, it is possible that $\mu((-\infty, x]) = \infty$ for every $x \in \mathbb{R}$. By contrast, complex measures take only finite values, so it is possible to take the (more convenient) normalization $F(-\infty) = 0$. This is also the standard practice in probability theory, where the function $F(x) = \mu((-\infty, x])$ is called the *cumulative distribution function* of μ .

A deep theorem in analysis that we will not have time to cover in this course is the Lebesgue differentiation theorem, which states the following.

THEOREM 10.24: LEBESGUE DIFFERENTIATION THEOREM

Let $\mu, \nu : \text{Borel}(\mathbb{R}) \rightarrow [0, \infty]$ be locally finite Borel measures. Then the limit

$$\lim_{\delta \rightarrow 0^+} \frac{\nu((x - \delta, x + \delta))}{\mu((x - \delta, x + \delta))}$$

exists for μ -a.e. $x \in \mathbb{R}$. Moreover, if $\nu = \nu_a + \nu_s$ is the Lebesgue decomposition of ν with respect to μ , then the limit is equal μ -a.e. to the Radon–Nikodym derivative $\frac{d\nu_a}{d\mu}$.

REMARK. The term “Lebesgue differentiation theorem” is sometimes used to refer to the following special case (corresponding to $\mu = \lambda$ and $d\nu = f d\lambda$). If $f \in L^1(\mathbb{R})$, then for a.e. $x \in \mathbb{R}$, $\lim_{\delta \rightarrow 0^+} \int_{x-\delta}^{x+\delta} f(t) dt = f(x)$.

This leads to the following description of the Lebesgue decomposition of a complex Borel measure on \mathbb{R} .

THEOREM 10.25

Let $F \in \text{NBV}(\mathbb{R})$. Then F is differentiable a.e. (with respect to the Lebesgue measure λ). Moreover, if $\mu_F = \mu_a + \mu_s$ is the Lebesgue decomposition of μ_F with respect to λ , then $F' = \frac{d\mu_a}{d\lambda}$ a.e.

If μ is absolutely continuous, then we may write $F_\mu(x) = \int_{-\infty}^x \frac{d\mu}{d\lambda}(t) dt = \int_{-\infty}^x F'(t) dt$. In Exercise 3.5, you showed the following property: for any $\varepsilon > 0$, there exists $\delta > 0$ such that if $E \in \text{Borel}(\mathbb{R})$ and $\lambda(E) < \delta$, then $|\mu|(E) = \int_E |F'(t)| dt < \varepsilon$. Inspired by this characterization, we define absolute continuity of functions.

DEFINITION 10.26

A function $F : \mathbb{R} \rightarrow \mathbb{C}$ is *absolutely continuous* if for every $\varepsilon > 0$, there exists $\delta > 0$ with the following property: if $a < b_1 \leq a_2 < b_2 \leq \dots \leq a_n < b_n$ and $\sum_{i=1}^n (b_i - a_i) < \delta$, then $\sum_{i=1}^n |F(b_i) - F(a_i)| < \varepsilon$.

We then have the following theorem.

THEOREM 10.27: FUNDAMENTAL THEOREM OF CALCULUS FOR THE LEBESGUE INTEGRAL

- (1) Suppose $F : \mathbb{R} \rightarrow \mathbb{C}$ is absolutely continuous. Then F is differentiable a.e. and $F(b) - F(a) = \int_a^b F'(x) dx$ for every $a, b \in \mathbb{R}$.
- (2) If $f \in L^1(\mathbb{R})$, then the function $F(x) = \int_{-\infty}^x f(t) dt$ is absolutely continuous, and $F'(x) = f(x)$ a.e.

4. The Dual of L^p

As another application of the Radon–Nikodym theorem, we will show that L^p and L^q are dual for conjugate exponents $p, q \in (1, \infty)$.

THEOREM 10.28: RIESZ–FRÉCHET THEOREM

Let (X, \mathcal{B}, μ) be a measure space. Let $p, q \in (1, \infty)$ with $\frac{1}{p} + \frac{1}{q} = 1$. For $g \in L^q(\mu)$, define $\varphi_g : L^p(\mu) \rightarrow \mathbb{C}$ by $\varphi_g(f) = \int_X fg d\mu$. Then the map $g \mapsto \varphi_g$ is an isometric isomorphism between $L^q(\mu)$ and $L^p(\mu)^*$.

PROOF. By Hölder's inequality, $\|fg\|_1 \leq \|f\|_p \|g\|_q$, so fg is an integrable function for $f \in L^p(\mu)$ and $g \in L^q(\mu)$. Therefore, φ_g is a well-defined map for $g \in L^q(\mu)$. Linearity of φ_g follows immediately from linearity of the integral. Moreover, the bound from Hölder's inequality shows that $\|\varphi_g\|_{L^p(\mu)^*} \leq \|g\|_q < \infty$, so φ_g is continuous by Exercise 8.2. In fact, $\|g\|_q = \|\varphi_g\|_{L^p(\mu)^*}$ by Exercise 8.5, so $g \mapsto \varphi_g$ is an isometric embedding of $L^q(\mu)$ into $L^p(\mu)^*$. It remains to check that this map is surjective. Let $\varphi \in L^p(\mu)^*$. We want to find a function $g \in L^q(\mu)$ such that $\varphi = \varphi_g$. We break the proof into intermediate cases.

CASE 1. μ is finite.

In this case, $\mathbb{1}_E \in L^p(\mu)$ for every $E \in \mathcal{B}$, so we may define a function $\nu : \mathcal{B} \rightarrow \mathbb{C}$ by $\nu(E) = \varphi(\mathbb{1}_E)$.

CLAIM 1. ν is a complex measure.

First, $\nu(\emptyset) = \varphi(\mathbb{1}_\emptyset) = \varphi(0) = 0$. Next, suppose $(E_n)_{n \in \mathbb{N}}$ is a sequence of disjoint measurable sets, and let $E = \bigsqcup_{n \in \mathbb{N}} E_n$. By linearity and continuity of φ , it suffices to show that $\sum_{n=1}^N \mathbb{1}_{E_n} \rightarrow \mathbb{1}_E$ in $L^p(\mu)$ as $N \rightarrow \infty$. But

$$\left\| \mathbb{1}_E - \sum_{n=1}^N \mathbb{1}_{E_n} \right\|_p = \left\| \sum_{n=N+1}^{\infty} \mathbb{1}_{E_n} \right\|_p = \mu \left(\bigsqcup_{n=N+1}^{\infty} E_n \right)^{1/p} \xrightarrow[N \rightarrow \infty]{} 0$$

by continuity of μ from above.

By construction, $\nu \ll \mu$, so by the Radon–Nikodym theorem, there is a μ -essentially unique measurable function $g : X \rightarrow \mathbb{C}$ such that $d\nu = g d\mu$.

CLAIM 2. $\varphi = \varphi_g$ on the space \mathcal{S} of simple functions.

Let $s = \sum_{j=1}^n c_j \mathbb{1}_{E_j} \in \mathcal{S}$. Then

$$\varphi(s) = \sum_{j=1}^n c_j \varphi(\mathbb{1}_{E_j}) = \sum_{j=1}^n c_j \nu(E_j) = \sum_{j=1}^n c_j \int_{E_j} g \, d\mu = \int_X sg \, d\mu = \varphi_g(s).$$

CLAIM 3. $\varphi = \varphi_g$ on $L^\infty(\mu)$.

(Note that $L^\infty(\mu) \subseteq L^p(\mu)$ since μ is a finite measure; see Exercise 8.6(a).) Given $f \in L^\infty(\mu)$, we may find a bounded sequence of simple functions $(s_n)_{n \in \mathbb{N}}$ such that $|s_n| \leq |f|$ and $s_n \rightarrow f$ a.e. by Proposition 3.7. Then by the bounded convergence theorem, $s_n \rightarrow f$ in $L^p(\mu)$, so $\varphi(f) = \lim_{n \rightarrow \infty} \varphi(s_n)$. Similarly, $s_n g \rightarrow fg$ a.e., and $|s_n g| \leq |fg| \in L^1(\mu)$, so by the dominated convergence theorem, $\varphi_g(f) = \lim_{n \rightarrow \infty} \varphi_g(s_n)$. By Claim 2, it follows that $\varphi(f) = \varphi_g(f)$.

CLAIM 4. $g \in L^q(\mu)$.

For $n \in \mathbb{N}$, let $E_n = \{|g| \leq n\}$, and let $g_n = g \mathbb{1}_{E_n}$ so that $|g_n| \leq |g|$ and $g_n \rightarrow g$ a.e. By the dominated convergence theorem, it follows that $g_n \rightarrow g$ in $L^1(\mu)$. If $g = 0$ a.e., then $g \in L^q(\mu)$, so assume $g \neq 0$. Then $g_n \neq 0$ for large n ; in particular, $\|g_n\|_q > 0$. Since μ is finite and g_n is bounded, we also have $\|g_n\|_q < \infty$.

Let $f_n = \frac{|g_n|^{q-2} g_n}{\|g_n\|_q^{q-1}}$ so that $f_n g = f_n g_n = \frac{|g_n|^q}{\|g_n\|_q^{q-1}}$. Then $\int_X f_n g \, d\mu = \|g_n\|_q$. Moreover,

$$\int_X |f_n|^p \, d\mu = \frac{1}{\|g_n\|_q^{p(q-1)}} \int_X |g_n|^{p(q-1)} \, d\mu = \frac{1}{\|g_n\|_q^q} \int_X |g_n|^q \, d\mu = 1.$$

Therefore, by Fatou's lemma and claim 3,

$$\|g\|_q \leq \liminf_{n \rightarrow \infty} \|g_n\|_q = \liminf_{n \rightarrow \infty} \int_X f_n g \, d\mu = \liminf_{n \rightarrow \infty} \varphi(f_n) \leq \|\varphi\|_{L^p(\mu)^*} < \infty.$$

By Claim 4, the functional φ_g is defined on all of $L^p(\mu)$.

CLAIM 5. $\varphi = \varphi_g$ on $L^p(\mu)$

Both φ and φ_g are continuous linear functionals on $L^p(\mu)$, and they agree on the dense set \mathcal{S} , so they must be equal.

Claims 4 and 5 together complete the proof in the finite case.

CASE 2. μ is σ -finite.

Write X as an increasing union $X = \bigcup_{n \in \mathbb{N}} X_n$ of measurable sets $X_n \in \mathcal{B}$ with $\mu(X_n) < \infty$. We may identify $L^p(\mu|_{X_n})$ with the subspace of $L^p(\mu)$ of functions that vanish outside of X_n . Using Case 1, let $g_n \in L^q(\mu|_{X_n})$ be the representative of $\varphi|_{L^p(\mu|_{X_n})}$. By uniqueness of g_n , we must have $g_n = g_m$ a.e. on X_m for $n > m$. Therefore, $g = \lim_{n \rightarrow \infty} g_n$ is defined a.e. on X and $\|g\|_q = \lim_{n \rightarrow \infty} \|g_n\|_q \leq \|\varphi\|_{L^p(\mu)^*} < \infty$. By the dominated convergence theorem and continuity of φ , we conclude that $\varphi = \varphi_g$.

CASE 3. μ is an arbitrary (positive) measure.

Let \mathcal{F} be the family of σ -finite subsets of X . For each $F \in \mathcal{F}$, we may use Case 2 to find $g_F \in L^q(\mu|_F)$ representing the functional $\varphi|_{L^p(\mu|_F)}$. Let $M = \sup_{F \in \mathcal{F}} \|g_F\|_q \leq \|\varphi\|_{L^p(\mu)^*} < \infty$. Let $(F_n)_{n \in \mathbb{N}}$ be a sequence in \mathcal{F} such that $\|g_{F_n}\|_q \rightarrow M$. Let $F = \bigcup_{n \in \mathbb{N}} F_n$, and let $g = g_F$. Then $\|g\|_q \geq \|g_{F_n}\|_q$ for every $n \in \mathbb{N}$, so $\|g\|_q = M$.

Suppose $A \in \mathcal{F}$ and $A \supseteq F$. Then

$$\int_X |g|^q d\mu = M^q \geq \int_X |g_A|^q d\mu = \int_X |g_{A \setminus F}|^q d\mu + \int_X |g|^q d\mu,$$

so $g_{A \setminus F} = 0$ a.e. and $g_A = g$ a.e.

We claim $\varphi = \varphi_g$. Let $f \in L^p(\mu)$. By Chebyshev's inequality (see Problem 6 on the exam study guide),

$$\mu \left(\left\{ |f| \geq \frac{1}{n} \right\} \right) \leq n^p \int_X |f|^p d\mu < \infty,$$

so $\{f \neq 0\} \in \mathcal{F}$. Let $A = F \cup \{f \neq 0\} \in \mathcal{F}$. Since f vanishes outside of A , we have $\varphi(f) = \varphi_{g_A}(f)$. Moreover, $g_A = g$ a.e., so $\varphi_{g_A}(f) = \varphi_g(f)$.

□

REMARK. Under the additional assumption that μ is σ -finite, the Riesz–Fréchet theorem holds also for $p = 1$ and $q = \infty$ by essentially the same argument presented above. (The only step requiring some modification is the proof of Claim 4.)

However, outside of extremely limited cases (such as counting measure on a finite set), the dual space $L^\infty(\mu)^*$ is much larger than $L^1(\mu)$. Providing a useful description of $L^\infty(\mu)^*$ requires the development of additional tools from functional analysis that will not be addressed in these notes.

COROLLARY 10.29

Let (X, \mathcal{B}, μ) be a measure space. If $\varphi : L^2(\mu) \rightarrow \mathbb{C}$ is a continuous linear functional, then there exists a unique $g \in L^2(\mu)$ such that $\varphi(f) = \langle f, g \rangle$ for every $f \in L^2(\mu)$.

REMARK. Corollary 10.29 can be proved using different techniques from functional analysis. It is a general fact of Hilbert spaces that every continuous linear functional on a Hilbert space can be represented as the inner product with an element of the Hilbert space. This is known as the Riesz representation theorem (not to be confused with the other Riesz representation theorem covered in Theorem 6.6 in these notes).

5. The Riesz Representation Theorem, Revisited

Let X be an LCH space. The Riesz representation theorem established a one-to-one correspondence between positive linear functionals on $C_c(X)$ and Radon measures on X . One may guess

that dropping the assumption of positivity on the linear functional corresponds to dropping the assumption of positivity on the measures. What we now want to show is that such a statement is true: the dual of $C_c(X)$ is a space of complex measures.

First we need some preliminaries. The space $C_c(X)$ is not complete in general, so when working with continuous linear functionals, it is more natural to work with its completion.

DEFINITION 10.30

Let X be an LCH space. We say that a function $f : X \rightarrow \mathbb{C}$ *vanishes at infinity* if for every $\varepsilon > 0$, there exists a compact set K such that $\sup_{x \notin K} |f(x)| < \varepsilon$. We denote by $C_0(X)$ the space of continuous functions on X that vanish at infinity.

The next proposition shows that the dual space $C_c(X)^*$ is the same as the dual space $C_0(X)^*$.

PROPOSITION 10.31

Let X be an LCH space. Then $C_0(X)$ is a Banach space with the norm $\|f\|_u = \sup_{x \in X} |f(x)|$, and $C_c(X)$ is dense in $C_0(X)$. In particular, if $\varphi : C_c(X)$ is a continuous linear functional, then φ has a unique extension to a continuous linear functional $\tilde{\varphi} : C_0(X) \rightarrow \mathbb{C}$.

PROOF. Let us first show that $C_0(X)$ is complete and then show that $C_c(X)$ is a dense subspace.

CLAIM 1. $C_0(X)$ is complete.

Let $(f_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in $C_0(X)$. Then for each $x \in X$, since $|f_n(x) - f_m(x)| \leq \|f_n - f_m\|_u$, the sequence $(f_n(x))_{n \in \mathbb{N}}$ is Cauchy. We may therefore define a limit pointwise by $f(x) = \lim_{n \rightarrow \infty} f_n(x)$.

Let $\varepsilon > 0$. Then choose $N \in \mathbb{N}$ so that $|f_n(x) - f_m(x)| < \varepsilon$ for every $n, m \geq N$ and every $x \in X$. Then for every $x \in X$ and every $n \geq N$, $|f_n(x) - f(x)| \leq \sup_{m \geq N} |f_n(x) - f_m(x)| \leq \varepsilon$. Hence, $f_n \rightarrow f$ uniformly. The uniform limit of continuous functions is continuous, so $f : X \rightarrow \mathbb{C}$ is continuous.

It remains to check that f vanishes at infinity. Let $\varepsilon > 0$. Let $n \in \mathbb{N}$ such that $\|f_n - f\|_u < \frac{\varepsilon}{2}$, and let $K \subseteq X$ be a compact set such that $\sup_{x \notin K} |f_n(x)| < \frac{\varepsilon}{2}$. Then $\sup_{x \notin K} |f(x)| < \varepsilon$, so $f \in C_0(X)$.

CLAIM 2. $C_c(X)$ is dense in $C_0(X)$.

Let $f \in C_0(X)$, and let $\varepsilon > 0$. Let K be a compact set such that $\sup_{x \notin K} |f(x)| < \varepsilon$. By Urysohn's lemma, let $g \in C_c(X)$ with $K \prec g$. Then $\text{supp}(fg) \subseteq \text{supp}(g)$, so $fg \in C_c(X)$. Moreover, $|fg| \leq |f|$, and $fg = f$ on K . Thus, $\|fg - f\|_u = \sup_{x \in X} (|f(x)g(x)| - |f(x)|) \leq \sup_{x \notin K} |f(x)| < \varepsilon$.

□

Now we need to find the appropriate space of measures to replace positive Radon measures. Note that if a positive linear functional is continuous (bounded) on $C_c(X)$, then it is represented by a finite measure, which guarantees regularity (Radon measures are inner regular on all (σ) -finite sets by Proposition 9.4). So the natural guess for the dual space of $C_0(X)$ is the space of regular complex measures, but what does "regular" mean in this context? To answer this question, we associate to each complex measure a positive measure capturing some of its behavior.

DEFINITION 10.32

Let (X, \mathcal{B}) be a measurable space, and let $\mu : \mathcal{B} \rightarrow \mathbb{C}$ be a complex measure. The *total variation* of μ is the measure $|\mu|$ defined by

$$|\mu|(E) = \sup \left\{ \sum_{n=1}^{\infty} |\mu(E_n)| : E = \bigsqcup_{n \in \mathbb{N}} E_n \right\}$$

for $E \in \mathcal{B}$. The *total variation norm* of μ is the quantity $\|\mu\|_{TV} = |\mu|(X)$.

The definition of the total variation of a complex measure is motivated by Proposition 10.10. For more on the total variation measure, see Exercise 10.3.

PROPOSITION 10.33

Let (X, \mathcal{B}) be a measurable space, and let $\mathcal{M}(X, \mathcal{B})$ be the space of complex measures on (X, \mathcal{B}) . Then $\|\cdot\|_{TV}$ is a norm on $\mathcal{M}(X, \mathcal{B})$.

PROOF. We need to check three properties.

CLAIM 1. $\|\cdot\|_{TV}$ is absolutely homogeneous.

For $\mu \in \mathcal{M}(X, \mathcal{B})$ and $c \in \mathbb{C}$,

$$\|c\mu\|_{TV} = \sup \left\{ \sum_{n=1}^{\infty} \underbrace{|c\mu(E_n)|}_{|c||\mu(E_n)|} : X = \bigsqcup_{n \in \mathbb{N}} E_n \right\} = |c| \|\mu\|_{TV}.$$

CLAIM 2. $\|\cdot\|_{TV}$ satisfies the triangle inequality.

Let $\mu, \nu \in \mathcal{M}(X, \mathcal{B})$. Then

$$\|\mu + \nu\|_{TV} = \sup \left\{ \sum_{n=1}^{\infty} \underbrace{|\mu(E_n) + \nu(E_n)|}_{\leq |\mu(E_n)| + |\nu(E_n)|} : X = \bigsqcup_{n \in \mathbb{N}} E_n \right\} \leq \|\mu\|_{TV} + \|\nu\|_{TV}.$$

CLAIM 3. $\|\cdot\|_{TV}$ is positive definite.

Suppose $\mu \in \mathcal{M}(X, \mathcal{B})$ and $\|\mu\|_{TV} = 0$. Then $\sum_{n=1}^{\infty} |\mu(E_n)| = 0$ for every $X = \bigsqcup_{n \in \mathbb{N}} E_n$. In particular, given $E \in \mathcal{B}$, we put $E_1 = E$, $E_2 = X \setminus E$, and $E_3 = E_4 = \dots = \emptyset$ to conclude that $|\mu(E)| + |\mu(X \setminus E)| = 0$. Hence, $\mu = 0$. □

DEFINITION 10.34

Let X be an LCH space. A complex measure $\mu : \text{Borel}(X) \rightarrow \mathbb{C}$ is *regular* if its total variation $|\mu|$ is regular.

We now have all of the relevant definitions in order to state a version of the Riesz representation theorem relating continuous linear functionals to complex measures.

THEOREM 10.35: RIESZ REPRESENTATION THEOREM (COMPLEX MEASURES)

Let X be an LCH space. Then $C_0(X)^*$ is isometrically isomorphic to the space of complex regular Borel measure on X with the total variation norm. That is, for any continuous linear functional $\varphi : C_0(X) \rightarrow \mathbb{C}$, there is a unique complex regular Borel measure μ on X such that

$$\varphi(f) = \int_X f \, d\mu$$

for every $f \in C_0(X)$, and $\|\varphi\|_{C_0(X)^*} = \|\mu\|_{TV}$.

PROOF. Let $C_0(X, \mathbb{R}) \subseteq C_0(X)$ be the space of real-valued continuous functions on X that vanish at infinity. Note that every element $f \in C_0(X)$ can be written as $f_1 + if_2$ with $f_1, f_2 \in C_0(X, \mathbb{R})$, and $\varphi(f) = \varphi(f_1) + i\varphi(f_2)$, so φ is determined by its values on $C_0(X, \mathbb{R})$. Let $\varphi_1, \varphi_2 \in C_0(X)^*$ such that $\varphi_j(f) \in \mathbb{R}$ for every $f \in C_0(X, \mathbb{R})$ and $\varphi = \varphi_1 + i\varphi_2$. (We define φ_1 by $\text{Re}(\varphi)$ on $C_0(X, \mathbb{R})$ and extend to $C_0(X)$ by linearity and similarly for φ_2 .) We want to prove a Jordan decomposition for φ_1 and φ_2 to be able to apply the Riesz representation theorem for positive linear functionals.

CLAIM 1. If $\psi : C_0(X, \mathbb{R}) \rightarrow \mathbb{R}$ is a bounded linear functional, then there exist positive bounded linear functionals ψ^+ and ψ^- such that $\psi = \psi^+ - \psi^-$.

For $f \in C_0(X, \mathbb{R})$, $f \geq 0$, define

$$\psi^+(f) = \sup \{\psi(g) : g \in C_0(X, \mathbb{R}), 0 \leq g \leq f\}.$$

Note that $0 = \psi(0) \leq \psi^+(f) \leq \sup_{0 \leq g \leq f} |\psi(g)| \leq \|\psi\|_{C_0(X, \mathbb{R})^*} \|f\|_u$. We need to show that ψ^+ is linear, which will be similar to our proof of linearity of the integral.

For $c \geq 0$, we have $\psi^+(cf) = c\psi^+(f)$, since ψ is linear. Let $f_1, f_2 \in C_0(X, \mathbb{R})$, $f_1, f_2 \geq 0$. If $0 \leq g_1 \leq f_1$ and $0 \leq g_2 \leq f_2$, then $0 \leq g_1 + g_2 \leq f_1 + f_2$, so $\psi^+(f_1 + f_2) \geq \psi^+(f_1) + \psi^+(f_2)$. On the other hand, given $0 \leq g \leq f_1 + f_2$, we may set $g_1 = \min\{g, f_1\}$ and $g_2 = g - g_1$ so that $0 \leq g_j \leq f_j$ and $\psi(g) = \psi(g_1) + \psi(g_2) \leq \psi^+(f_1) + \psi^+(f_2)$. Hence, $\psi^+(f_1 + f_2) \leq \psi^+(f_1) + \psi^+(f_2)$. This shows that ψ^+ extends to a linear functional on $C_0(X, \mathbb{R})$ by $\psi^+(f) = \psi^+(f^+) - \psi^+(f^-)$. Moreover, for $f \in C_0(X, \mathbb{R})$,

$$|\psi^+(f)| = |\psi^+(f^+) - \psi^+(f^-)| \leq \psi^+(f^+) + \psi^+(f^-) = \psi^+(|f|) \leq \|\psi\|_{C_0(X, \mathbb{R})^*} \|f\|_u,$$

so $\psi^+ \in C_0(X, \mathbb{R})^*$.

Let $\psi^- = \psi^+ - \psi \in C_0(X, \mathbb{R})^*$. Suppose $f \geq 0$. Then $\psi^+(f) \geq \psi(f)$, so $\psi^-(f) \geq 0$. That is, ψ^- is a positive bounded linear functional.

By Claim 1, we may decompose $\varphi_j = \varphi_j^+ - \varphi_j^-$ for $j \in \{1, 2\}$. By the Riesz representation theorem, there is a finite regular Borel measure μ_j^\pm such that $\varphi_j^\pm(f) = \int_X f \, d\mu_j^\pm$ for every $f \in C_c(X)$ and hence for every $f \in C_0(X)$ by continuity. Define $\mu = \mu_1^+ - \mu_1^- + i(\mu_2^+ - \mu_2^-)$. Then μ is a complex regular Borel measure on X and $\varphi(f) = \int_X f \, d\mu$.

CLAIM 2. $\|\mu\|_{TV} = \|\varphi\|_{C_0(X)^*}$.

Let $f \in C_0(X)$ with $\|f\|_u \leq 1$. Then

$$|\varphi(f)| = \left| \int_X f \, d\mu \right| = \left| \int_X f \frac{d\mu}{d|\mu|} \, d|\mu| \right| \leq \int_X |f| \, d|\mu| \leq |\mu|(X) = \|\mu\|_{TV},$$

where we have used that $\left| \frac{d\mu}{d|\mu|} \right| = 1$ $|\mu|$ -a.e. (see Exercise 10.3). Therefore, $\|\varphi\|_{C_0(X)^*} \leq \|\mu\|_{TV}$.

Let $\varepsilon > 0$. Let $g = \frac{d\mu}{d|\mu|}$. Note that $|g| = 1$ $|\mu|$ -a.e. By Lusin's theorem (Theorem 9.9), there is a continuous function $f \in C_c(X)$ such that $|\mu|(\{f \neq \bar{g}\}) < \frac{\varepsilon}{2}$ and $\|f\|_u \leq 1$. Thus,

$$\|\varphi\|_{C_0(X)^*} \geq |\varphi(f)| = \left| \int_X f \, d\mu \right| = \left| \int_X fg \, d|\mu| \right| \geq |\mu|(\{f = \bar{g}\}) - |\mu|(\{f \neq \bar{g}\}) > \|\mu\|_{TV} - \varepsilon.$$

It remains to check the uniqueness part of the theorem.

CLAIM 3. If μ and ν are complex regular Borel measures on X and $\int_X f \, d\mu = \int_X f \, d\nu$ for every $f \in C_0(X)$, then $\mu = \nu$.

Let $\rho = \mu - \nu$. Then $\int_X f \, d\rho = 0$ for every $f \in C_0(X)$, so by Claim 2, $\|\rho\|_{TV} = 0$. Thus, $\rho = 0$ by Proposition 10.33. □

Chapter Notes

The presentation of material in this chapter is heavily influenced by the book of Folland [2]. The first 3 sections correspond to material from [2, Chapter 3], section 4 to [2, Section 6.2], and the final section to [2, Section 7.3]. The largest point of departure from [2] is the proofs of the Hahn decomposition theorem and the Radon–Nikodym theorem. The construction of the Hahn decomposition provided in these notes uses the original argument of Hahn, which, for reasons still somewhat mysterious to me, has largely been displaced by other proofs in the most commonly used measure theory texts. It is also quite common for the Lebesgue decomposition and Radon–Nikodym theorem to be proved simultaneously (as in different arguments presented in [2] and [7, 9]); this can provide a slick and clever proof, but the end result is to reveal the existence of a Radon–Nikodym derivative without much indication of what that derivative looks like. This is in contrast with the proof presented in these notes, which builds the Radon–Nikodym derivative explicitly through level sets using the Hahn decomposition theorem. I learned of Hahn's proof of the Hahn decomposition theorem, the level set construction of the Radon–Nikodym derivative, and the role of s -finite measures in the whole theory from one of my graduate school professors, Neil Falkner. Additional historical comments and sketches of many of the proofs from this chapter appear in one of his articles [1].

Exercises

10.1 Prove Proposition 10.10.

10.2 Let (X, \mathcal{B}) be a measurable space.

(a) Let $\mu, \nu, \rho : \mathcal{B} \rightarrow [0, \infty]$ be σ -finite measures, and suppose $\rho \ll \nu \ll \mu$. Prove the “chain rule”

$$\frac{d\rho}{d\mu} = \frac{d\rho}{d\nu} \frac{d\nu}{d\mu}.$$

(b) Suppose μ, ν are finite (positive) measures on (X, \mathcal{B}) , and $\mu \approx \nu$. Show that the Radon–Nikodym derivative $f = \frac{d\nu}{d\mu}$ satisfies $0 < f < \infty$ μ -a.e., and $\frac{d\mu}{d\nu} = \frac{1}{f}$ ν -a.e.

10.3 Let (X, \mathcal{B}) be a measurable space.

(a) Let $\mu : \mathcal{B} \rightarrow [-\infty, \infty]$ be a signed measure, and let $|\mu|$ be the total variation measure. Show that $\mu \ll |\mu|$ and describe the Radon–Nikodym derivative $\frac{d\mu}{d|\mu|}$.

(b) Suppose $\mu : \mathcal{B} \rightarrow \mathbb{C}$ is a complex measure. Define the *total variation measure* $|\mu|$ by

$$|\mu|(E) = \sup \left\{ \sum_{n=1}^{\infty} |\mu(E_n)| : E = \bigsqcup_{n \in \mathbb{N}} E_n \right\}$$

for $E \in \mathcal{B}$. Suppose ν is a σ -finite positive measure on (X, \mathcal{B}) such that $\mu \ll \nu$. (For example, applying the Jordan decomposition theorem to the real and imaginary parts of μ and writing $\mu = \nu_1 - \nu_2 + i(\nu_3 - \nu_4)$ as a combination of positive finite measures, one can take $\nu = \sum_{j=1}^4 \nu_j$.)

Prove that $|\mu|$ is a measure and $\frac{d|\mu|}{d\nu} = \left| \frac{d\mu}{d\nu} \right|$ a.e. Conclude that there exists a measurable function $\theta : X \rightarrow [0, 1)$ such that $\frac{d\mu}{d|\mu|}(x) = e^{2\pi i \theta(x)}$ for $|\mu|$ -a.e. $x \in X$.

10.4 Let (X, \mathcal{B}) be a measurable space, and suppose ν, μ are positive measures on (X, \mathcal{B}) such that $d\nu = g d\mu$. Let $p, q \in (1, \infty)$ be conjugate exponents. Prove that if $g \in L^p(\mu)$, then

$$\nu(A) \leq \|g\|_p \mu(A)^{1/q}.$$

Bibliography

- [1] Neil Falkner. Hahn's proof of the Hahn decomposition theorem, and related matters, *Amer. Math. Monthly* **126** (2019), 264–268.
- [2] Gerald B. Folland. *Real Analysis*, 2nd edition (John Wiley & Sons, Inc., 1999).
- [3] R. K. Getoor. *Excessive Measures*, Probab. Appl. (Birkhäuser Boston, 1990).
- [4] Alexander S. Kechris. *Classical Descriptive Set Theory*, Grad. Texts in Math., **156** (Springer, 1995).
- [5] J. E. Littlewood. *Lectures On The Theory Of Functions* (Oxford University Press, 1944).
- [6] John C. Oxtoby. *Measure and Category*, 2nd edition, Grad. Texts in Math., **2** (Springer, 1980).
- [7] Walter Rudin. *Real and Complex Analysis*, 3rd edition (McGraw-Hill Book Co., 1987).
- [8] Robert M. Solovay. A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. Math. (2)* **92** (1970) 1–56.
- [9] Elias M. Stein and Rami Shakarchi. *Real Analysis* (Princeton University Press, 2005).
- [10] Terence Tao. *An Introduction to Measure Theory*, Grad. Stud. Math., **126** (American Mathematical Society, 2011). Preliminary version available online at <https://terrytao.wordpress.com/wp-content/uploads/2012/12/gsm-126-tao5-measure-book.pdf>
- [11] Anush Tserunyan. *Introduction to Descriptive Set Theory*. Online lecture notes, available at: https://www.math.mcgill.ca/atserunyan/Teaching_notes/dst_lectures.pdf