

October 15, 2024

## Problem Set 5

**Exercise 1.** Consider the representation of the group  $U(1) = \{e^{i\theta}, \theta \in [0, 2\pi[ \} \subset \mathbb{C}$  in  $V = \mathbb{C}^2$  given by the rotation matrix

$$\rho(e^{i\theta}) = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

Decompose  $V$  into a direct sum of two irreducible unitary complex representations of  $U(1)$ .

**Exercise 2.** Let  $A$  be an associative algebra over a field  $k$ . For a representation  $V$  of  $A$ , consider the vector space  $\text{End}_A(V)$  of endomorphisms of the representation  $V$  (linear maps  $V \rightarrow V$  commuting with the action of  $A$  in  $V$ ). Let  $V$  be the left regular representation,  $V = A$ . Show that  $\text{End}_A(A)$  is an associative algebra isomorphic to  $A^{\text{op}}$ , the algebra  $A$  with the opposite multiplication.

**Exercise 3.** Let  $A = \text{Mat}_d(k)$  for a field  $k$ . Prove that the algebra  $A$  is semisimple, meaning that any finite dimensional representation of  $A$  over  $k$  is isomorphic to a direct sum of irreducible representations.

*Hint:* Consider the basis of matrices with a single nonzero matrix element  $\{E_{ij}\}$  in  $A$ . Show that for a representation  $V$  of  $A$ , we have  $V = \bigoplus_{i=1}^d E_{ii}V$  and that for  $v \in E_{11}V$ , the linear span of  $\{E_{11}v, E_{21}v, \dots, E_{d1}v\}$  is a subrepresentation of  $V$  isomorphic to  $k^d$ . Conclude by choosing a basis in  $E_{11}V$ .

**Exercise 4.** Let  $A$  be a finite dimensional algebra, and  $\text{Rad}(A)$  the set of all elements of  $A$  that act by 0 in all irreducible representations of  $A$ .

(a) Show that  $\text{Rad}(A)$  is a two-sided ideal in  $A$ .

(b) Let  $I \subset A$  be a two-sided nilpotent ideal, meaning that there exist  $n \in \mathbb{N}$  such that  $x^n = 0$  for all  $x \in I$ . Show that  $I \subset \text{Rad}(A)$ .

**Exercise 5.** Recall that the character of a finite dimensional representation  $V$  of an algebra  $A$  over a field  $k$  is defined as  $\chi_V(a) = \text{Tr}_V \rho(a)$ . Show that if  $V$  is a finite dimensional representation of  $A$ , and  $W \subset V$  a subrepresentation, then the character  $\chi_V = \chi_W + \chi_{V/W}$ .

**Exercise 6.** (a) Construct all possible representations of the cyclic group  $C_2 = \langle t \mid t^2 = 1 \rangle$  in  $V$ , where  $V$  is a two-dimensional vector space over the field  $\mathbb{F}_2$ . Decompose the obtained representations into a direct sum of irreducibles.

(b) For the obtained irreducible representations, consider the intertwiners  $\phi : V \rightarrow V$  that commute with the action of the group  $C_3$ . Show how the Schur's lemma fails in the case of the field  $\mathbb{F}_2$ , which is not algebraically closed.