

Differential equations of movement

- Consider an infinitely small cube subjected to a force represented by the vector:

$$\mathbf{f}(\mathbf{x}, t) = \begin{pmatrix} f_1(\mathbf{x}, t) \\ f_2(\mathbf{x}, t) \\ f_3(\mathbf{x}, t) \end{pmatrix}.$$


- Stress is not uniform. The variation of the stress between two opposite sides is linear and

$$d\sigma_{ij} = \partial_{x_k} \sigma_{ij} dx_k \quad (*)$$

- Newton 2nd law of motion taking into account internal forces $\mathbf{f}^i$ is

$$\mathbf{f}^i(\mathbf{x}, t) + \mathbf{f}(\mathbf{x}, t) = \rho(\mathbf{x}) \ddot{\mathbf{u}}(\mathbf{x}, t) dx_1 dx_2 dx_3$$

where $\rho(\mathbf{x})$ is the material density, $\ddot{\mathbf{u}} = \partial_{tt} \mathbf{u}$ is the acceleration, $dx_1 dx_2 dx_3$ is the volume of the cube.



Credit: [N]

Equilibrium of forces in the x_1 direction:

$$(\cancel{\sigma_{11}} + d\sigma_{11} - \cancel{\sigma_{11}}) dx_2 dx_3 + (\cancel{\sigma_{21}} + d\sigma_{21} - \cancel{\sigma_{21}}) dx_1 dx_3 + (\cancel{\sigma_{31}} + d\sigma_{31} - \cancel{\sigma_{31}}) dx_1 dx_2 + f_1 dx_1 dx_2 dx_3 = \rho \ddot{u}_1 dx_1 dx_2 dx_3$$

After accounting $(*)$, the latter equation is:

$$\frac{\partial \sigma_{11}}{\partial x_1} + \frac{\partial \sigma_{21}}{\partial x_2} + \frac{\partial \sigma_{31}}{\partial x_3} + f_1 = \rho \ddot{u}_1$$

Similarly, the equilibrium of forces in the x_2 and x_3 directions leads to:

$$\frac{\partial \sigma_{1j}}{\partial x_1} + \frac{\partial \sigma_{2j}}{\partial x_2} + \frac{\partial \sigma_{3j}}{\partial x_3} + f_j = \rho \ddot{u}_j \quad (j=1,2,3)$$

The equilibrium equations can be written in matrix form as

$$\nabla^T \sigma + f = \rho \ddot{u}$$

where

$$\nabla^T = \begin{pmatrix} \partial_x & 0 & 0 & 0 & \partial_z & \partial_y \\ 0 & \partial_y & 0 & \partial_z & 0 & \partial_x \\ 0 & 0 & \partial_z & \partial_y & \partial_x & 0 \end{pmatrix}, \quad \sigma = \begin{pmatrix} \sigma_{11} \\ \sigma_{22} \\ \sigma_{33} \\ \sigma_{23} \\ \sigma_{31} \\ \sigma_{12} \end{pmatrix}.$$

Since $\sigma = C \varepsilon = C \nabla u$ we obtain

$$\boxed{\nabla^T C \nabla u + f = \rho \ddot{u}}$$

- ① Introduce an admissible (avoid divergence of integral) **virtual displacement**:

$$\delta \mathbf{u}(\mathbf{x}) = \begin{pmatrix} \delta u_1(\mathbf{x}) \\ \delta u_2(\mathbf{x}) \\ \delta u_3(\mathbf{x}) \end{pmatrix}$$

- ② Multiply the differential equation by $\delta \mathbf{u}^T$ and integrate it over the spatial domain Ω .

$$\int_{\Omega} \delta \mathbf{u}^T (\nabla^T C \nabla \mathbf{u} + \mathbf{f}) d\Omega = \int_{\Omega} \rho \delta \mathbf{u}^T \ddot{\mathbf{u}} d\Omega$$

- ③ Apply the divergence theorem to the first term:


$$-\int_{\Omega} (\nabla \delta \mathbf{u})^T C \nabla \mathbf{u} d\Omega + \int_{\Gamma} \delta \mathbf{u}^T \mathbf{N}^T C \nabla \mathbf{u} d\Gamma + \int_{\Omega} \delta \mathbf{u}^T \mathbf{f} d\Omega = \int_{\Omega} \rho \delta \mathbf{u}^T \ddot{\mathbf{u}} d\Omega$$

Let's clarify how the term $\int_{\Omega} (\delta \mathbf{u})^T \nabla^T C \nabla \mathbf{u} d\Omega$ is transformed into

$$-\int_{\Omega} (\nabla \delta \mathbf{u})^T C \nabla \mathbf{u} d\Omega + \int_{\Gamma} (\delta \mathbf{u})^T \mathbf{N}^T C \nabla \mathbf{u} d\Gamma$$

by applying the divergence theorem:

Div. theorem: $\int_{\Omega} \partial_{x_i} u d\Omega = \int_{\Gamma} u n_i d\Gamma$



$\vec{n}$ outward pointing normal vector.

$$\int_{\Omega} (\nabla u)^T \nabla^T C \nabla u \, d\Omega = \int_{\Omega} (\nabla u)^T \nabla^T \sigma \, d\Omega$$

$$= \sum_{j=1}^3 \int_{\Omega} u_j \left(\frac{\partial \sigma_{1j}}{\partial x_1} + \frac{\partial \sigma_{2j}}{\partial x_2} + \frac{\partial \sigma_{3j}}{\partial x_3} \right) d\Omega$$

$$= \sum_{i=1}^3 \sum_{j=1}^3 \int_{\Omega} u_j \frac{\partial \sigma_{ij}}{\partial x_i} d\Omega$$

$$= \sum_{i=1}^3 \sum_{j=1}^3 \int_{\Omega} \underbrace{\frac{\partial}{\partial x_i} (u_j \sigma_{ij}) - \frac{\partial u_j}{\partial x_i} \sigma_{ij}}_{\text{apply div. thm.}} d\Omega$$

$$= \sum_{i=1}^3 \sum_{j=1}^3 \int_{\Gamma} u_j \sigma_{ij} n_i \, d\Gamma - \int_{\Omega} \frac{\partial u_j}{\partial x_i} \sigma_{ij} \, d\Omega$$

$$= \int_{\Gamma} (\nabla u)^T N^T \sigma \, d\Gamma - \int_{\Omega} (\nabla \nabla u)^T \sigma \, d\Omega$$

$$= \int_{\Gamma} (\nabla u)^T N^T C \nabla u \, d\Gamma - \int_{\Omega} (\nabla \nabla u)^T C \nabla u \, d\Omega$$