


The four laws of black hole mechanics

- we saw black holes generically form as a result of gravitational collapse (assuming cosmic censorship)
- due to the uniqueness theorems, these final states are characterised by a small # of parameters (M, J), just as a thermodynamic system is
- we will now further explore the analogy between b.h. & thermodynamics
- a useful (though not very detailed) review w/ refs to the original literature is: R. Wald: "The thermodyn. of b.h." in *Living reviews in relativity*

Setup: consider a **stationary** black hole space-time. It has been shown that in vacuum or electrovac GR, the horizon must be a **Killing horizon**, i.e. a null hypersurface to which a Killing vector field is normal (and thus $\xi^2 = 0$ on $\mathcal{H}$)

- as long as the b.h.s are non-extremal & we look @ the maximally analytically extended spt, these horizons are **bifurcate Killing horizons**



- pair of orthogonal null surfaces =
= Killing horizons w.r.t the **same** K.V.


• the Killing vect. ξ^μ vanishes on B

Example : Rindler space

- patch of (2d) Minkowski in coord. adapted to the experience of an accelerated observer

$$ds^2 = -r^2 dt^2 + dr^2 + d\vec{x}_\perp^2, \text{ obtained via } \begin{cases} x^0 = r \sinh t \\ x^1 = r \cosh t > 0 \end{cases}$$

↑ patch $\mathbb{R}^{1,1} \times \mathbb{R}^{d-2}$



- $\xi \equiv \frac{\partial}{\partial t}$ is a K.V. $\rightarrow$ boost in Minkowski

$$\partial_t = \frac{\partial x^0}{\partial t} \partial_{x^0} + \frac{\partial x^1}{\partial t} \partial_{x^1} = (x^1 \partial_0 + x^0 \partial_1)$$

- the lines $x^1 = \pm x^0$ form a bifurcate Killing horizon, which has 2 components $\cap \underbrace{\mathcal{B}}_{x^0=x^1=0}$ w/ $\xi|_{\mathcal{B}} = 0$

- check I: $\xi^2 = X_0^2 - X_1^2 = 0$ for $x^1 = \pm x^0$

- in Rindler coord $x^0 = \pm x^1$ is obtained via $r \rightarrow 0, t \rightarrow \pm\infty$ w/ $r e^{\mp t}$ fixed

- more precisely, by intro. $u, v = x^0 \pm x^1 = \pm r e^{\pm t}$, the horizon's components are at $v=0, u$ fixed & $u=0, v$ fixed < 0

- in terms of these $\xi = u \partial_u - v \partial_v$

check II

- $\xi \perp \text{horizon} \Rightarrow \xi_\mu \propto n_\mu \propto \partial_r = \frac{\partial u}{\partial r} \partial_u + \frac{\partial v}{\partial r} \partial_v = e^t \partial_u - e^{-t} \partial_v \begin{cases} \propto \partial_u, & t > 0 \\ \propto -\partial_v, & t < 0 \end{cases}$

Other examples: $\kappa = 2M$ in Schwarzschild, for ∂_t

$\kappa = \kappa_+$ in Kerr, for $\partial_t + \Omega_H \partial_\phi$ w/ $\Omega_H =$ angular velocity of the horizon

Some properties of Killing horizons :

(see e.g. Townsend 9707012)

- let l^μ be the normal to the null hypersurface $\mathcal{N}$
- l^μ null $\Rightarrow l^\mu$ is also a tg. vector in $\mathcal{N}$, i.e. $l^\mu = \frac{dx^\mu}{d\lambda}$ for some curve $x^\mu(\lambda)$ in $\mathcal{N}$
- the curves $x^\mu(\lambda)$ are geodesics: $l^\lambda \nabla_\lambda l^\mu \propto l^\mu$ (see e.g. Townsend)
generators of $\mathcal{N}$

$l_\mu \perp \Rightarrow l_\mu = f \partial_\mu S$ $\leftarrow$ function that specifies hypersurf.

$$\begin{aligned} \Rightarrow l^\lambda \nabla_\lambda l_\mu &= l^\lambda \nabla_\lambda (f \partial_\mu S) = l^\lambda \partial_\lambda f \frac{l_\mu}{f} + l^\lambda f \nabla_\lambda \partial_\mu S = l^\lambda \partial_\lambda \ln f l_\mu + \\ &+ l^\lambda f \nabla_\mu \partial_\lambda S = \# l_\mu = l^\lambda f \nabla_\mu \ln f \underbrace{\quad}_{\text{bound}} = \underbrace{l^2}_{\text{bound}} \frac{\partial_\mu f}{f} + \frac{1}{2} \nabla_\mu l^2 \\ &\quad \perp \nabla t^\mu = \text{tg. vect to } \mathcal{N} \Rightarrow \propto l^\mu \text{ q.e.d.} \end{aligned}$$

• affine param: $l^\lambda \nabla_\lambda l^\mu = 0$

• since $\xi^\mu \propto l^\mu$, the Killing vect $\perp \mathcal{N}$ satisfies

$$\boxed{\xi^\mu \nabla_\mu \xi^\nu = \kappa \xi^\nu}$$

$\uparrow$ geodesic eqn. in non-affine param
 ξ has a natural normaliz. @ ∞

surface gravity

• this leads to the 1st interpretation of surface gravity:

since $\xi^\mu = f l^\mu$ (affine param), we have $\xi^\mu \nabla_\mu (f l^\nu) = \xi^\mu \partial_\mu \ln f \xi^\nu$

so κ measures the failure of the Killing param, $v: \xi^\mu \partial_\mu v = 1$, to equal 1
 can show $f = 1/\kappa$ Affine Param.

• for the **2nd interpretation** note that proper acceleration particle along ξ^μ

$$\text{is } a^\mu = u^\lambda \nabla_\lambda u^\mu, \text{ w/ } u^\mu = \frac{\xi^\mu}{\sqrt{-\xi^2}}$$

$$\Rightarrow a^\mu = \frac{\xi^\lambda}{\sqrt{-\xi^2}} \nabla_\lambda \frac{\xi^\mu}{\sqrt{-\xi^2}} = \frac{\xi^\lambda \nabla_\lambda \xi^\mu}{-\xi^2} + \underbrace{\xi^\lambda \xi^\mu \xi^\nu \nabla_\lambda \xi_\nu}_{=0}$$

Noting that $K^2 = \lim_{\rightarrow \mathcal{N}} \frac{(\xi^\mu \nabla_\mu \xi_\lambda)(\xi^\nu \nabla_\nu \xi^\lambda)}{\xi^2}$ acceleration orbits $\xi \times \xi^2$ redshift

so $K = \lim_{\rightarrow \text{hor}} (a V)$ force that must be exerted @ infinity to hold a unit test mass in place, hence the name

$\uparrow$ redshift factor $\sqrt{-\xi^2}$

• also possible to show that $K^2 = -\frac{1}{2} \nabla^\mu \xi^\nu \nabla_\mu \xi_\nu / \mathcal{N}$ (see e.g. Townsend) Wald p. 332

• **3rd interpretation**: K also given by $2\pi/\text{periodicity}$ of the euclidean time coord. chosen to ensure that Euclidean b.h. geom. is smooth (no conical defect)

Example: Rindler space

can also be obt. from $\perp$ + affine parameter.

• $\xi^\mu = \partial_t$, while null geod || future horizon is ∂_u , $\xi \approx u \partial_u$ near future hor

• $\Rightarrow f = u \quad \kappa = \xi^\lambda \partial_\lambda \ln f = u \partial_u \ln u = 1$

• also from periodicity: $dr^2 + r^2 dt_E^2 \Rightarrow t_E \sim t_E + 2\pi \Rightarrow \kappa = 1$

• otherwise $\xi^\lambda \nabla_\lambda \xi_\mu = -\xi^\lambda \nabla_\mu \xi_\lambda = -\frac{1}{2} \partial_\mu \xi^2 = -\frac{1}{2} \left(V \delta_\mu^u + u \delta_\mu^v \right) = \pm \xi_\mu \Big|_{\substack{\nearrow \\ \text{near hor.}}} \Rightarrow \kappa = \pm 1$

$\underbrace{-r^2 = UV}_{\nearrow}$

Exercise: ① Compute the surface gravity associated w/ the Killing horizon of the Schwarzschild black hole

② Compare the result to $\frac{1}{2\pi} \times$ the periodicity of the Euclidean Schwarzschild time necessary to have a smooth geom.

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \frac{dr^2}{1 - \frac{2M}{r}} + r^2 d\Omega_2^2 \quad t = -t_E$$

near $r = 2M + \delta^2$ $dt_E^2 \left(1 - \frac{2M}{2M + \delta^2}\right) + \frac{(2\delta d\delta)^2}{2M + \delta^2 - 2M} = \frac{\delta^2 dt_E^2}{2M} + 8M d\delta^2 \approx 8M \left(d\delta^2 + \frac{\delta^2 dt_E^2}{16M^2} \right)$

$\Rightarrow$ no conical singularity if $t_E \sim t_E + 2\pi \cdot 4M \Rightarrow \boxed{\kappa = \frac{1}{4M}} \times G$

Zeroth law of b.h. mechanics

The surface gravity, κ , of a stationary b.h. is constant over the event horizon.

Pf: 2 steps : 1st, prove that $k = \text{const.}$ on orbits of ξ , ($\xi^\mu \partial_\mu k^2 = 0$)

(Hint: use K.V. id $\nabla_\sigma \nabla_\mu \xi^\nu = R^\nu_{\mu\sigma\rho} \xi^\rho$)

2nd, prove that $\kappa = \text{const}$ on $\mathcal{B}$ ($t^\lambda \nabla_\lambda \kappa^2 = -\nabla^\mu \xi^\nu t^\lambda R_{\nu\mu\lambda\sigma} \xi^\sigma|_{\mathcal{W}}$)
 $t^\lambda t^\lambda t_{\mu\lambda} \mathcal{W} \rightarrow \mathcal{B}$

since $k = \text{const}$ on orbits of $\mathbb{G}$, $k = \text{const}$ on $\mathcal{N}$ if it is const on $\mathcal{B}$

The reason this is called the zeroth law is that, as we will show, κ is analogous to a temperature, so this is reminiscent of the zeroth law of thermodynamics: temperature is uniform everywhere in a system in thermal equilibrium.

- note our proof is purely geometrical, but makes use of the bifurcate structure of the Killing horizon. I a different proof which does not make use of it, but uses instead Einstein's eqns + dominant energy cond. (see e.g. Wald's book)

The first law

$$\frac{\kappa}{2\pi} \frac{\delta A}{4G} = \delta M - \Omega_H \delta J \quad (+..)$$

- several different ways to show it

①. Check nearby stationary solutions satisfy it

Example : Schwarzschild

$$R_s = 2GM \Rightarrow A = 16\pi G^2 M^2, K = \frac{1}{4GM} \frac{1}{8\pi GM} \frac{\delta(16\pi G^2 M^2)}{4G} = 8M$$

- note that since b.h. absorb energy but do not emit (classically) $\delta M > 0 \Rightarrow \delta A > 0$

2. Physical process version of the 1st law

Consider a quasistatic process in which mass is added to the black hole. We thus start w/ a stationary black hole & alter it by some infinitesimal physical process. We assume the b.h. then settles down to a new stationary final state.

- $\Delta T_{\mu\nu}$ = stress energy of matter thrown in. The change in the mass & angular momentum of the b.h. are

$$\Delta M = \int_0^\infty d\lambda \int d^2\Omega \Delta T_{\mu\nu} \underbrace{\xi^\mu}_{\substack{\text{affine param} \\ \text{on } \mathcal{H}}} \underbrace{k^\nu}_{\substack{\perp \mathcal{H}}} \quad \Delta J = - \int_0^\infty d\lambda \int d^2\Omega \Delta T_{\mu\nu} \underbrace{\varphi^\mu}_{\substack{\perp \mathcal{H}}} \underbrace{k^\nu}_{\substack{\perp \mathcal{H}}}$$

- the change in the area of the horizon is $\Delta A = \int d^2\Omega d\lambda \theta(\lambda)$

- a Killing horizon has const. cross-sectional area, so $\theta=0$ (in fact, $B_{\mu\nu}=0$ for a Killing horizon. See e.g. Townsend)

- then, by Raychaudhuri's eqn, $\frac{d\theta}{d\lambda} = -8\pi G \Delta T_{\mu\nu} k^\mu k^\nu$ (drop θ^2 terms)

$$\begin{aligned} \Rightarrow \Delta A &= \int_0^\infty d\lambda \int_{S^2} d^2\Omega \theta = \int_{S^2} d^2\Omega \lambda \theta \Big|_0^\infty - \int_0^\infty d\lambda \int_{S^2} d^2\Omega \lambda \frac{d\theta}{d\lambda} = \\ &= 8\pi G \int_0^\infty d\lambda \int_{S^2} d^2\Omega \lambda \Delta T_{\mu\nu} k^\mu k^\nu = \frac{8\pi G}{k} (\Delta M - \Omega_H \Delta J) \end{aligned}$$

$\frac{1}{k} \xi^\mu$ since $\xi^\mu = \xi^\mu + \Omega_H \varphi^\mu$

$$\Rightarrow \frac{k \Delta A}{8\pi G} = \Delta M - \Omega_H \Delta J$$

- looks like "1st law" of thermodyn $T dS = dM + \dots \Rightarrow A_H \propto \text{entropy}$
 $k \propto \text{temperature}$

- same argument + the null energy condition, can be used to show that $\Delta\mathcal{A} > 0 \approx 2^{\text{nd}}$ law of thermodyn.

The third law : it is impossible, by any procedure, no matter how idealized, to reduce κ (surface gravity) to zero by a finite sequence of operations.

(clearly, cannot require that $S \rightarrow 0$ as $T \rightarrow 0$, violated by extremal b.h.s)

- very hard to bring to extremality by adding charge / spin.
- counterexample recently constr. by mathematicians (finely-tuned collapse $\Rightarrow$ exactly extremal Reissner - Nordström in finite time)